



Full wwPDB X-ray Structure Validation Report i

Dec 2, 2025 – 02:14 am GMT

PDB ID : 2C7R / pdb_00002c7r
Title : HhaI DNA methyltransferase (T250G mutant) complex with oligonucleotide containing 2-aminopurine as a target base (GPGC:GMGC) and SAH
Authors : Daujotyte, D.; Grazulis, S.
Deposited on : 2005-11-27
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	1337	-	-	X	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 3471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MODIFICATION METHYLASE HHAI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	327	Total	C 2612	N 1666	O 446	S 487	13	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	250	GLY	THR	engineered mutation	UNP P05102


- Molecule 2 is a DNA chain called 5'-D(*G*GP*AP*TP*GP*(5CM)*GP*CP*TP*GP*AP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	C	12	Total	C 251	N 118	O 48	P 73	12	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*G*TP*CP*AP*GP*(2PR)*GP*CP*AP*TP*CP*C)-3'.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	D	12	Total	C 246	N 116	O 46	P 72	12	0	0	0

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	26	14	6	5	1	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	229	Total O 229 229	0	0
7	C	21	Total O 21 21	0	0
7	D	21	Total O 21 21	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: MODIFICATION METHYLASE HHAI



- Molecule 2: 5'-D(*G*GP*AP*TP*GP*(5CM)*GP*CP*TP*GP*AP*C)-3'



- Molecule 3: 5'-D(*G*TP*CP*AP*GP*(2PR)*GP*CP*AP*TP*CP*C)-3'



4 Data and refinement statistics i

Xtriaze (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value			Source
Space group	H 3 2			Depositor
Cell constants a, b, c, α , β , γ	95.62 \AA 90.00 $^{\circ}$	95.62 \AA 90.00 $^{\circ}$	315.76 \AA 120.00 $^{\circ}$	Depositor
Resolution (\AA)	29.50 – 1.90			Depositor
% Data completeness (in resolution range)	99.8 (29.50-1.90)			Depositor
R_{merge}	0.07			Depositor
R_{sym}	(Not available)			Depositor
Refinement program	CNS 1.0			Depositor
R , R_{free}	0.199, 0.217			Depositor
Estimated twinning fraction	No twinning to report.			Xtriaze
Total number of atoms	3471			wwPDB-VP
Average B, all atoms (\AA^2)	18.0			wwPDB-VP

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5CM, GOL, SAH, SO4, 2PR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	2/2667 (0.1%)	1.00	10/3593 (0.3%)
2	C	0.40	0/258	0.78	0/393
3	D	0.40	0/250	0.99	1/379 (0.3%)
All	All	0.55	2/3175 (0.1%)	0.98	11/4365 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
3	D	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	GLU	C-N	-16.84	1.09	1.33
1	A	221	GLN	C-N	-12.75	1.16	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	GLU	CA-C-N	18.63	148.74	121.72
1	A	220	GLU	C-N-CA	18.63	148.74	121.72
1	A	220	GLU	O-C-N	-12.55	104.55	122.36
1	A	305	SER	N-CA-C	8.29	120.47	110.19
3	D	433	DC	C4'-C3'-O3'	7.68	121.52	110.00
1	A	146	SER	N-CA-C	-6.74	101.83	110.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ALA	N-CA-C	6.72	119.18	111.11
1	A	78	GLY	N-CA-C	-5.50	100.13	113.18
1	A	173	ASN	N-CA-C	5.42	117.89	111.33
1	A	77	ALA	N-CA-C	5.39	116.36	107.20
1	A	308	ILE	N-CA-C	5.24	115.86	110.36

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	409	DC	Sidechain
3	D	432	DC	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2612	0	2594	50	0
2	C	251	0	137	6	0
3	D	246	0	135	3	0
4	A	26	0	19	0	0
5	A	35	0	0	0	0
6	A	30	0	40	7	0
7	A	229	0	0	6	0
7	C	21	0	0	0	0
7	D	21	0	0	1	0
All	All	3471	0	2925	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:LYS:HE3	7:A:2066:HOH:O	1.21	1.28

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:412:DA:H2”	2:C:413:DC:H5’	1.29	1.07
1:A:50:GLU:OE2	1:A:56[A]:LYS:HD3	1.57	1.02
1:A:275:HIS:HD2	1:A:277:ARG:H	1.08	0.94
6:A:1340:GOL:H32	7:A:2227:HOH:O	1.68	0.92
1:A:82:GLN:H	1:A:82:GLN:HE21	1.19	0.85
1:A:50:GLU:OE2	1:A:56[A]:LYS:CD	2.26	0.83
1:A:275:HIS:CD2	1:A:277:ARG:H	2.00	0.70
1:A:111:LYS:HB3	7:A:2066:HOH:O	1.92	0.69
2:C:406:DG:H1’	2:C:407:5CM:H5”	1.77	0.66
1:A:203:GLU:OE2	6:A:1340:GOL:H11	1.96	0.66
1:A:138:ASN:O	1:A:142:GLU:HG3	1.97	0.64
1:A:5:LYS:N	6:A:1337:GOL:O3	2.29	0.63
1:A:56[B]:LYS:HE2	7:A:2034:HOH:O	1.98	0.63
1:A:127:HIS:HD2	1:A:132:THR:OG1	1.80	0.62
2:C:412:DA:C2’	2:C:413:DC:H5’	2.18	0.62
1:A:156:ASP:HB3	1:A:185:PRO:HB3	1.81	0.62
1:A:43:LYS:O	1:A:47:GLU:HG3	2.01	0.61
2:C:409:DC:H2”	2:C:410:DT:H71	1.83	0.60
1:A:42:ASP:O	1:A:46:GLN:HG3	2.01	0.60
1:A:256:GLY:HA3	2:C:406:DG:O6	2.02	0.59
1:A:5:LYS:H	6:A:1337:GOL:HO3	1.53	0.56
1:A:65:ASN:HB3	1:A:68:THR:HG23	1.87	0.55
1:A:82:GLN:HE21	1:A:82:GLN:N	1.99	0.53
1:A:80:PRO:HB3	1:A:82:GLN:HE22	1.75	0.52
1:A:152:LEU:HD22	6:A:1336:GOL:H11	1.93	0.51
2:C:412:DA:H2”	2:C:413:DC:C5’	2.20	0.51
1:A:5:LYS:HG3	6:A:1337:GOL:O3	2.10	0.50
1:A:155:LEU:HD12	1:A:155:LEU:C	2.37	0.50
1:A:90:GLN:O	1:A:127:HIS:HE1	1.95	0.50
1:A:5:LYS:N	6:A:1337:GOL:HO3	2.09	0.49
1:A:137:LYS:NZ	1:A:141:ASN:HD21	2.10	0.49
1:A:262:THR:OG1	1:A:273:LYS:HB2	2.12	0.49
1:A:25:ARG:HH12	1:A:55:GLU:CD	2.21	0.48
1:A:111:LYS:CB	7:A:2066:HOH:O	2.57	0.46
1:A:137:LYS:HZ3	1:A:141:ASN:HD21	1.63	0.46
1:A:59:GLY:O	1:A:60:ASP:C	2.58	0.46
3:D:427:2PR:C8	3:D:427:2PR:H5”	2.47	0.45
3:D:427:2PR:H8	3:D:427:2PR:H2”	1.77	0.45
1:A:225:LYS:C	1:A:247:ILE:HG12	2.42	0.45
3:D:427:2PR:H2”	7:D:2010:HOH:O	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:ILE:HG13	1:A:180:PHE:CZ	2.53	0.44
1:A:25:ARG:O	1:A:29:GLU:HG3	2.17	0.44
1:A:75:LEU:C	1:A:75:LEU:HD23	2.43	0.43
1:A:101:PHE:CE1	1:A:136:VAL:HG13	2.54	0.43
1:A:125:ALA:HB2	1:A:166:ILE:HD12	1.99	0.43
1:A:297:GLN:O	1:A:301:GLN:HG3	2.19	0.43
1:A:209:ARG:HB3	1:A:211:ASP:OD1	2.19	0.42
1:A:127:HIS:CD2	1:A:132:THR:OG1	2.68	0.41
1:A:84:PHE:CE2	1:A:92:GLY:HA2	2.55	0.41
1:A:221:GLN:HG3	7:A:2134:HOH:O	2.21	0.41
1:A:202:VAL:HB	1:A:205:LEU:HD12	2.03	0.41
1:A:260:ALA:C	1:A:261:LYS:HG2	2.46	0.41
1:A:9:LEU:CD1	1:A:28:LEU:HD13	2.52	0.40
1:A:70:PRO:O	1:A:72:HIS:HD2	2.04	0.40
1:A:79:PHE:HB2	1:A:80:PRO:HD2	2.03	0.40
1:A:157:TYR:CE2	1:A:183:PRO:HB2	2.56	0.40
1:A:17:LEU:O	1:A:18:PHE:C	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	326/327 (100%)	316 (97%)	9 (3%)	1 (0%)	37 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	283/282 (100%)	278 (98%)	5 (2%)	54 52

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	56[A]	LYS
1	A	56[B]	LYS
1	A	82	GLN
1	A	232	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
1	A	82	GLN
1	A	90	GLN
1	A	127	HIS
1	A	129	ASN
1	A	141	ASN
1	A	181	GLN
1	A	268	ASN
1	A	275	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2PR	D	427	3	17,23,24	1.11	2 (11%)	16,33,36	1.74	3 (18%)
2	5CM	C	407	2,3	17,21,22	0.74	1 (5%)	24,30,33	1.63	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2PR	D	427	3	-	1/3/21/22	0/3/3/3
2	5CM	C	407	2,3	-	0/7/21/22	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	427	2PR	C6-N1	3.39	1.39	1.32
2	C	407	5CM	C4-N3	2.48	1.38	1.34
3	D	427	2PR	C8-N7	-2.20	1.30	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	407	5CM	C5-C4-N3	-5.22	116.04	121.67
3	D	427	2PR	C6-N1-C2	4.81	125.33	116.43
2	C	407	5CM	C4-N3-C2	4.50	126.77	120.69
3	D	427	2PR	C2-N3-C4	-3.02	111.91	115.36
2	C	407	5CM	N1-C2-N3	-2.34	114.54	118.81
3	D	427	2PR	C2'-C1'-N9	-2.31	108.95	114.27
2	C	407	5CM	O2-C2-N1	2.00	123.03	118.89

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	427	2PR	C4'-C5'-O5'-P

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	427	2PR	3	0
2	C	407	5CM	1	0

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	1338	-	5,5,5	0.93	0	5,5,5	0.41	0
5	SO4	A	1330	-	4,4,4	1.88	2 (50%)	6,6,6	0.92	0
6	GOL	A	1336	-	5,5,5	0.92	0	5,5,5	0.40	0
5	SO4	A	1331	-	4,4,4	1.90	2 (50%)	6,6,6	0.87	0
4	SAH	A	1328	-	24,28,28	1.27	4 (16%)	25,40,40	2.41	4 (16%)
5	SO4	A	1329	-	4,4,4	1.82	2 (50%)	6,6,6	0.89	0
5	SO4	A	1333	-	4,4,4	1.88	2 (50%)	6,6,6	0.89	0
6	GOL	A	1339	-	5,5,5	0.92	0	5,5,5	0.40	0
5	SO4	A	1335	-	4,4,4	1.93	2 (50%)	6,6,6	0.97	0
6	GOL	A	1340	-	5,5,5	0.92	0	5,5,5	0.40	0
5	SO4	A	1332	-	4,4,4	1.87	2 (50%)	6,6,6	0.90	0
6	GOL	A	1337	-	5,5,5	0.90	0	5,5,5	0.40	0
5	SO4	A	1334	-	4,4,4	1.86	2 (50%)	6,6,6	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	1336	-	-	4/4/4/4	-
4	SAH	A	1328	-	-	3/11/31/31	0/3/3/3
6	GOL	A	1340	-	-	4/4/4/4	-
6	GOL	A	1339	-	-	2/4/4/4	-
6	GOL	A	1337	-	-	2/4/4/4	-
6	GOL	A	1338	-	-	2/4/4/4	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1328	SAH	C4-N3	3.26	1.40	1.35
5	A	1331	SO4	O1-S	3.16	1.63	1.46
5	A	1335	SO4	O1-S	3.16	1.63	1.46
5	A	1330	SO4	O1-S	3.12	1.62	1.46
5	A	1333	SO4	O1-S	3.10	1.62	1.46
5	A	1332	SO4	O1-S	3.09	1.62	1.46
5	A	1334	SO4	O1-S	3.08	1.62	1.46
5	A	1329	SO4	O1-S	2.96	1.62	1.46
4	A	1328	SAH	C8-N7	-2.52	1.30	1.34
4	A	1328	SAH	C2'-C1'	-2.50	1.50	1.53
5	A	1335	SO4	O3-S	-2.19	1.29	1.47
5	A	1333	SO4	O3-S	-2.09	1.30	1.47
5	A	1329	SO4	O3-S	-2.08	1.30	1.47
5	A	1331	SO4	O3-S	-2.07	1.30	1.47
5	A	1332	SO4	O3-S	-2.07	1.30	1.47
4	A	1328	SAH	O-C	2.06	1.28	1.22
5	A	1330	SO4	O3-S	-2.06	1.31	1.47
5	A	1334	SO4	O3-S	-2.05	1.31	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1328	SAH	OXT-C-O	-9.40	102.76	124.09
4	A	1328	SAH	O4'-C1'-C2'	-5.11	99.45	106.93
4	A	1328	SAH	C5-C6-N6	2.96	124.85	120.35
4	A	1328	SAH	CB-CA-N	2.85	117.63	110.17

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1336	GOL	C1-C2-C3-O3
6	A	1337	GOL	O1-C1-C2-C3
6	A	1338	GOL	O1-C1-C2-C3
6	A	1340	GOL	C1-C2-C3-O3
6	A	1340	GOL	O2-C2-C3-O3
4	A	1328	SAH	OXT-C-CA-CB
6	A	1338	GOL	O1-C1-C2-O2
6	A	1339	GOL	O1-C1-C2-C3
6	A	1340	GOL	O1-C1-C2-C3
6	A	1336	GOL	O2-C2-C3-O3
6	A	1337	GOL	O1-C1-C2-O2
6	A	1339	GOL	O1-C1-C2-O2
6	A	1340	GOL	O1-C1-C2-O2
4	A	1328	SAH	OXT-C-CA-N
6	A	1336	GOL	O1-C1-C2-O2
6	A	1336	GOL	O1-C1-C2-C3
4	A	1328	SAH	CB-CG-SD-C5'

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1336	GOL	1	0
6	A	1340	GOL	2	0
6	A	1337	GOL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	221:GLN	C	222:THR	N	1.16
1	A	220:GLU	C	221:GLN	N	1.09

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

EDS was not executed - this section is therefore empty.