



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 08:31 AM EDT

PDB ID : 2CDV
Title : REFINED STRUCTURE OF CYTOCHROME C3 AT 1.8 ANGSTROMS RESOLUTION
Authors : Higuchi, Y.; Kusunoki, M.; Matsuura, Y.; Yasuoka, N.; Kakudo, M.
Deposited on : 1983-11-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	107	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	107	801	489	151	150	11	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ASN	ASP	conflict	UNP P00132

- Molecule 2 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	47	Total	O	0	0
			47	47		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: CYTOCHROME C3

Chain A:  77% 23%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.90Å 68.10Å 34.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	0.176 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1020	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.97	0/818	1.49	5/1089 (0.5%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	12	ASP	CB-CG-OD1	14.44	131.30	118.30
1	A	59	ASP	CB-CG-OD2	7.27	124.84	118.30
1	A	53	ASP	CB-CG-OD2	-7.26	111.77	118.30
1	A	53	ASP	CB-CG-OD1	6.43	124.09	118.30
1	A	12	ASP	CB-CG-OD2	-6.38	112.56	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	782	16	0
2	A	172	0	120	7	0
3	A	47	0	0	2	0
All	All	1020	0	902	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:111:HEC:HBC3	2:A:111:HEC:HMC1	1.59	0.85
1:A:96:GLU:HG2	1:A:104:LYS:HD2	1.67	0.76
2:A:111:HEC:HMC1	2:A:111:HEC:CBC	2.28	0.62
1:A:90:ASP:O	1:A:94:LYS:HG3	2.02	0.59
1:A:16:GLN:HA	1:A:16:GLN:OE1	2.07	0.54
1:A:5:PRO:HD2	1:A:22:HIS:CE1	2.47	0.49
1:A:67:HIS:NE2	2:A:109:HEC:O2A	2.45	0.49
1:A:83:HIS:HE1	2:A:111:HEC:C4A	2.27	0.47
1:A:50:GLY:HA2	1:A:63:LYS:HZ1	1.79	0.46
1:A:15:LYS:C	1:A:17:PRO:HD3	2.36	0.46
1:A:52:HIS:ND1	1:A:62:ALA:O	2.51	0.43
1:A:66:TYR:HA	3:A:156:HOH:O	2.18	0.42
1:A:88:GLY:O	1:A:94:LYS:HE2	2.19	0.42
1:A:104:LYS:HB3	2:A:111:HEC:HMA3	2.02	0.42
1:A:4:ALA:HA	1:A:5:PRO:HD3	1.81	0.42
1:A:5:PRO:HD3	2:A:110:HEC:C4A	2.50	0.41
2:A:111:HEC:HMB1	2:A:111:HEC:HBB3	2.02	0.40
1:A:1:ALA:HA	1:A:2:PRO:HD2	1.93	0.40
1:A:94:LYS:HE2	3:A:128:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	105/107 (98%)	102 (97%)	3 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	85/85 (100%)	83 (98%)	2 (2%)	49 36

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LYS
1	A	95	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	109	1	32,50,50	2.19	9 (28%)	30,82,82	2.77	13 (43%)
2	HEC	A	111	1	32,50,50	2.21	6 (18%)	30,82,82	2.40	12 (40%)
2	HEC	A	108	1	32,50,50	1.95	6 (18%)	30,82,82	1.97	9 (30%)
2	HEC	A	110	1	32,50,50	2.06	7 (21%)	30,82,82	2.55	12 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	109	1	-	0/10/54/54	-
2	HEC	A	111	1	-	4/10/54/54	-
2	HEC	A	108	1	-	5/10/54/54	-
2	HEC	A	110	1	-	2/10/54/54	-

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	109	HEC	C3C-C2C	-7.15	1.32	1.40
2	A	111	HEC	C2B-C3B	-6.98	1.32	1.40
2	A	111	HEC	C3C-C2C	-6.53	1.33	1.40
2	A	108	HEC	C2B-C3B	-5.88	1.34	1.40
2	A	110	HEC	C2B-C3B	-5.61	1.34	1.40
2	A	110	HEC	C3C-C2C	-5.57	1.34	1.40
2	A	109	HEC	C2B-C3B	-4.96	1.35	1.40
2	A	108	HEC	C3C-C2C	-4.33	1.35	1.40
2	A	109	HEC	CBC-CAC	-3.63	1.36	1.49
2	A	108	HEC	CBB-CAB	-3.53	1.36	1.49
2	A	110	HEC	CBB-CAB	-3.43	1.36	1.49
2	A	110	HEC	CBC-CAC	-3.40	1.36	1.49
2	A	111	HEC	CBB-CAB	-3.26	1.37	1.49
2	A	108	HEC	CBC-CAC	-3.15	1.37	1.49
2	A	111	HEC	CBC-CAC	-3.08	1.38	1.49
2	A	109	HEC	CBB-CAB	-2.86	1.38	1.49
2	A	109	HEC	C3A-C4A	2.49	1.48	1.42
2	A	110	HEC	CMD-C2D	2.47	1.56	1.51
2	A	108	HEC	CAA-C2A	2.37	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	110	HEC	CMC-C2C	2.27	1.57	1.51
2	A	109	HEC	CMD-C2D	2.25	1.56	1.51
2	A	109	HEC	CAD-C3D	2.19	1.57	1.52
2	A	110	HEC	O2D-CGD	-2.18	1.23	1.30
2	A	109	HEC	C3D-C2D	-2.18	1.31	1.37
2	A	111	HEC	C4B-C3B	2.13	1.46	1.43
2	A	109	HEC	CAA-C2A	2.05	1.55	1.52
2	A	111	HEC	C3D-C2D	-2.04	1.31	1.37
2	A	108	HEC	O2D-CGD	-2.02	1.24	1.30

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	108	HEC	CBB-CAB-C3B	-5.95	113.57	127.49
2	A	111	HEC	CBC-CAC-C3C	-5.83	113.85	127.49
2	A	109	HEC	CMC-C2C-C3C	5.75	132.58	125.82
2	A	109	HEC	CMC-C2C-C1C	-5.67	120.15	128.46
2	A	109	HEC	CBC-CAC-C3C	-5.14	115.46	127.49
2	A	110	HEC	CMC-C2C-C1C	-5.11	120.96	128.46
2	A	111	HEC	CBB-CAB-C3B	-4.98	115.84	127.49
2	A	109	HEC	CBD-CAD-C3D	-4.90	104.31	112.54
2	A	110	HEC	CBC-CAC-C3C	-4.85	116.15	127.49
2	A	110	HEC	CMD-C2D-C1D	-4.75	121.49	128.46
2	A	109	HEC	CBB-CAB-C3B	-4.72	116.44	127.49
2	A	109	HEC	CMB-C2B-C1B	-4.59	121.73	128.46
2	A	110	HEC	CMB-C2B-C1B	-4.49	121.88	128.46
2	A	110	HEC	CMC-C2C-C3C	4.46	131.07	125.82
2	A	110	HEC	CMB-C2B-C3B	4.15	130.70	125.82
2	A	109	HEC	CMD-C2D-C1D	-4.00	122.60	128.46
2	A	111	HEC	CMD-C2D-C1D	-3.83	122.85	128.46
2	A	110	HEC	CBB-CAB-C3B	-3.78	118.65	127.49
2	A	111	HEC	CMC-C2C-C1C	-3.72	123.00	128.46
2	A	108	HEC	CMC-C2C-C1C	-3.46	123.39	128.46
2	A	111	HEC	CMB-C2B-C1B	-3.38	123.50	128.46
2	A	111	HEC	CMC-C2C-C3C	3.35	129.76	125.82
2	A	108	HEC	CMB-C2B-C3B	3.14	129.51	125.82
2	A	111	HEC	O1A-CGA-CBA	-3.03	113.48	123.09
2	A	108	HEC	CMB-C2B-C1B	-3.02	124.03	128.46
2	A	109	HEC	CMA-C3A-C2A	2.95	130.51	124.94
2	A	109	HEC	CMB-C2B-C3B	2.92	129.25	125.82
2	A	110	HEC	O1D-CGD-CBD	-2.91	113.87	123.09
2	A	108	HEC	CBC-CAC-C3C	-2.89	120.73	127.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	111	HEC	O2D-CGD-CBD	2.77	122.76	114.00
2	A	108	HEC	CMA-C3A-C2A	2.72	130.08	124.94
2	A	108	HEC	CMD-C2D-C1D	-2.67	124.55	128.46
2	A	111	HEC	CMB-C2B-C3B	2.61	128.89	125.82
2	A	109	HEC	O1A-CGA-CBA	-2.60	114.83	123.09
2	A	110	HEC	CMA-C3A-C2A	2.60	129.84	124.94
2	A	110	HEC	CMD-C2D-C3D	2.48	129.63	124.94
2	A	111	HEC	O1D-CGD-CBD	-2.44	115.36	123.09
2	A	109	HEC	CMD-C2D-C3D	2.43	129.52	124.94
2	A	110	HEC	C1D-C2D-C3D	2.40	108.66	107.00
2	A	110	HEC	O2D-CGD-CBD	2.34	121.39	114.00
2	A	111	HEC	CMD-C2D-C3D	2.33	129.33	124.94
2	A	109	HEC	O2D-CGD-O1D	2.28	129.20	123.33
2	A	108	HEC	O1A-CGA-CBA	-2.09	116.47	123.09
2	A	111	HEC	CBD-CAD-C3D	-2.05	109.09	112.54
2	A	109	HEC	O1D-CGD-CBD	-2.03	116.66	123.09
2	A	108	HEC	O1D-CGD-CBD	-2.03	116.67	123.09

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	111	HEC	CAA-CBA-CGA-O2A
2	A	111	HEC	CAA-CBA-CGA-O1A
2	A	108	HEC	CAD-CBD-CGD-O1D
2	A	108	HEC	CAD-CBD-CGD-O2D
2	A	111	HEC	CAD-CBD-CGD-O2D
2	A	108	HEC	CAA-CBA-CGA-O2A
2	A	108	HEC	C3A-C2A-CAA-CBA
2	A	108	HEC	CAA-CBA-CGA-O1A
2	A	111	HEC	CAD-CBD-CGD-O1D
2	A	110	HEC	CAA-CBA-CGA-O2A
2	A	110	HEC	CAA-CBA-CGA-O1A


There are no ring outliers.


3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	109	HEC	1	0
2	A	111	HEC	5	0
2	A	110	HEC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.