



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 02:59 pm GMT

PDB ID : 2J5E  
Title : Crystal structure of EGFR kinase domain in complex with an irreversible inhibitor 13-jab  
Authors : Yun, C.-H.; Eck, M.J.  
Deposited on : 2006-09-14  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPIDERMAL GROWTH FACTOR RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	310	2483	1599	416	452	16	0	0	0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	60	Total	O	0	0
			60	60		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: EPIDERMAL GROWTH FACTOR RECEPTOR



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.02Å 145.02Å 145.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 3.10 24.87 – 3.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.00-3.10) 100.0 (24.87-3.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.60 (at 3.11Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.186 , 0.255 0.186 , 0.257	Depositor DCC
$R_{free}$ test set	658 reflections (7.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.6	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.036 for -l,-k,-h	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2544	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CY0, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	5/2506 (0.2%)	0.77	1/3391 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	734	GLU	CB-CG	7.20	1.65	1.52
1	A	734	GLU	CD-OE2	6.50	1.32	1.25
1	A	745	LYS	CE-NZ	6.37	1.65	1.49
1	A	734	GLU	CD-OE1	5.65	1.31	1.25
1	A	734	GLU	CG-CD	5.12	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1018	ILE	C-N-CD	-5.30	108.94	120.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2483	0	2501	114	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	60	0	0	3	0
All	All	2544	0	2501	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:724:GLY:HA2	1:A:748:ARG:H	1.10	1.10
1:A:835:HIS:HD2	1:A:837:ASP:H	1.11	0.94
1:A:857:GLY:C	1:A:858:LEU:HD23	1.90	0.91
1:A:854:THR:HG23	1:A:855:ASP:N	1.83	0.91
1:A:858:LEU:HD23	1:A:858:LEU:N	1.91	0.85
1:A:1019:PRO:O	1:A:1020:GLN:CB	2.30	0.80
1:A:846:LYS:HE3	1:A:1014:ASP:OD1	1.81	0.80
1:A:696:GLY:N	1:A:863:GLY:HA2	1.96	0.80
1:A:1018:ILE:O	1:A:1018:ILE:HG23	1.83	0.78
1:A:724:GLY:HA2	1:A:748:ARG:N	1.95	0.77
1:A:768:SER:O	1:A:831:ARG:NH1	2.19	0.75
1:A:825:MET:HE3	1:A:853:ILE:HD13	1.69	0.75
1:A:724:GLY:CA	1:A:748:ARG:H	1.98	0.73
1:A:747:LEU:HD12	1:A:748:ARG:O	1.89	0.73
1:A:1007:MET:HA	3:A:2058:HOH:O	1.91	0.70
1:A:783:THR:O	1:A:784:SER:C	2.30	0.68
1:A:793:MET:HE1	1:A:852:LYS:HD2	1.76	0.67
1:A:918:ILE:H	1:A:918:ILE:HD12	1.58	0.67
1:A:926:ILE:HG23	1:A:931:GLU:HB3	1.76	0.66
1:A:835:HIS:HD2	1:A:837:ASP:N	1.90	0.66
1:A:835:HIS:CD2	1:A:837:ASP:H	2.04	0.66
1:A:820:GLN:HE21	1:A:851:VAL:HG22	1.63	0.63
1:A:756:ASN:ND2	1:A:786:VAL:HG23	2.13	0.63
1:A:718:LEU:HD13	1:A:728:LYS:HB2	1.80	0.63
1:A:877:PRO:O	1:A:881:MET:HG2	1.99	0.62
1:A:767:ALA:HB2	1:A:777:LEU:HD23	1.81	0.62
1:A:882:ALA:HB1	1:A:884:GLU:OE2	2.01	0.61
1:A:932:ARG:HH22	1:A:953:ILE:HD11	1.66	0.61
1:A:965:ILE:O	1:A:969:SER:HB2	2.01	0.60
1:A:858:LEU:N	1:A:858:LEU:CD2	2.64	0.60
1:A:1018:ILE:HG13	1:A:1019:PRO:HD2	1.85	0.59
1:A:820:GLN:NE2	1:A:851:VAL:H	2.01	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:823:LYS:HA	1:A:965:ILE:HD11	1.84	0.58
1:A:834:VAL:HG12	1:A:836:ARG:HG3	1.85	0.58
1:A:802:VAL:HG12	1:A:809:ILE:HD13	1.83	0.58
1:A:985:GLU:O	1:A:987:MET:N	2.38	0.57
1:A:1018:ILE:O	1:A:1018:ILE:CG2	2.52	0.56
1:A:797:CY0:CAJ	3:A:2037:HOH:O	2.53	0.56
1:A:751:THR:O	1:A:753:PRO:HD3	2.06	0.55
1:A:879:LYS:HE3	1:A:914:PRO:O	2.06	0.55
1:A:854:THR:CG2	1:A:855:ASP:N	2.56	0.55
1:A:825:MET:CE	1:A:853:ILE:HD13	2.36	0.55
1:A:708:LYS:HB2	1:A:711:GLU:HG3	1.89	0.54
1:A:854:THR:HG23	1:A:855:ASP:H	1.68	0.54
1:A:908:MET:HG3	1:A:939:CYS:SG	2.47	0.54
1:A:836:ARG:HG2	1:A:891:TYR:CD1	2.43	0.54
1:A:1017:LEU:O	1:A:1018:ILE:HB	2.07	0.54
1:A:748:ARG:O	1:A:749:GLU:HB2	2.07	0.53
1:A:886:ILE:O	1:A:886:ILE:HG22	2.08	0.53
1:A:700:ASN:C	1:A:700:ASN:HD22	2.10	0.53
1:A:756:ASN:ND2	1:A:786:VAL:CG2	2.70	0.53
1:A:793:MET:CE	1:A:852:LYS:HD2	2.38	0.53
1:A:795:PHE:HB2	1:A:845:VAL:HB	1.91	0.53
1:A:858:LEU:HD12	1:A:876:VAL:HG13	1.94	0.50
1:A:756:ASN:HA	1:A:759:ILE:HG22	1.93	0.50
1:A:1018:ILE:O	1:A:1019:PRO:O	2.30	0.50
1:A:867:LYS:HD3	1:A:867:LYS:O	2.10	0.50
1:A:713:LYS:HG2	1:A:732:ILE:HD12	1.94	0.49
1:A:781:CYS:HB3	1:A:787:GLN:HE21	1.77	0.49
1:A:820:GLN:HE22	1:A:851:VAL:H	1.61	0.49
1:A:836:ARG:NH1	1:A:858:LEU:HB3	2.28	0.48
1:A:791:GLN:HG3	3:A:2014:HOH:O	2.12	0.48
1:A:791:GLN:OE1	1:A:1012:ASP:OD1	2.31	0.48
1:A:909:THR:HB	1:A:912:SER:OG	2.14	0.48
1:A:955:ALA:HA	1:A:958:ARG:NE	2.29	0.48
1:A:724:GLY:HA3	1:A:747:LEU:HA	1.95	0.47
1:A:815:LEU:O	1:A:819:VAL:HG23	2.14	0.47
1:A:955:ALA:O	1:A:958:ARG:HB2	2.14	0.47
1:A:972:ALA:O	1:A:975:PRO:HD3	2.13	0.47
1:A:909:THR:HG22	1:A:936:PRO:HB3	1.96	0.47
1:A:715:ILE:O	1:A:716:LYS:HG3	2.14	0.47
1:A:936:PRO:O	1:A:939:CYS:HB2	2.14	0.47
1:A:808:ASN:ND2	1:A:988:HIS:O	2.47	0.47

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:897:VAL:O	1:A:900:TYR:HB3	2.15	0.46
1:A:898:TRP:CE3	1:A:958:ARG:NH1	2.84	0.46
1:A:759:ILE:HG13	1:A:788:LEU:HD13	1.98	0.46
1:A:804:GLU:O	1:A:805:HIS:ND1	2.49	0.46
1:A:811:SER:HB2	1:A:979:LEU:HB2	1.97	0.45
1:A:825:MET:HE3	1:A:853:ILE:CD1	2.40	0.45
1:A:898:TRP:HE3	1:A:958:ARG:NH1	2.14	0.45
1:A:700:ASN:C	1:A:700:ASN:ND2	2.69	0.45
1:A:825:MET:CE	1:A:853:ILE:CD1	2.94	0.45
1:A:778:LEU:HB2	1:A:789:ILE:O	2.16	0.45
1:A:932:ARG:NH1	1:A:951:TRP:O	2.50	0.45
1:A:774:VAL:HG13	1:A:856:PHE:HZ	1.82	0.45
1:A:952:MET:O	1:A:958:ARG:NH1	2.49	0.45
1:A:783:THR:O	1:A:785:THR:N	2.51	0.44
1:A:846:LYS:HE2	1:A:1012:ASP:OD2	2.17	0.44
1:A:752:SER:O	1:A:754:LYS:N	2.51	0.44
1:A:811:SER:OG	1:A:975:PRO:HB2	2.17	0.44
1:A:835:HIS:HE1	1:A:854:THR:O	2.01	0.44
1:A:796:GLY:HA2	1:A:797:CY0:HAK	2.00	0.44
1:A:932:ARG:HH22	1:A:953:ILE:CD1	2.30	0.44
1:A:820:GLN:NE2	1:A:851:VAL:HG22	2.32	0.43
1:A:926:ILE:CG2	1:A:931:GLU:HB3	2.45	0.43
1:A:793:MET:HG3	1:A:844:LEU:CB	2.48	0.43
1:A:908:MET:HG2	1:A:947:MET:HE1	1.99	0.43
1:A:977:ARG:O	1:A:977:ARG:HG2	2.18	0.43
1:A:813:TYR:CD1	1:A:813:TYR:N	2.85	0.43
1:A:984:ASP:O	1:A:985:GLU:CD	2.57	0.42
1:A:736:GLU:HB3	1:A:738:VAL:HG22	2.00	0.42
1:A:902:VAL:O	1:A:906:GLU:HG3	2.19	0.42
1:A:705:ARG:HG2	1:A:707:LEU:HD23	2.01	0.42
1:A:793:MET:HG3	1:A:844:LEU:HB2	2.01	0.42
1:A:983:GLY:O	1:A:984:ASP:O	2.38	0.42
1:A:829:GLU:HG2	1:A:893:HIS:CG	2.55	0.42
1:A:918:ILE:HG12	1:A:926:ILE:HD11	2.02	0.41
1:A:861:LEU:H	1:A:861:LEU:HD12	1.86	0.41
1:A:748:ARG:O	1:A:749:GLU:CB	2.69	0.41
1:A:797:CY0:OAC	1:A:797:CY0:HAM	2.20	0.41
1:A:730:LEU:HD22	1:A:739:LYS:HB3	2.04	0.40
1:A:740:ILE:O	1:A:742:VAL:HG13	2.21	0.40
1:A:759:ILE:HG13	1:A:788:LEU:CD1	2.51	0.40
1:A:853:ILE:HG22	1:A:854:THR:N	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	303/327 (93%)	253 (84%)	36 (12%)	14 (5%)	<b>2</b> <b>15</b>

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	749	GLU
1	A	1019	PRO
1	A	697	GLU
1	A	756	ASN
1	A	784	SER
1	A	872	GLU
1	A	984	ASP
1	A	720	SER
1	A	750	ALA
1	A	753	PRO
1	A	752	SER
1	A	808	ASN
1	A	1018	ILE
1	A	937	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	267/286 (93%)	231 (86%)	36 (14%)	4 16

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	700	ASN
1	A	705	ARG
1	A	747	LEU
1	A	751	THR
1	A	757	LYS
1	A	784	SER
1	A	785	THR
1	A	786	VAL
1	A	807	ASP
1	A	808	ASN
1	A	831	ARG
1	A	832	ARG
1	A	833	LEU
1	A	841	ARG
1	A	847	THR
1	A	854	THR
1	A	855	ASP
1	A	858	LEU
1	A	867	LYS
1	A	875	LYS
1	A	878	ILE
1	A	884	GLU
1	A	892	THR
1	A	895	SER
1	A	908	MET
1	A	912	SER
1	A	913	LYS
1	A	925	SER
1	A	945	MET
1	A	960	LYS
1	A	967	GLU
1	A	985	GLU
1	A	1008	ASP
1	A	1010	VAL
1	A	1011	VAL
1	A	1017	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	700	ASN
1	A	756	ASN
1	A	787	GLN
1	A	812	GLN
1	A	816	ASN
1	A	820	GLN
1	A	835	HIS
1	A	842	ASN
1	A	893	HIS
1	A	935	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CY0	A	797	1	29,30,31	1.32	3 (10%)	34,39,41	2.28	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CY0	A	797	1	-	3/15/17/19	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	797	CY0	C6-C5	-2.99	1.41	1.44
1	A	797	CY0	CAY-NAS	-2.71	1.36	1.41
1	A	797	CY0	CAX-NAT	-2.65	1.34	1.40

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	797	CY0	C2-N1-C6	5.51	121.31	116.59
1	A	797	CY0	C6-C5-C4	5.46	119.32	115.88
1	A	797	CY0	CAM-C5-C6	-4.82	120.53	124.88
1	A	797	CY0	N3-C2-N1	-4.46	121.71	128.68
1	A	797	CY0	C5-C6-N1	-4.20	117.89	121.35
1	A	797	CY0	C2-N3-C4	3.91	120.79	115.40
1	A	797	CY0	C5-C4-N3	-3.63	118.97	122.83
1	A	797	CY0	NAT-C6-N1	3.09	122.86	118.72

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	797	CY0	SAU-CAN-CAO-CAW
1	A	797	CY0	CAN-CAO-CAW-NAS
1	A	797	CY0	CAN-CAO-CAW-OAC

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	797	CY0	3	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/327 (94%)	-0.30	8 (2%) 56 33	29, 51, 86, 112	5 (1%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	696	GLY	3.1
1	A	750	ALA	3.1
1	A	990	PRO	2.8
1	A	749	GLU	2.7
1	A	1019	PRO	2.3
1	A	987	MET	2.1
1	A	1005	GLU	2.1
1	A	922	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CY0	A	797	28/29	0.92	0.18	52,66,69,72	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	3021	1/1	0.97	0.12	52,52,52,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.