



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2022 – 12:06 PM EST

PDB ID : 2KBZ
Title : NMR structure of protein gp15 of bacteriophage SPP1
Authors : Gallopin, M.; Gilquin, B.; Zinn-Justin, S.
Deposited on : 2008-12-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.27
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	99	

2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:42-A:102, A:116-A:133 (79)	0.75	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 4, 5, 6, 8, 9, 11, 12, 14, 15, 17, 19
2	10, 16, 18, 20
3	1, 7
Single-model clusters	13

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1586 atoms, of which 797 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 15 protein (Bacteriophage SPP1 complete nucleotide sequence).


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	99	1586	502	797	134	148	5	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)



4.2.2 Score per residue for model 2 (medoid)


- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)






4.2.3 Score per residue for model 3

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.4 Score per residue for model 4

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.5 Score per residue for model 5

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.6 Score per residue for model 6

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.7 Score per residue for model 7

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.8 Score per residue for model 8

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)



4.2.9 Score per residue for model 9


- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)



4.2.10 Score per residue for model 10


- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)






4.2.11 Score per residue for model 11

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.12 Score per residue for model 12

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.13 Score per residue for model 13

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.14 Score per residue for model 14

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.15 Score per residue for model 15

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.16 Score per residue for model 16

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.17 Score per residue for model 17

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.18 Score per residue for model 18

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)




4.2.19 Score per residue for model 19

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)



4.2.20 Score per residue for model 20

- Molecule 1: 15 protein (Bacteriophage SPP1 complete nucleotide sequence)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
INCA	refinement	1.0

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.98±0.02	0±0/651 (0.0± 0.0%)	1.64±0.04	10±3/883 (1.1± 0.3%)
All	All	0.98	0/13020 (0.0%)	1.64	192/17660 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.2±0.9
All	All	0	44

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	98	PHE	CB-CG-CD1	9.90	127.73	120.80	3	7
1	A	132	TYR	CB-CG-CD1	9.46	126.67	121.00	20	9
1	A	99	TYR	CB-CG-CD1	-8.78	115.73	121.00	14	5
1	A	60	VAL	CA-C-N	8.53	140.97	117.10	18	18
1	A	98	PHE	CB-CG-CD2	-8.17	115.08	120.80	11	8
1	A	59	MET	CG-SD-CE	-7.84	87.66	100.20	19	1
1	A	91	PHE	CB-CG-CD1	7.78	126.24	120.80	2	2
1	A	129	LEU	CB-CG-CD2	7.75	124.18	111.00	1	2
1	A	132	TYR	CB-CG-CD2	-7.48	116.51	121.00	10	8
1	A	99	TYR	CB-CG-CD2	-7.29	116.63	121.00	17	3
1	A	133	ARG	C-N-CA	7.29	139.92	121.70	5	1
1	A	100	MET	CG-SD-CE	-7.02	88.97	100.20	15	2
1	A	73	ASN	N-CA-CB	-6.95	98.10	110.60	8	2
1	A	75	PHE	CB-CG-CD1	6.90	125.63	120.80	3	1
1	A	117	PHE	CB-CG-CD2	-6.86	116.00	120.80	8	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	116	ASN	C-N-CA	6.84	138.81	121.70	20	1
1	A	55	TYR	CB-CG-CD1	6.81	125.08	121.00	5	4
1	A	125	ILE	CA-CB-CG1	6.75	123.83	111.00	6	2
1	A	96	ALA	CB-CA-C	6.74	120.22	110.10	15	9
1	A	129	LEU	CA-CB-CG	6.59	130.47	115.30	1	1
1	A	132	TYR	CA-CB-CG	6.53	125.80	113.40	19	1
1	A	65	GLU	CB-CA-C	6.47	123.33	110.40	2	4
1	A	91	PHE	CB-CG-CD2	-6.46	116.28	120.80	2	2
1	A	121	ILE	CA-C-N	6.40	135.02	117.10	6	1
1	A	95	ALA	CB-CA-C	-6.39	100.51	110.10	15	1
1	A	56	LEU	CB-CG-CD2	6.26	121.64	111.00	5	7
1	A	121	ILE	CB-CA-C	6.25	124.11	111.60	6	1
1	A	60	VAL	CA-C-O	-6.18	107.13	120.10	18	11
1	A	66	PHE	CB-CG-CD1	6.15	125.11	120.80	9	4
1	A	69	ASP	C-N-CA	6.15	137.07	121.70	1	7
1	A	72	HIS	N-CA-CB	5.99	121.39	110.60	14	2
1	A	100	MET	C-N-CA	5.95	136.57	121.70	16	2
1	A	102	ASN	N-CA-CB	5.88	121.19	110.60	12	2
1	A	130	ASN	CA-C-N	5.88	133.56	117.10	2	6
1	A	49	ASN	N-CA-CB	5.84	121.11	110.60	4	11
1	A	75	PHE	CB-CG-CD2	-5.80	116.74	120.80	3	2
1	A	99	TYR	N-CA-CB	5.73	120.92	110.60	15	2
1	A	120	GLU	C-N-CA	5.71	135.97	121.70	17	3
1	A	56	LEU	CB-CA-C	5.64	120.93	110.20	12	2
1	A	70	GLU	N-CA-CB	-5.59	100.54	110.60	9	2
1	A	60	VAL	O-C-N	-5.52	110.61	121.10	12	1
1	A	70	GLU	CA-CB-CG	5.50	125.51	113.40	10	2
1	A	79	ASP	N-CA-CB	5.47	120.46	110.60	18	1
1	A	55	TYR	CB-CG-CD2	-5.45	117.73	121.00	5	2
1	A	48	THR	C-N-CA	5.39	135.18	121.70	13	1
1	A	96	ALA	N-CA-CB	-5.38	102.56	110.10	12	1
1	A	57	THR	CA-CB-CG2	-5.33	104.94	112.40	10	1
1	A	60	VAL	CG1-CB-CG2	5.30	119.38	110.90	12	1
1	A	90	ILE	N-CA-CB	5.29	122.96	110.80	5	1
1	A	56	LEU	CA-CB-CG	5.24	127.36	115.30	6	4
1	A	117	PHE	CB-CG-CD1	5.23	124.46	120.80	8	1
1	A	131	PRO	C-N-CA	5.23	134.77	121.70	17	4
1	A	88	VAL	CB-CA-C	5.21	121.31	111.40	13	1
1	A	132	TYR	C-N-CA	5.21	134.72	121.70	19	1
1	A	68	LYS	CB-CA-C	5.21	120.81	110.40	1	1
1	A	73	ASN	CB-CA-C	5.20	120.80	110.40	3	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	101	THR	C-N-CA	5.19	134.68	121.70	14	1
1	A	62	LEU	N-CA-CB	5.14	120.67	110.40	11	1
1	A	90	ILE	CA-CB-CG1	5.12	120.73	111.00	5	1
1	A	121	ILE	CA-CB-CG1	5.08	120.66	111.00	20	1
1	A	72	HIS	CA-CB-CG	5.06	122.21	113.60	7	1
1	A	99	TYR	N-CA-C	5.02	124.55	111.00	9	1
1	A	57	THR	CA-C-N	5.02	128.24	117.20	16	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	75	PHE	Sidechain	17
1	A	132	TYR	Sidechain	15
1	A	99	TYR	Sidechain	8
1	A	117	PHE	Sidechain	2
1	A	55	TYR	Sidechain	2

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	636	637	637	4±2
All	All	12720	12740	12740	72

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:121:ILE:HG22	1:A:122:PRO:HD2	0.67	1.65	19	1
1:A:121:ILE:HG22	1:A:122:PRO:CD	0.63	2.22	19	1
1:A:88:VAL:O	1:A:92:VAL:HG13	0.59	1.97	11	2
1:A:88:VAL:O	1:A:92:VAL:HG23	0.53	2.02	4	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:57:THR:HG22	1:A:92:VAL:HG12	0.53	1.81	18	1
1:A:56:LEU:HD13	1:A:57:THR:H	0.52	1.63	1	1
1:A:63:LEU:HD22	1:A:121:ILE:HD12	0.51	1.82	17	1
1:A:56:LEU:HD21	1:A:93:ALA:HA	0.51	1.82	13	2
1:A:47:ILE:HG23	1:A:48:THR:H	0.51	1.65	9	1
1:A:132:TYR:CG	1:A:133:ARG:N	0.51	2.78	14	2
1:A:60:VAL:CG1	1:A:92:VAL:HG12	0.50	2.36	11	2
1:A:60:VAL:HG13	1:A:63:LEU:HD23	0.50	1.81	4	1
1:A:88:VAL:HG11	1:A:132:TYR:CG	0.50	2.42	19	1
1:A:59:MET:HE3	1:A:99:TYR:CE2	0.50	2.42	20	1
1:A:121:ILE:HD13	1:A:121:ILE:H	0.49	1.67	9	1
1:A:121:ILE:HB	1:A:122:PRO:HD2	0.49	1.83	20	6
1:A:56:LEU:HB2	1:A:96:ALA:HB1	0.49	1.83	12	2
1:A:89:LEU:HA	1:A:92:VAL:HG22	0.49	1.83	1	1
1:A:125:ILE:HD12	1:A:125:ILE:H	0.48	1.68	2	1
1:A:60:VAL:HG11	1:A:92:VAL:HA	0.48	1.84	16	1
1:A:64:VAL:HG12	1:A:91:PHE:CE2	0.48	2.43	7	1
1:A:56:LEU:HD23	1:A:57:THR:HG23	0.48	1.86	17	2
1:A:60:VAL:HG13	1:A:61:PRO:HD3	0.48	1.85	15	1
1:A:60:VAL:HG11	1:A:92:VAL:HG12	0.48	1.86	11	1
1:A:68:LYS:HG2	1:A:132:TYR:CE2	0.47	2.45	20	1
1:A:56:LEU:O	1:A:60:VAL:HG12	0.47	2.10	15	1
1:A:60:VAL:HB	1:A:92:VAL:HG12	0.46	1.86	8	1
1:A:63:LEU:HD22	1:A:117:PHE:CB	0.46	2.41	16	1
1:A:68:LYS:HD3	1:A:132:TYR:CE2	0.46	2.46	7	1
1:A:57:THR:HG22	1:A:92:VAL:HG23	0.45	1.88	3	2
1:A:62:LEU:HD12	1:A:117:PHE:CE2	0.45	2.46	11	1
1:A:60:VAL:HG21	1:A:92:VAL:HA	0.45	1.89	11	1
1:A:99:TYR:CE1	1:A:121:ILE:HG23	0.44	2.47	19	1
1:A:52:HIS:CE1	1:A:55:TYR:CD2	0.44	3.05	5	1
1:A:60:VAL:HG21	1:A:92:VAL:O	0.44	2.12	8	1
1:A:122:PRO:HG2	1:A:125:ILE:HD11	0.44	1.89	8	2
1:A:99:TYR:CD1	1:A:121:ILE:HG23	0.44	2.47	19	1
1:A:99:TYR:CE1	1:A:120:GLU:HB2	0.44	2.47	17	1
1:A:117:PHE:CD1	1:A:117:PHE:N	0.43	2.85	9	1
1:A:63:LEU:HD22	1:A:121:ILE:HB	0.43	1.89	2	1
1:A:64:VAL:HG21	1:A:88:VAL:HG13	0.43	1.91	2	1
1:A:59:MET:HB3	1:A:117:PHE:CZ	0.43	2.49	8	1
1:A:121:ILE:H	1:A:121:ILE:CD1	0.42	2.26	9	1
1:A:68:LYS:HG2	1:A:132:TYR:CD2	0.42	2.50	13	1
1:A:64:VAL:O	1:A:68:LYS:HB2	0.42	2.15	17	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:88:VAL:HG21	1:A:132:TYR:HA	0.42	1.91	19	1
1:A:132:TYR:CD2	1:A:133:ARG:N	0.42	2.88	12	1
1:A:99:TYR:CE1	1:A:121:ILE:CG2	0.41	3.03	15	1
1:A:91:PHE:CD2	1:A:129:LEU:HB2	0.41	2.50	6	1
1:A:63:LEU:HD22	1:A:117:PHE:HB3	0.41	1.92	16	1
1:A:60:VAL:CG1	1:A:92:VAL:HG22	0.41	2.45	18	1
1:A:94:LYS:HA	1:A:97:GLN:HG2	0.41	1.91	13	3
1:A:47:ILE:HG22	1:A:52:HIS:HB3	0.41	1.93	14	1
1:A:129:LEU:O	1:A:132:TYR:CD2	0.41	2.73	17	1
1:A:52:HIS:CE1	1:A:100:MET:SD	0.41	3.13	2	1
1:A:91:PHE:CE2	1:A:129:LEU:HD23	0.40	2.51	12	1
1:A:60:VAL:HG12	1:A:64:VAL:HG23	0.40	1.93	14	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/99 (80%)	67±2 (85±2%)	7±2 (8±2%)	5±2 (7±2%)	2	18
All	All	1580/1980 (80%)	1342 (85%)	132 (8%)	106 (7%)	2	18

All 20 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	49	ASN	18
1	A	72	HIS	14
1	A	132	TYR	13
1	A	50	ASP	12
1	A	121	ILE	11
1	A	116	ASN	6
1	A	102	ASN	6
1	A	78	LYS	4
1	A	133	ARG	4
1	A	123	SER	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	75	PHE	3
1	A	47	ILE	2
1	A	51	LYS	2
1	A	131	PRO	2
1	A	71	CYS	1
1	A	48	THR	1
1	A	52	HIS	1
1	A	117	PHE	1
1	A	120	GLU	1
1	A	82	GLU	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/88 (82%)	57±3 (79±4%)	15±3 (21±4%)	3	32
All	All	1440/1760 (82%)	1140 (79%)	300 (21%)	3	32

All 53 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	56	LEU	19
1	A	66	PHE	19
1	A	45	LEU	16
1	A	73	ASN	16
1	A	125	ILE	15
1	A	132	TYR	15
1	A	116	ASN	13
1	A	90	ILE	12
1	A	121	ILE	11
1	A	88	VAL	10
1	A	102	ASN	9
1	A	60	VAL	9
1	A	98	PHE	7
1	A	129	LEU	7
1	A	68	LYS	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	97	GLN	7
1	A	133	ARG	6
1	A	42	LYS	6
1	A	47	ILE	6
1	A	52	HIS	6
1	A	100	MET	6
1	A	57	THR	5
1	A	126	LEU	5
1	A	70	GLU	5
1	A	64	VAL	5
1	A	82	GLU	5
1	A	71	CYS	4
1	A	89	LEU	4
1	A	99	TYR	4
1	A	117	PHE	4
1	A	84	ILE	3
1	A	48	THR	3
1	A	75	PHE	3
1	A	101	THR	2
1	A	54	GLU	2
1	A	63	LEU	2
1	A	128	LYS	2
1	A	94	LYS	2
1	A	83	SER	2
1	A	58	GLU	2
1	A	59	MET	2
1	A	130	ASN	1
1	A	86	SER	1
1	A	72	HIS	1
1	A	51	LYS	1
1	A	43	ARG	1
1	A	49	ASN	1
1	A	78	LYS	1
1	A	123	SER	1
1	A	124	THR	1
1	A	65	GLU	1
1	A	62	LEU	1
1	A	81	ASN	1

6.3.3 RNA

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided