



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 15, 2024 – 07:26 AM EST

PDB ID : 2NTE
Title : Crystal Structure of the BARD1 BRCT Domains
Authors : Birrane, G.; Varma, A.K.; Soni, A.; Ladas, J.A.A.
Deposited on : 2006-11-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	210	
1	B	210	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	301	-	-	X	-
3	CL	B	305	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BRCA1-associated RING domain protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	210	Total 1693	C 1086	N 290	O 306	S 11	0	0	0
1	B	209	Total 1689	C 1084	N 289	O 305	S 11	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	2	Total Cl 2 2	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	133	Total O 133 133	0	2
5	B	105	Total O 105 105	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BRCA1-associated RING domain protein 1



- Molecule 1: BRCA1-associated RING domain protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.44Å 75.80Å 116.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.87 – 1.90 19.87 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.8 (19.87-1.90) 93.7 (19.87-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.57 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.247 0.200 , 0.255	Depositor DCC
R_{free} test set	1910 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3653	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.09	3/1735 (0.2%)	0.97	4/2347 (0.2%)
1	B	1.01	1/1731 (0.1%)	0.87	0/2341
All	All	1.05	4/3466 (0.1%)	0.92	4/4688 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	620	CYS	CB-SG	-6.61	1.71	1.82
1	A	728	SER	CB-OG	5.52	1.49	1.42
1	A	676	TYR	CD1-CE1	-5.17	1.31	1.39
1	A	736	TYR	CE1-CZ	-5.14	1.31	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	612	ASP	CB-CG-OD1	6.16	123.84	118.30
1	A	741	ASP	CB-CG-OD2	6.00	123.70	118.30
1	A	658	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	658	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	776	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1693	0	1703	31	0
1	B	1689	0	1701	37	0
2	A	5	0	0	0	0
2	B	5	0	0	2	0
3	A	1	0	0	0	0
3	B	2	0	0	2	0
4	A	12	0	18	4	0
4	B	8	0	12	3	0
5	A	133	0	0	3	0
5	B	105	0	0	6	0
All	All	3653	0	3434	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:614:VAL:HG23	1:A:653:ILE:HG21	1.32	1.10
1:B:701:GLN:HE21	4:B:308:EDO:H11	0.93	1.10
1:B:714:THR:HG22	5:B:214:HOH:O	1.60	1.00
1:B:701:GLN:NE2	4:B:308:EDO:H11	1.79	0.96
1:B:701:GLN:HE21	4:B:308:EDO:C1	1.83	0.88
1:A:587:GLU:O	1:A:590:VAL:HG23	1.79	0.81
1:B:575:SER:OG	2:B:301:SO4:O3	2.05	0.74
1:B:609:VAL:HG21	1:B:653:ILE:HD11	1.71	0.72
1:A:730:GLN:HE22	1:A:774:PRO:HA	1.57	0.69
1:B:711:SER:O	1:B:712:ASP:C	2.34	0.66
1:A:664:ARG:HG3	1:A:664:ARG:NH1	2.12	0.64
1:B:712:ASP:O	1:B:716:THR:HG23	1.99	0.63
1:B:587:GLU:O	1:B:590:VAL:HG22	1.99	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:724:ALA:HB3	1:A:731:ARG:HD2	1.82	0.62
1:A:572:LEU:HD12	1:A:606:HIS:HB2	1.84	0.59
1:B:771:GLU:HG2	1:B:773:LEU:HD23	1.84	0.59
1:A:664:ARG:HH11	1:A:664:ARG:CG	2.16	0.59
1:B:609:VAL:HG21	1:B:653:ILE:CD1	2.32	0.58
1:B:609:VAL:CG2	1:B:653:ILE:HD11	2.33	0.58
1:A:587:GLU:O	1:A:590:VAL:CG2	2.51	0.58
1:A:743:CYS:SG	1:A:745:TYR:HB3	2.42	0.58
1:B:708:LYS:H	1:B:714:THR:HG21	1.69	0.58
1:A:692:ILE:O	1:A:696:THR:HG23	2.04	0.57
1:B:570:LEU:HD22	1:B:606:HIS:HE1	1.68	0.57
1:A:664:ARG:HG3	1:A:664:ARG:HH11	1.69	0.57
1:A:590:VAL:HG23	5:A:54:HOH:O	2.05	0.56
1:A:776:ASP:OD1	1:A:777:SER:OG	2.23	0.56
1:A:642:ARG:HB2	1:A:644:VAL:HG12	1.89	0.55
1:B:570:LEU:HD22	1:B:606:HIS:CE1	2.42	0.54
1:B:709:PRO:O	1:B:710:ASP:OD1	2.26	0.53
1:B:667:LEU:CD1	1:B:667:LEU:N	2.71	0.52
1:A:664:ARG:NH1	1:A:664:ARG:CG	2.73	0.51
1:A:627:GLY:HA3	1:A:664:ARG:HG2	1.93	0.51
1:B:693:LYS:NZ	5:B:144:HOH:O	2.31	0.51
1:B:626:ASN:HB2	5:B:56:HOH:O	2.11	0.51
1:B:619:LYS:NZ	2:B:301:SO4:O1	2.44	0.50
1:B:621:MET:HE3	1:B:767:VAL:HB	1.93	0.50
1:A:634:GLU:HG3	5:A:53:HOH:O	2.11	0.50
1:A:614:VAL:CG2	1:A:653:ILE:HG21	2.23	0.49
1:A:640:LEU:O	1:A:643:LYS:NZ	2.28	0.49
1:A:621:MET:CE	4:A:310:EDO:H12	2.43	0.48
1:B:711:SER:O	1:B:713:VAL:N	2.46	0.48
1:B:579:SER:O	1:B:583:LYS:HG3	2.14	0.48
1:B:710:ASP:HA	3:B:305:CL:CL	2.52	0.46
1:B:730:GLN:HE22	1:B:774:PRO:HA	1.80	0.46
1:B:661:ARG:O	1:B:665:GLU:HG3	2.16	0.46
1:A:632:LYS:HD3	1:A:634:GLU:CD	2.36	0.45
1:B:715:GLN:HB2	3:B:305:CL:CL	2.52	0.45
1:B:745:TYR:OH	5:B:111:HOH:O	2.20	0.45
1:B:621:MET:HE3	1:B:767:VAL:CG1	2.47	0.45
1:A:730:GLN:NE2	1:A:774:PRO:HA	2.27	0.45
1:A:703:LEU:HD22	4:A:309:EDO:H22	1.99	0.44
1:A:590:VAL:CG2	5:A:54:HOH:O	2.63	0.43
1:A:621:MET:HE1	4:A:310:EDO:H12	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:587:GLU:HG2	1:A:640:LEU:HD21	2.00	0.43
1:A:748:GLU:H	1:A:748:GLU:CD	2.22	0.43
1:B:693:LYS:CE	5:B:144:HOH:O	2.66	0.43
1:B:708:LYS:N	1:B:714:THR:HG21	2.32	0.43
1:B:665:GLU:O	1:B:667:LEU:HD13	2.19	0.42
1:B:621:MET:HE3	1:B:767:VAL:HG11	2.02	0.42
1:A:632:LYS:HD3	1:A:634:GLU:OE1	2.19	0.42
1:B:601:ASP:HB3	1:B:603:THR:H	1.85	0.42
1:A:586:SER:O	1:A:590:VAL:HG22	2.19	0.41
1:A:713:VAL:HG22	4:A:307:EDO:O2	2.20	0.41
1:B:686:HIS:O	1:B:687:PRO:C	2.59	0.41
1:A:587:GLU:HG2	1:A:640:LEU:CD2	2.51	0.41
1:B:777:SER:HB3	5:B:116:HOH:O	2.20	0.41
1:B:667:LEU:N	1:B:667:LEU:HD12	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/210 (99%)	204 (98%)	3 (1%)	1 (0%)	25	17
1	B	207/210 (99%)	202 (98%)	3 (1%)	2 (1%)	13	5
All	All	415/420 (99%)	406 (98%)	6 (1%)	3 (1%)	19	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	710	ASP
1	B	712	ASP
1	A	655	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/189 (100%)	180 (95%)	9 (5%)	21	14
1	B	189/189 (100%)	181 (96%)	8 (4%)	25	18
All	All	378/378 (100%)	361 (96%)	17 (4%)	23	16

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	572	LEU
1	A	590	VAL
1	A	593	LYS
1	A	632	LYS
1	A	643	LYS
1	A	664	ARG
1	A	713	VAL
1	A	741	ASP
1	A	744	ASN
1	B	572	LEU
1	B	580	GLU
1	B	601	ASP
1	B	619	LYS
1	B	637	LYS
1	B	667	LEU
1	B	711	SER
1	B	727	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	663	ASN
1	A	686	HIS
1	A	690	ASN
1	A	730	GLN
1	A	744	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	663	ASN
1	B	701	GLN
1	B	730	GLN
1	B	744	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	301	-	4,4,4	0.33	0	6,6,6	1.07	0
4	EDO	A	307	-	3,3,3	0.28	0	2,2,2	0.12	0
4	EDO	B	311	-	3,3,3	1.11	0	2,2,2	0.12	0
4	EDO	A	310	-	3,3,3	0.47	0	2,2,2	0.78	0
4	EDO	A	309	-	3,3,3	1.17	0	2,2,2	1.46	0
2	SO4	A	302	-	4,4,4	0.32	0	6,6,6	0.86	0
4	EDO	B	308	-	3,3,3	0.56	0	2,2,2	1.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	307	-	-	0/1/1/1	-
4	EDO	B	311	-	-	0/1/1/1	-
4	EDO	A	310	-	-	1/1/1/1	-
4	EDO	A	309	-	-	0/1/1/1	-
4	EDO	B	308	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	308	EDO	O1-C1-C2-O2
4	A	310	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	SO4	2	0
4	A	307	EDO	1	0
4	A	310	EDO	2	0
4	A	309	EDO	1	0
4	B	308	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Warning: The R factor obtained from EDS is 0.2598, which does not match the depositor's R factor of 0.19618. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/210 (100%)	1.26	33 (15%) 6 5	41, 50, 66, 78	0
1	B	209/210 (99%)	1.01	24 (11%) 11 11	36, 50, 61, 66	0
All	All	419/420 (99%)	1.13	57 (13%) 8 8	36, 50, 63, 78	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	743	CYS	6.2
1	A	742	LEU	6.0
1	A	726	PRO	4.6
1	A	580	GLU	4.1
1	A	727	ASP	4.1
1	A	745	TYR	4.0
1	B	656	GLY	3.6
1	A	683	PHE	3.6
1	A	685	HIS	3.6
1	A	597	TYR	3.6
1	B	569	PRO	3.4
1	A	777	SER	3.3
1	B	713	VAL	3.3
1	B	668	LEU	3.2
1	B	777	SER	3.1
1	A	709	PRO	3.1
1	A	583	LYS	3.0
1	A	741	ASP	2.8
1	B	662	LEU	2.8
1	B	663	ASN	2.8
1	A	614	VAL	2.8
1	B	651	TYR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	660	SER	2.7
1	B	714	THR	2.6
1	A	713	VAL	2.6
1	B	590	VAL	2.5
1	A	653	ILE	2.5
1	B	591	ILE	2.5
1	A	613	ALA	2.5
1	A	654	PRO	2.4
1	A	612	ASP	2.4
1	A	664	ARG	2.4
1	A	629	TRP	2.3
1	A	628	CYS	2.3
1	A	772	LEU	2.3
1	A	770	PHE	2.3
1	A	746	HIS	2.3
1	B	626	ASN	2.3
1	A	625	LEU	2.3
1	B	742	LEU	2.3
1	A	569	PRO	2.2
1	B	710	ASP	2.2
1	B	709	PRO	2.2
1	B	580	GLU	2.2
1	A	680	TRP	2.2
1	B	647	GLN	2.2
1	B	601	ASP	2.2
1	B	653	ILE	2.2
1	B	725	ARG	2.1
1	A	591	ILE	2.1
1	B	613	ALA	2.1
1	A	610	PRO	2.1
1	B	624	ILE	2.1
1	A	626	ASN	2.1
1	A	668	LEU	2.1
1	B	667	LEU	2.0
1	A	630	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	310	4/4	0.74	0.19	57,58,60,61	0
4	EDO	B	308	4/4	0.76	0.24	61,66,66,66	0
2	SO4	A	302	5/5	0.79	0.15	61,62,66,69	0
4	EDO	B	311	4/4	0.79	0.16	48,48,49,50	0
4	EDO	A	309	4/4	0.81	0.21	41,50,51,57	0
2	SO4	B	301	5/5	0.81	0.14	64,66,68,70	0
3	CL	B	305	1/1	0.83	0.11	90,90,90,90	0
3	CL	A	304	1/1	0.86	0.12	63,63,63,63	0
3	CL	B	306	1/1	0.90	0.12	70,70,70,70	0
4	EDO	A	307	4/4	0.95	0.16	47,47,47,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.