



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 22, 2021 – 04:02 PM EDT

PDB ID : 2O8M  
Title : Crystal structure of the S139A mutant of Hepatitis C Virus NS3/4A protease  
Authors : Fischmann, T.O.; Prongay, A.J.; Madison, V.M.; Yao, N.  
Deposited on : 2006-12-12  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.22  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.22


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	 2% 80% 12% • 8%
1	B	200	 69% 8% 22%
2	C	23	 4% 83% 17%
2	D	23	 4% 65% 9% • 22%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	185	1357	846	247	254	10	0	0	1
1	B	155	1138	711	207	212	8	0	0	1

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9ELS8
A	-10	ALA	-	expression tag	UNP Q9ELS8
A	-9	SER	-	expression tag	UNP Q9ELS8
A	-8	MET	-	expression tag	UNP Q9ELS8
A	-7	THR	-	expression tag	UNP Q9ELS8
A	-6	GLY	-	expression tag	UNP Q9ELS8
A	-5	GLY	-	expression tag	UNP Q9ELS8
A	-4	GLN	-	expression tag	UNP Q9ELS8
A	-3	GLN	-	expression tag	UNP Q9ELS8
A	-2	MET	-	expression tag	UNP Q9ELS8
A	-1	GLY	-	expression tag	UNP Q9ELS8
A	119	ARG	GLN	conflict	UNP Q9ELS8
A	139	ALA	SER	conflict	UNP Q9ELS8
A	182	GLY	-	expression tag	UNP Q9ELS8
A	183	SER	-	expression tag	UNP Q9ELS8
A	184	HIS	-	expression tag	UNP Q9ELS8
A	185	HIS	-	expression tag	UNP Q9ELS8
A	186	HIS	-	expression tag	UNP Q9ELS8
A	187	HIS	-	expression tag	UNP Q9ELS8
A	188	HIS	-	expression tag	UNP Q9ELS8
A	189	HIS	-	expression tag	UNP Q9ELS8
B	-11	MET	-	expression tag	UNP Q9ELS8
B	-10	ALA	-	expression tag	UNP Q9ELS8
B	-9	SER	-	expression tag	UNP Q9ELS8
B	-8	MET	-	expression tag	UNP Q9ELS8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	THR	-	expression tag	UNP Q9ELS8
B	-6	GLY	-	expression tag	UNP Q9ELS8
B	-5	GLY	-	expression tag	UNP Q9ELS8
B	-4	GLN	-	expression tag	UNP Q9ELS8
B	-3	GLN	-	expression tag	UNP Q9ELS8
B	-2	MET	-	expression tag	UNP Q9ELS8
B	-1	GLY	-	expression tag	UNP Q9ELS8
B	119	ARG	GLN	conflict	UNP Q9ELS8
B	139	ALA	SER	conflict	UNP Q9ELS8
B	182	GLY	-	expression tag	UNP Q9ELS8
B	183	SER	-	expression tag	UNP Q9ELS8
B	184	HIS	-	expression tag	UNP Q9ELS8
B	185	HIS	-	expression tag	UNP Q9ELS8
B	186	HIS	-	expression tag	UNP Q9ELS8
B	187	HIS	-	expression tag	UNP Q9ELS8
B	188	HIS	-	expression tag	UNP Q9ELS8
B	189	HIS	-	expression tag	UNP Q9ELS8

- Molecule 2 is a protein called Protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	23	Total	C	N	O	S	0	0	0
			161	107	29	24	1			
2	D	18	Total	C	N	O	S	0	0	1
			118	76	23	18	1			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	219	LYS	-	expression tag	UNP P27958
C	220	LYS	-	expression tag	UNP P27958
C	240	LYS	-	expression tag	UNP P27958
C	241	LYS	-	expression tag	UNP P27958
D	219	LYS	-	expression tag	UNP P27958
D	220	LYS	-	expression tag	UNP P27958
D	240	LYS	-	expression tag	UNP P27958
D	241	LYS	-	expression tag	UNP P27958

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0
3	B	1	Total 1	Zn 1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Na 1	0	0
4	C	1	Total 1	Na 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	117	Total 117	O 117	0	0
5	B	92	Total 92	O 92	0	0
5	C	13	Total 13	O 13	0	0
5	D	13	Total 13	O 13	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protease




- Molecule 1: Protease



- Molecule 2: Protease



- Molecule 2: Protease



## 4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	224.64Å 224.64Å 75.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.00 70.27 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-2.00) 94.8 (70.27-2.00)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 2.00Å)	Xtrriage
Refinement program	BUSTER	Depositor
R, $R_{free}$	0.194 , 0.228 0.197 , 0.232	Depositor DCC
$R_{free}$ test set	2343 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.7	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 59.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3013	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1383	0.54	0/1885
1	B	0.40	0/1162	0.54	0/1587
2	C	0.70	1/162 (0.6%)	0.55	0/216
2	D	0.54	0/118	0.56	0/158
All	All	0.43	1/2825 (0.0%)	0.54	0/3846

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	222	CYS	CB-SG	-6.04	1.72	1.82

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1357	0	1367	21	0
1	B	1138	0	1139	9	0
2	C	161	0	194	1	0
2	D	118	0	138	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	117	0	0	3	0
5	B	92	0	0	0	0
5	C	13	0	0	0	0
5	D	13	0	0	0	0
All	All	3013	0	2838	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:ARG:HD2	1:A:134:TYR:CD2	2.15	0.80
1:A:161:ARG:HH11	1:A:161:ARG:HG2	1.47	0.80
1:A:127:LEU:HD12	1:B:99:CYS:HA	1.68	0.73
1:A:130:ARG:HD2	1:A:134:TYR:CE2	2.29	0.68
1:B:132:ILE:HD13	1:B:132:ILE:O	1.95	0.66
1:A:161:ARG:HG2	1:A:161:ARG:NH1	2.12	0.61
1:A:106:LEU:HD11	5:A:1083:HOH:O	2.07	0.55
1:A:132:ILE:HD12	1:A:132:ILE:O	2.08	0.53
1:A:172:VAL:O	1:A:175:LEU:HB2	2.10	0.52
1:B:122:SER:HB2	1:B:169:PHE:O	2.10	0.51
1:A:82:LEU:HD11	1:A:175:LEU:HD12	1.93	0.49
1:B:123:ARG:HD3	1:B:167:VAL:O	2.12	0.49
1:A:32:GLU:HA	1:A:92:ARG:O	2.14	0.47
1:A:82:LEU:CD1	1:A:175:LEU:HD12	2.46	0.46
1:A:106:LEU:HD13	1:A:143:LEU:CD1	2.46	0.46
1:B:145:CYS:HB2	1:B:146:PRO:HD2	1.98	0.46
1:A:155:ARG:NH2	1:A:168:ASP:HB3	2.31	0.45
1:B:110:HIS:O	1:B:111:ALA:HB3	2.17	0.45
5:A:1110:HOH:O	2:C:234:LYS:HD2	2.18	0.43
2:D:220:LYS:HG3	2:D:221:GLY:H	1.84	0.43
1:A:109:ARG:HG2	5:A:1031:HOH:O	2.18	0.43
1:A:43:PHE:N	1:A:43:PHE:CD1	2.87	0.42
1:A:126:LEU:HD11	1:A:167:VAL:HG13	2.01	0.42
1:A:161:ARG:NH1	1:A:161:ARG:CG	2.80	0.42
2:D:220:LYS:HG3	2:D:221:GLY:N	2.34	0.42
1:A:54:THR:HG22	1:A:83:VAL:CG2	2.50	0.42
1:A:54:THR:CG2	1:A:83:VAL:HG22	2.50	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:THR:HG22	1:A:83:VAL:HG22	2.02	0.41
1:B:104:LEU:HD22	1:B:151:VAL:HG21	2.02	0.40
1:B:145:CYS:HB2	1:B:146:PRO:CD	2.51	0.40
1:A:130:ARG:CD	1:A:134:TYR:CE2	3.02	0.40
1:B:43:PHE:CD1	1:B:43:PHE:N	2.89	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/200 (92%)	179 (98%)	4 (2%)	0	100	100
1	B	153/200 (76%)	149 (97%)	4 (3%)	0	100	100
2	C	21/23 (91%)	21 (100%)	0	0	100	100
2	D	16/23 (70%)	16 (100%)	0	0	100	100
All	All	373/446 (84%)	365 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/160 (92%)	138 (94%)	9 (6%)	18	14

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	124/160 (78%)	119 (96%)	5 (4%)	31	29
2	C	18/19 (95%)	16 (89%)	2 (11%)	6	3
2	D	13/19 (68%)	11 (85%)	2 (15%)	2	1
All	All	302/358 (84%)	284 (94%)	18 (6%)	19	14

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LEU
1	A	24	ARG
1	A	37	SER
1	A	102	SER
1	A	127	LEU
1	A	128	SER
1	A	161	ARG
1	A	165	LYS
1	A	172	VAL
1	B	92	ARG
1	B	132	ILE
1	B	168	ASP
1	B	172	VAL
1	B	181	SER
2	C	228	ARG
2	C	240	LYS
2	D	220	LYS
2	D	222	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/200 (92%)	0.15	3 (1%) 72 70	24, 35, 56, 73	0
1	B	155/200 (77%)	0.03	1 (0%) 89 88	27, 38, 58, 86	0
2	C	23/23 (100%)	0.19	1 (4%) 35 34	26, 31, 66, 82	0
2	D	18/23 (78%)	0.33	1 (5%) 24 23	25, 33, 56, 76	0
All	All	381/446 (85%)	0.11	6 (1%) 72 70	24, 36, 58, 86	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	GLY	4.7
2	D	237	ILE	3.9
1	A	-2	MET	2.4
2	C	241	LYS	2.2
1	A	181	SER	2.1
1	B	180	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NA	A	1002	1/1	0.97	0.09	23,23,23,23	0
3	ZN	A	201	1/1	0.99	0.14	38,38,38,38	0
4	NA	C	1001	1/1	0.99	0.05	21,21,21,21	0
3	ZN	B	201	1/1	1.00	0.15	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.