

Full wwPDB X-ray Structure Validation Report (i)

Oct 23, 2024 – 10:51 PM EDT

PDB ID	:	2OX9
Title	:	Mouse Scavenger Receptor C-type Lectin carbohydrate-recognition domain.
Authors	:	Weis, W.I.; Feinberg, H.; Drickamer, K.; Taylor, M.E.
Deposited on	:	2007-02-20
Resolution	:	1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} {\rm Whole \ archive} \\ (\#{\rm Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	3187 (1.96-1.96)
Clashscore	180529	3412(1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
			4%				
1	А	140	76%		17%		7%
			9%				
1	В	140	60%	30%		•	9%
			22%				
1	С	140	66%	24%		•	6%
			12%				
1	D	140	66%	25%		•	7%
2	Ε	3	100%				



Mol	Chain	Length	Quality of chain	
2	F	3	100%	
2	G	3	100%	
2	Н	3	67%	33%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Δ	120	Total	С	Ν	0	S	0	0	0	
	A	130	1079	683	181	209	6	0	0	0	
1	р	197	Total	С	Ν	0	S	0	0	0	0
	D	121	1066	679	176	205	6		0	0	
1	1 C	131	Total	С	Ν	0	S	0	0	0	
			1087	689	182	210	6	0	0	0	
1	П	120	Total	С	Ν	0	S	0	0	0	
	130	1079	683	181	209	6	0	0	0		

• Molecule 1 is a protein called Collectin placenta 1.

• Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[beta-D-galactopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	Е	3	Total C N O	0	0	0
			36 20 1 15			
2	F	3	Total C N O	0	0	0
2			36 20 1 15			0
2	С	2	Total C N O	0	0	0
2 G	5	36 20 1 15	0	0	U	
9	Ц	3	2 Total C N O O	0	0	0
	п	3	36 20 1 15	0	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	4	Total Ca 4 4	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	4	Total Ca 4 4	0	0
3	С	4	Total Ca 4 4	0	0
3	D	4	Total Ca 4 4	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	122	Total O 122 122	0	0
4	В	89	Total O 89 89	0	0
4	С	65	Total O 65 65	0	0
4	D	81	Total O 81 81	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Collectin placenta 1





 \bullet Molecule 2: alpha-L-fucopyranose-(1-3)-[beta-D-galactopyranose-(1-4)]2-acetamido-2-deoxy-bet a-D-glucopyranose

Chain E:

100%

NAG1 FUC2 GAL3

 \bullet Molecule 2: alpha-L-fucopyranose-(1-3)-[beta-D-galactopyranose-(1-4)]2-acetamido-2-deoxy-bet a-D-glucopyranose

Chain F:

100%

NAG1 FUC2 GAL3

 \bullet Molecule 2: alpha-L-fucopyranose-(1-3)-[beta-D-galactopyranose-(1-4)]2-acetamido-2-deoxy-bet a-D-glucopyranose

Chain G:

100%

67%

NAG1 FUC2 GAL3

 \bullet Molecule 2: alpha-L-fucopyranose-(1-3)-[beta-D-galactopyranose-(1-4)]2-acetamido-2-deoxy-bet a-D-glucopyranose

33%

Chain H:

NAG1 FUC2 GAL3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	48.00Å 53.76Å 59.08Å	Depositor
a, b, c, α , β , γ	67.75° 76.70° 85.37°	Depositor
Bosolution(A)	53.39 - 1.95	Depositor
Resolution (A)	53.39 - 1.95	EDS
% Data completeness	(Not available) (53.39-1.95)	Depositor
(in resolution range)	91.9 (53.39-1.95)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) > 1$	$6.11 (at 1.95 \text{\AA})$	Xtriage
Refinement program	CNS 1.1	Depositor
P. P.	0.224 , 0.273	Depositor
n, n_{free}	0.218 , 0.268	DCC
R_{free} test set	1785 reflections $(5.01%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	18.7	Xtriage
Anisotropy	0.486	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 46.9	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4828	wwPDB-VP
Average B, all atoms $(Å^2)$	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.28% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, CA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.36	0/1115	0.62	0/1508	
1	В	0.35	0/1101	0.60	0/1490	
1	С	0.32	0/1123	0.59	0/1520	
1	D	0.36	0/1115	0.60	0/1508	
All	All	0.35	0/4454	0.60	0/6026	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1079	0	963	18	0
1	В	1066	0	960	30	0
1	С	1087	0	974	33	0
1	D	1079	0	963	30	0
2	Е	36	0	32	0	0
2	F	36	0	31	0	0
2	G	36	0	31	0	0
2	Н	36	0	31	1	0
3	A	4	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	4	0	0	0	0
3	С	4	0	0	0	0
3	D	4	0	0	0	0
4	А	122	0	0	5	0
4	В	89	0	0	6	0
4	С	65	0	0	2	0
4	D	81	0	0	5	0
All	All	4828	0	3985	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:615:THR:HG22	1:D:616:ASP:H	1.45	0.81
1:A:668:LEU:HD11	1:A:685:VAL:CG1	2.11	0.80
1:B:648:ARG:HG2	1:B:649:GLU:OE1	1.82	0.80
1:A:668:LEU:HD11	1:A:685:VAL:HG12	1.62	0.80
1:C:735:GLU:H	1:C:735:GLU:CD	1.89	0.76
1:B:733:GLU:H	1:B:733:GLU:CD	1.88	0.76
1:B:676:GLU:HG2	1:B:678:LYS:HE3	1.67	0.74
1:C:615:THR:HG23	1:C:616:ASP:H	1.51	0.74
1:C:615:THR:HG23	1:C:616:ASP:N	2.05	0.72
1:D:615:THR:HG22	1:D:616:ASP:N	2.05	0.70
1:C:632:LYS:O	1:C:636:GLU:HG3	1.93	0.67
1:C:661:ARG:HB3	1:C:662:GLU:OE2	1.96	0.65
1:D:733:GLU:HG2	4:D:241:HOH:O	1.98	0.63
1:A:677:TRP:O	1:A:685:VAL:HG13	1.98	0.63
1:A:646:ASN:HB3	4:A:234:HOH:O	1.99	0.62
1:C:712:ILE:HD13	1:C:718:ASN:HB2	1.80	0.61
1:A:609:PRO:O	1:A:610:HIS:HB2	2.01	0.61
1:C:659:VAL:O	1:C:661:ARG:HG3	2.01	0.61
1:C:609:PRO:O	1:C:610:HIS:HB2	2.01	0.59
1:B:679:TRP:HZ3	4:B:239:HOH:O	1.84	0.59
1:C:668:LEU:HD11	1:C:685:VAL:CG2	2.32	0.59
1:B:625:LYS:HB3	1:B:725:ILE:HG22	1.84	0.59
1:D:674:GLU:O	1:D:675:SER:HB2	2.03	0.58
1:D:695:PRO:HD2	4:D:224:HOH:O	2.01	0.58
1:C:612:LYS:HE2	1:C:657:HIS:HD2	1.69	0.58
1:B:733:GLU:CD	1:B:733:GLU:N	2.55	0.58



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:677:TRP:O	1:C:685:VAL:HG23	2.05	0.57
1:B:668:LEU:HG	1:B:677:TRP:HE3	1.71	0.55
1:C:668:LEU:HD11	1:C:685:VAL:HG22	1.89	0.55
1:A:670:ASP:OD1	1:A:707:ASP:HA	2.07	0.54
1:C:668:LEU:HG	1:C:677:TRP:HE3	1.73	0.54
1:D:609:PRO:O	1:D:610:HIS:HB2	2.08	0.54
1:C:612:LYS:HE2	1:C:657:HIS:CD2	2.43	0.53
1:A:668:LEU:CD1	1:A:685:VAL:HG12	2.36	0.53
1:B:721:GLN:HE21	1:B:723:ASP:CG	2.11	0.53
1:B:609:PRO:O	1:B:610:HIS:HB2	2.08	0.53
1:D:644:PHE:HA	1:D:679:TRP:CE3	2.44	0.52
1:C:614:PHE:O	1:C:615:THR:HG22	2.10	0.52
1:D:698:TRP:CD2	2:H:3:GAL:H62	2.44	0.52
1:B:609:PRO:HG2	4:B:158:HOH:O	2.09	0.52
1:C:702:HIS:CG	1:C:703:GLY:H	2.28	0.52
1:A:632:LYS:O	1:A:636:GLU:HG3	2.09	0.52
1:B:686:ASP:O	1:B:688:LYS:HE2	2.10	0.52
1:C:632:LYS:NZ	4:C:842:HOH:O	2.42	0.52
1:B:667:GLY:N	4:B:239:HOH:O	2.42	0.51
1:D:721:GLN:HG2	1:D:723:ASP:OD1	2.10	0.51
1:C:721:GLN:HE21	1:C:723:ASP:CG	2.14	0.51
1:C:674:GLU:O	1:C:675:SER:HB2	2.12	0.50
1:C:702:HIS:CD2	1:C:703:GLY:H	2.29	0.50
1:A:661:ARG:HG2	4:A:200:HOH:O	2.12	0.50
1:C:691:LYS:HG3	1:C:716:GLN:NE2	2.26	0.50
1:B:674:GLU:O	1:B:675:SER:HB2	2.12	0.50
1:B:690:TRP:CD2	1:B:695:PRO:HD3	2.47	0.49
1:C:626:GLU:HB3	1:C:630:ASP:HB2	1.93	0.48
1:C:650:GLU:O	1:C:654:ILE:HG13	2.15	0.47
1:D:688:LYS:HA	1:D:717:TRP:CH2	2.49	0.47
1:C:611:TRP:CH2	1:C:732:LYS:HE2	2.49	0.47
1:B:712:ILE:HD13	1:B:718:ASN:HB2	1.97	0.47
1:B:668:LEU:HG	1:B:677:TRP:CE3	2.50	0.47
1:B:670:ASP:OD1	1:B:707:ASP:HA	2.14	0.47
1:B:666:ILE:HG13	4:B:239:HOH:O	2.15	0.46
1:C:659:VAL:CG1	1:C:661:ARG:HE	2.29	0.46
1:D:632:LYS:O	1:D:636:GLU:HG3	2.16	0.46
1:D:725:ILE:O	1:D:726:ASN:ND2	2.49	0.46
1:A:674:GLU:O	1:A:675:SER:HB2	2.16	0.46
1:C:659:VAL:O	1:C:660:GLY:C	2.54	0.46
1:A:684:PRO:HG3	4:A:275:HOH:O	2.16	0.46



	louo pugom	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:668:LEU:HG	1:A:677:TRP:HE3	1.81	0.45
1:D:644:PHE:HA	1:D:679:TRP:CZ3	2.51	0.45
1:D:615:THR:CG2	1:D:616:ASP:H	2.15	0.45
1:D:662:GLU:H	1:D:662:GLU:CD	2.20	0.45
1:D:615:THR:CG2	1:D:616:ASP:N	2.73	0.44
1:C:663:SER:HB2	1:C:726:ASN:ND2	2.33	0.44
1:D:696:ASP:O	1:D:697:ASN:C	2.55	0.44
1:B:655:LYS:HG3	4:B:104:HOH:O	2.18	0.44
1:D:668:LEU:HG	1:D:677:TRP:HE3	1.83	0.44
1:D:686:ASP:HB3	4:D:154:HOH:O	2.16	0.44
1:B:658:THR:HA	1:B:661:ARG:HD2	2.00	0.44
1:A:668:LEU:HB3	1:A:709:ALA:HB3	2.00	0.44
1:D:721:GLN:NE2	1:D:723:ASP:OD1	2.45	0.44
1:C:674:GLU:O	1:C:675:SER:CB	2.66	0.43
1:A:684:PRO:HD2	4:A:230:HOH:O	2.18	0.43
1:D:661:ARG:NH2	4:D:89:HOH:O	2.52	0.43
1:C:659:VAL:O	1:C:661:ARG:CG	2.67	0.43
1:D:697:ASN:ND2	1:D:700:SER:HB3	2.34	0.43
1:B:641:HIS:O	1:B:642:LEU:C	2.56	0.42
1:B:627:ILE:HA	1:B:724:GLU:O	2.20	0.42
1:B:615:THR:N	1:D:649:GLU:OE2	2.41	0.42
1:B:632:LYS:O	1:B:636:GLU:HG3	2.20	0.42
1:D:637:ASP:HB2	4:D:243:HOH:O	2.19	0.42
1:D:670:ASP:HB3	1:D:677:TRP:CD2	2.55	0.42
1:B:620:TYR:O	1:B:729:ILE:HA	2.20	0.42
1:A:690:TRP:CD2	1:A:695:PRO:HD3	2.55	0.42
1:C:652:GLN:NE2	4:C:844:HOH:O	2.46	0.41
1:C:659:VAL:O	1:C:661:ARG:N	2.54	0.41
1:D:658:THR:HG21	1:D:729:ILE:HD11	2.02	0.41
1:A:652:GLN:NE2	4:A:160:HOH:O	2.50	0.41
1:D:620:TYR:HB3	1:D:730:CYS:HB2	2.03	0.41
1:B:649:GLU:CD	1:B:649:GLU:H	2.24	0.41
1:B:607:CYS:N	4:B:279:HOH:O	2.53	0.41
1:D:611:TRP:CE2	1:D:620:TYR:HB2	2.56	0.41
1:A:667:GLY:O	1:A:679:TRP:HA	2.20	0.40
1:B:669:THR:HA	1:B:677:TRP:CZ3	2.57	0.40
1:B:618:CYS:SG	1:B:734:ARG:HG3	2.61	0.40
1:C:691:LYS:HG3	1:C:716:GLN:HE22	1.86	0.40
1:D:617:LYS:HD3	1:D:733:GLU:HA	2.04	0.40
1:A:615:THR:H	1:C:649:GLU:CD	2.24	0.40
1:B:690:TRP:CE2	1:B:695:PRO:HD3	2.57	0.40



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:D:688:LYS:HA	1:D:717:TRP:CZ3	2.56	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Per	cen	ntiles
1	А	128/140~(91%)	124 (97%)	3~(2%)	1 (1%)	1	.6	8
1	В	123/140~(88%)	115~(94%)	8~(6%)	0	10	0	100
1	С	129/140~(92%)	116 (90%)	9~(7%)	4 (3%)	•	3	0
1	D	128/140~(91%)	119 (93%)	7~(6%)	2(2%)	5	8	2
All	All	508/560~(91%)	474 (93%)	27 (5%)	7 (1%)	9	9	3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	660	GLY
1	D	697	ASN
1	С	659	VAL
1	С	689	ASN
1	С	712	ILE
1	D	712	ILE
1	А	712	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	115/124~(93%)	113~(98%)	2(2%)	56 52
1	В	115/124~(93%)	113~(98%)	2(2%)	56 52
1	С	116/124~(94%)	115 (99%)	1 (1%)	75 75
1	D	115/124~(93%)	114 (99%)	1 (1%)	75 75
All	All	461/496 (93%)	455 (99%)	6 (1%)	65 62

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	637	ASP
1	А	659	VAL
1	В	646	ASN
1	В	688	LYS
1	С	735	GLU
1	D	662	GLU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such side chains are listed below:

Mol	Chain	Res	Type
1	А	726	ASN
1	В	716	GLN
1	В	721	GLN
1	В	726	ASN
1	С	652	GLN
1	С	657	HIS
1	С	721	GLN
1	С	726	ASN
1	D	726	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Dog	Link	Bo	ond leng	\mathbf{ths}	В	ond ang	les
MOI	туре	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	NAG	Е	1	2	$15,\!15,\!15$	0.41	0	21,21,21	0.56	0
2	FUC	Е	2	2	10,10,11	0.46	0	14,14,16	0.33	0
2	GAL	Е	3	2,3	$11,\!11,\!12$	0.46	0	$15,\!15,\!17$	0.31	0
2	NAG	F	1	2	$15,\!15,\!15$	0.42	0	21,21,21	0.55	0
2	FUC	F	2	2	10,10,11	0.46	0	14,14,16	0.28	0
2	GAL	F	3	2,3	11,11,12	0.46	0	$15,\!15,\!17$	0.35	0
2	NAG	G	1	2	$15,\!15,\!15$	0.49	0	21,21,21	0.54	0
2	FUC	G	2	2	10,10,11	0.47	0	14,14,16	0.39	0
2	GAL	G	3	2,3	11,11,12	0.37	0	$15,\!15,\!17$	0.34	0
2	NAG	Н	1	2	$15,\!15,\!15$	0.45	0	21,21,21	0.57	0
2	FUC	Н	2	2	10,10,11	0.53	0	14,14,16	0.30	0
2	GAL	Н	3	2,3	11,11,12	0.42	0	$15,\!15,\!17$	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	Е	1	2	-	0/6/26/26	0/1/1/1
2	FUC	Е	2	2	-	-	0/1/1/1
2	GAL	Е	3	2,3	-	0/2/19/22	0/1/1/1
2	NAG	F	1	2	-	0/6/26/26	0/1/1/1
2	FUC	F	2	2	-	-	0/1/1/1
2	GAL	F	3	2,3	-	0/2/19/22	0/1/1/1
2	NAG	G	1	2	-	2/6/26/26	0/1/1/1
2	FUC	G	2	2	-	-	0/1/1/1
2	GAL	G	3	2,3	-	1/2/19/22	0/1/1/1
2	NAG	Н	1	2	-	2/6/26/26	0/1/1/1
2	FUC	Н	2	2	-	-	0/1/1/1
2	GAL	Н	3	2,3	-	0/2/19/22	0/1/1/1



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Н	1	NAG	O5-C5-C6-O6
2	Н	1	NAG	C4-C5-C6-O6
2	G	1	NAG	C4-C5-C6-O6
2	G	1	NAG	O5-C5-C6-O6
2	G	3	GAL	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Н	3	GAL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	130/140~(92%)	0.41	6 (4%) 38 44	6, 19, 32, 44	0
1	В	127/140~(90%)	0.57	12 (9%) 15 19	9, 20, 34, 38	0
1	С	131/140~(93%)	1.43	31 (23%) 2 2	14, 29, 45, 62	0
1	D	130/140~(92%)	0.81	17 (13%) 8 10	8, 23, 40, 48	0
All	All	518/560~(92%)	0.81	66 (12%) 9 11	6, 23, 40, 62	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	660	GLY	6.8
1	С	659	VAL	4.7
1	С	661	ARG	4.3
1	D	606	GLY	4.2
1	D	701	GLY	4.0
1	D	699	GLY	3.9
1	С	713	TYR	3.8
1	С	714	ALA	3.8
1	С	737	VAL	3.6
1	С	698	TRP	3.5
1	С	685	VAL	3.5
1	D	609	PRO	3.3
1	С	715	GLY	3.3
1	С	712	ILE	3.2
1	С	607	CYS	3.1
1	С	702	HIS	3.0
1	D	698	TRP	3.0
1	D	607	CYS	3.0
1	В	684	PRO	3.0
1	С	696	ASP	2.8
1	С	723	ASP	2.8



Mol	Chain	Res	Type	RSRZ
1	D	703	GLY	2.8
1	А	723	ASP	2.7
1	В	685	VAL	2.7
1	С	722	CYS	2.6
1	В	723	ASP	2.6
1	D	702	HIS	2.6
1	D	705	GLY	2.6
1	В	698	TRP	2.6
1	В	722	CYS	2.6
1	D	734	ARG	2.6
1	В	683	SER	2.5
1	D	735	GLU	2.5
1	С	648	ARG	2.5
1	С	658	THR	2.5
1	D	722	CYS	2.5
1	С	662	GLU	2.5
1	С	684	PRO	2.5
1	С	694	GLN	2.4
1	А	735	GLU	2.4
1	С	690	TRP	2.4
1	С	652	GLN	2.3
1	С	623	LEU	2.3
1	С	725	ILE	2.3
1	D	704	PRO	2.2
1	С	663	SER	2.2
1	В	607	CYS	2.2
1	С	654	ILE	2.2
1	В	672	GLU	2.2
1	С	720	PHE	2.2
1	В	686	ASP	2.1
1	В	655	LYS	2.1
1	С	625	LYS	2.1
1	С	703	GLY	2.1
1	А	722	CYS	2.1
1	D	610	HIS	2.1
1	D	615	THR	2.1
1	В	652	GLN	2.1
1	С	705	GLY	2.1
1	В	610	HIS	2.1
1	А	684	PRO	2.0
1	D	700	SER	2.0
1	А	646	ASN	2.0



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	С	686	ASP	2.0
1	А	675	SER	2.0
1	D	661	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	NAG	G	1	15/15	0.68	0.22	$49,\!53,\!56,\!56$	0
2	NAG	Е	1	15/15	0.72	0.17	30,37,42,43	0
2	FUC	Н	2	10/11	0.74	0.17	40,42,43,43	0
2	FUC	G	2	10/11	0.76	0.20	52,53,54,55	0
2	GAL	G	3	11/12	0.79	0.20	45,46,47,50	0
2	NAG	Н	1	15/15	0.80	0.15	35,42,46,46	0
2	FUC	F	2	10/11	0.87	0.12	30,31,34,36	0
2	NAG	F	1	15/15	0.87	0.11	27,31,34,36	0
2	GAL	F	3	11/12	0.90	0.12	$16,\!21,\!26,\!31$	0
2	GAL	Н	3	11/12	0.90	0.09	26,31,34,35	0
2	FUC	E	2	10/11	0.91	0.10	25,27,29,29	0
2	GAL	Е	3	11/12	0.95	0.07	18,23,25,26	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	CA	С	802	1/1	0.89	0.12	$37,\!37,\!37,\!37$	0
3	CA	С	801	1/1	0.94	0.06	$27,\!27,\!27,\!27$	0
3	CA	D	803	1/1	0.94	0.07	$28,\!28,\!28,\!28$	0
3	CA	D	802	1/1	0.97	0.04	24,24,24,24	0
3	CA	С	803	1/1	0.97	0.05	$27,\!27,\!27,\!27$	0
3	CA	А	803	1/1	0.98	0.04	18,18,18,18	0
3	CA	В	801	1/1	0.98	0.03	18,18,18,18	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
3	CA	В	804	1/1	0.98	0.03	$13,\!13,\!13,\!13$	0
3	CA	А	802	1/1	0.98	0.03	18,18,18,18	0
3	CA	В	802	1/1	0.99	0.04	$17,\!17,\!17,\!17$	0
3	CA	В	803	1/1	0.99	0.03	24,24,24,24	0
3	CA	С	804	1/1	0.99	0.03	16, 16, 16, 16	0
3	CA	D	801	1/1	0.99	0.04	$17,\!17,\!17,\!17$	0
3	CA	А	804	1/1	0.99	0.02	$15,\!15,\!15,\!15$	0
3	CA	А	801	1/1	0.99	0.03	16, 16, 16, 16	0
3	CA	D	804	1/1	0.99	0.03	11,11,11,11	0

6.5 Other polymers (i)

There are no such residues in this entry.

