



# Full wwPDB X-ray Structure Validation Report i

Jun 12, 2024 – 08:03 AM EDT

PDB ID : 2R7R  
Title : Crystal Structure of Rotavirus SA11 VP1/RNA (UGUGACC) complex  
Authors : Lu, X.; Harrison, S.C.; Tao, Y.J.; Patton, J.T.; Nibert, M.L.  
Deposited on : 2007-09-09  
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

**X-RAY DIFFRACTION**

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|--------------------------|--|
| $R_{free}$            | 130704                   | 3163 (2.60-2.60)                                   |
| Clashscore            | 141614                   | 3518 (2.60-2.60)                                   |
| Ramachandran outliers | 138981                   | 3455 (2.60-2.60)                                   |
| Sidechain outliers    | 138945                   | 3455 (2.60-2.60)                                   |
| RSRZ outliers         | 127900                   | 3104 (2.60-2.60)                                   |
| RNA backbone          | 3102                     | 1040 (2.90-2.30)                                   |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain   |
|-----|-------|--------|--|
| 1   | X     | 7      | 29% <span style="background-color: yellow; width: 71%;">71%</span>     |
| 2   | A     | 1095   | % <span style="background-color: green; width: 55%;">55%</span> 40% .. |

## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 8844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(\*UP\*GP\*UP\*GP\*AP\*CP\*C)-3').

| Mol | Chain | Residues | Atoms |    |    |    |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|---------|-------|
|     |       |          | Total | C  | N  | O  | P |         |         |       |
| 1   | X     | 7        | 145   | 66 | 25 | 48 | 6 | 0       | 0       | 0     |

- Molecule 2 is a protein called RNA-dependent RNA polymerase.

| Mol | Chain | Residues | Atoms |      |      |      |    | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|---------|-------|
|     |       |          | Total | C    | N    | O    | S  |         |         |       |
| 2   | A     | 1073     | 8699  | 5579 | 1448 | 1634 | 38 | 0       | 0       | 0     |





## 4 Data and refinement statistics i

| Property  | Value   | Source           |
|---|---|------------------|
| Space group   | P 21 21 21  | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 76.35 Å   112.75 Å   143.79 Å<br>90.00°   90.00°   90.00° | Depositor        |
| Resolution (Å)  | 50.00 – 2.60<br>40.59 – 2.60                              | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 84.5 (50.00-2.60)<br>84.5 (40.59-2.60)                    | Depositor<br>EDS |
| $R_{merge}$   | 0.07  | Depositor        |
| $R_{sym}$   | (Not available)   | Depositor        |
| $< I/\sigma(I) >$ <sup>1</sup>  | 1.65 (at 2.61 Å)  | Xtriage          |
| Refinement program  | CNS   | Depositor        |
| $R$ , $R_{free}$  | 0.237 , 0.287<br>0.232 , 0.280                            | Depositor<br>DCC |
| $R_{free}$ test set   | 3030 reflections (7.88%)                                  | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 52.0  | Xtriage          |
| Anisotropy  | 0.351   | Xtriage          |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.31 , 45.1   | EDS              |
| L-test for twinning <sup>2</sup>  | $<  L  > = 0.49$ , $< L^2 > = 0.32$                       | Xtriage          |
| Estimated twinning fraction   | No twinning to report.                                    | Xtriage          |
| $F_o, F_c$ correlation  | 0.93  | EDS              |
| Total number of atoms   | 8844  | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 52.0  | wwPDB-VP         |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



























*Continued from previous page...*

| Mol | Chain | Res  | Type |
|-----|-------|------|------|
| 2   | A     | 556  | GLN  |
| 2   | A     | 557  | THR  |
| 2   | A     | 584  | ILE  |
| 2   | A     | 600  | ASN  |
| 2   | A     | 619  | HIS  |
| 2   | A     | 671  | SER  |
| 2   | A     | 708  | ASP  |
| 2   | A     | 721  | ARG  |
| 2   | A     | 784  | ILE  |
| 2   | A     | 791  | LEU  |
| 2   | A     | 807  | THR  |
| 2   | A     | 821  | PHE  |
| 2   | A     | 824  | ASN  |
| 2   | A     | 860  | MET  |
| 2   | A     | 868  | LYS  |
| 2   | A     | 872  | ILE  |
| 2   | A     | 880  | ASP  |
| 2   | A     | 892  | LEU  |
| 2   | A     | 895  | GLN  |
| 2   | A     | 900  | MET  |
| 2   | A     | 916  | SER  |
| 2   | A     | 917  | ARG  |
| 2   | A     | 922  | GLU  |
| 2   | A     | 939  | VAL  |
| 2   | A     | 1022 | PHE  |
| 2   | A     | 1027 | PRO  |
| 2   | A     | 1047 | LYS  |
| 2   | A     | 1088 | GLU  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (32) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2   | A     | 36  | ASN  |
| 2   | A     | 48  | ASN  |
| 2   | A     | 63  | ASN  |
| 2   | A     | 143 | ASN  |
| 2   | A     | 185 | HIS  |
| 2   | A     | 289 | ASN  |
| 2   | A     | 294 | ASN  |
| 2   | A     | 308 | GLN  |
| 2   | A     | 473 | GLN  |
| 2   | A     | 496 | GLN  |

*Continued on next page...*



*Continued from previous page...*

| Mol | Chain | Res  | Type |
|-----|-------|------|------|
| 1   | X     | 1103 | U    |

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.