



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 12:58 pm BST

PDB ID : 2VNO
Title : Family 51 carbohydrate binding module from a family 98 glycoside hydrolase produced by *Clostridium perfringens* in complex with blood group B-trisaccharide ligand.
Authors : Gregg, K.J.; Finn, R.; Abbott, D.W.; Boraston, A.B.
Deposited on : 2008-02-05
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	 4% 86% 9% • 5%
1	B	180	 2% 88% 7% • •
2	C	3	 100%
2	D	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GAL	C	3	X	-	-	-

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 3327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CPE0329.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	171	1335	847	219	267	2	0	4	0
1	B	173	1350	856	220	272	2	0	5	0

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[beta-D-galactopyranose-(1-3)]beta-D-galactopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	3	33	18	15	0	0	0
2	D	3	33	18	15	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	286	286	286	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	288	Total 288	O 288	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CPE0329



- Molecule 1: CPE0329



- Molecule 2: alpha-L-fucopyranose-(1-2)-[beta-D-galactopyranose-(1-3)]beta-D-galactopyranose



- Molecule 2: alpha-L-fucopyranose-(1-2)-[beta-D-galactopyranose-(1-3)]beta-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	69.69Å 98.55Å 49.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 1.45 19.67 – 1.45	Depositor EDS
% Data completeness (in resolution range)	95.4 (100.00-1.45) 95.5 (19.67-1.45)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.180 , 0.226 0.187 , 0.237	Depositor DCC
R_{free} test set	2917 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3327	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6220e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GAL, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.73	3/1371 (0.2%)	1.58	9/1855 (0.5%)
1	B	1.00	2/1387 (0.1%)	0.96	3/1880 (0.2%)
All	All	2.72	5/2758 (0.2%)	1.30	12/3735 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	189[A]	GLU	CG-CD	94.44	2.93	1.51
1	A	189[B]	GLU	CG-CD	94.44	2.93	1.51
1	A	121	TYR	CD1-CE1	6.75	1.49	1.39
1	B	151	TYR	CD2-CE2	-6.02	1.30	1.39
1	B	163	TYR	CE1-CZ	5.16	1.45	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	189[A]	GLU	CG-CD-OE1	-25.74	66.82	118.30
1	A	189[B]	GLU	CG-CD-OE1	-25.74	66.82	118.30
1	A	189[A]	GLU	CB-CG-CD	-20.57	58.66	114.20
1	A	189[B]	GLU	CB-CG-CD	-20.57	58.66	114.20
1	A	189[A]	GLU	CG-CD-OE2	19.84	157.98	118.30
1	A	189[B]	GLU	CG-CD-OE2	19.84	157.98	118.30
1	B	40	ARG	NE-CZ-NH2	-9.37	115.61	120.30
1	A	194	ASP	CB-CG-OD2	-6.73	112.24	118.30
1	A	180[A]	ARG	NE-CZ-NH1	6.65	123.62	120.30
1	A	180[B]	ARG	NE-CZ-NH1	6.65	123.62	120.30
1	B	194	ASP	CB-CG-OD2	-6.58	112.38	118.30
1	B	40	ARG	NE-CZ-NH1	5.74	123.17	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1335	0	1298	13	0
1	B	1350	0	1311	15	0
2	C	33	0	30	0	0
2	D	33	0	30	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	286	0	0	5	2
4	B	288	0	0	7	3
All	All	3327	0	2669	30	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:LYS:CE	1:B:190:LYS:CG	2.51	0.88
1:A:39:SER:N	4:A:2004:HOH:O	2.12	0.82
1:B:210:VAL:C	4:B:2282:HOH:O	2.19	0.81
1:A:208:ASP:CA	1:A:208:ASP:CG	2.53	0.76
1:B:85:GLU:CB	1:B:85:GLU:CD	2.54	0.76
1:B:59:LYS:CG	4:B:2047:HOH:O	2.33	0.75
1:A:54:HIS:HA	1:A:104:SER:HB2	1.69	0.74
1:A:190:LYS:CD	1:A:190:LYS:NZ	2.50	0.74
2:D:1:GAL:C3	2:D:3:GAL:C1	2.67	0.72
1:B:38:GLU:OE2	1:B:38:GLU:CG	2.38	0.72
1:B:210:VAL:C	4:B:2280:HOH:O	2.30	0.70
1:A:85:GLU:CD	4:A:2106:HOH:O	2.30	0.70
2:D:1:GAL:O3	2:D:3:GAL:C2	2.42	0.68
1:B:57:ASP:OD2	4:B:2050:HOH:O	2.14	0.66
1:A:51:ASN:ND2	4:A:2020:HOH:O	2.33	0.61
1:A:176:GLU:OE2	1:A:176:GLU:OE1	2.20	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:LYS:HG2	4:B:2047:HOH:O	2.00	0.58
1:B:54:HIS:HA	1:B:104:SER:HB2	1.88	0.55
1:B:59:LYS:HG3	4:B:2047:HOH:O	2.03	0.54
1:B:85:GLU:O	1:B:85:GLU:OE1	2.30	0.49
1:B:190:LYS:CG	1:B:190:LYS:NZ	2.76	0.48
1:A:113:ALA:HA	1:A:208:ASP:O	2.14	0.47
1:B:133[A]:GLU:OE2	1:B:133[A]:GLU:HA	2.15	0.46
1:B:76:GLN:NE2	4:B:2097:HOH:O	2.49	0.45
1:A:85:GLU:OE2	4:A:2106:HOH:O	2.21	0.44
1:A:85:GLU:OE1	1:A:85:GLU:N	2.48	0.43
1:A:39:SER:N	4:A:2001:HOH:O	2.52	0.43
1:A:59:LYS:HD3	1:A:59:LYS:HA	1.84	0.42
1:A:161:LEU:HD12	1:A:161:LEU:N	2.34	0.42
1:B:85:GLU:CD	1:B:85:GLU:CA	2.88	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2178:HOH:O	4:B:2158:HOH:O[4_455]	2.03	0.17
4:B:2051:HOH:O	4:B:2055:HOH:O[2_655]	2.04	0.16
4:A:2067:HOH:O	4:B:2238:HOH:O[4_455]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/180 (96%)	171 (99%)	1 (1%)	0	100	100
1	B	176/180 (98%)	174 (99%)	2 (1%)	0	100	100
All	All	348/360 (97%)	345 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/154 (94%)	142 (98%)	3 (2%)	53	19
1	B	146/154 (95%)	145 (99%)	1 (1%)	84	65
All	All	291/308 (94%)	287 (99%)	4 (1%)	67	37

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	A	104	SER
1	A	200	ASP
1	B	104	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	173	ASN
1	B	67	HIS
1	B	76	GLN
1	B	173	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAL	C	1	2	12,12,12	0.69	0	17,17,17	0.92	1 (5%)
2	FUC	C	2	2	10,10,11	0.95	1 (10%)	14,14,16	1.14	1 (7%)
2	GAL	C	3	2	11,11,12	0.81	1 (9%)	15,15,17	0.89	0
2	GAL	D	1	2	12,12,12	0.67	0	17,17,17	1.31	2 (11%)
2	FUC	D	2	2	10,10,11	0.65	0	14,14,16	0.99	1 (7%)
2	GAL	D	3	2	11,11,12	0.69	0	15,15,17	1.91	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	C	1	2	-	0/2/22/22	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1
2	GAL	C	3	2	1/1/4/5	0/2/19/22	0/1/1/1
2	GAL	D	1	2	-	0/2/22/22	0/1/1/1
2	FUC	D	2	2	-	-	0/1/1/1
2	GAL	D	3	2	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	GAL	C2-C3	2.21	1.55	1.52
2	C	2	FUC	O2-C2	-2.17	1.38	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	GAL	C1-O5-C5	3.99	117.59	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GAL	O3-C3-C2	-3.70	101.79	110.35
2	D	3	GAL	O5-C1-C2	3.50	116.17	110.77
2	D	3	GAL	C1-C2-C3	3.29	113.71	109.67
2	C	1	GAL	O3-C3-C2	-2.56	104.44	110.35
2	C	2	FUC	C3-C4-C5	2.15	113.12	109.77
2	D	1	GAL	O5-C1-C2	2.14	114.11	110.28
2	D	2	FUC	C6-C5-C4	-2.12	109.16	113.07
2	D	3	GAL	O3-C3-C2	2.03	113.88	109.99

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	3	GAL	C1


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3	GAL	2	0
2	D	1	GAL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	171/180 (95%)	0.53	7 (4%) 37 40	11, 15, 23, 33	1 (0%)
1	B	173/180 (96%)	0.53	4 (2%) 60 63	11, 14, 24, 31	0
All	All	344/360 (95%)	0.53	11 (3%) 47 50	11, 14, 24, 33	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	GLY	4.0
1	A	208	ASP	2.9
1	B	80	ILE	2.9
1	B	157	PHE	2.9
1	B	57	ASP	2.8
1	A	192	TRP	2.4
1	A	209	PHE	2.1
1	B	159	ASN	2.1
1	A	156	GLN	2.0
1	A	57	ASP	2.0
1	A	174	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FUC	C	2	10/11	0.84	0.18	17,19,22,23	0
2	GAL	C	1	12/12	0.89	0.15	15,19,23,27	0
2	GAL	C	3	11/12	0.90	0.12	13,14,17,18	0
2	FUC	D	2	10/11	0.90	0.18	17,19,21,22	0
2	GAL	D	1	12/12	0.91	0.12	16,18,22,23	0
2	GAL	D	3	11/12	0.93	0.09	12,14,16,17	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CA	A	1213	1/1	1.00	0.07	13,13,13,13	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CA	B	1214	1/1	1.00	0.07	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.