



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 05:20 PM EST

PDB ID : 2W1Z  
Title : ROP2 from *Toxoplasma gondii*: A virulence factor with a protein- kinase fold and no enzymatic activity.  
Authors : Labesse, G.; Gelin, M.; Bessin, Y.; Lebrun, M.; Arold, S.T.; Dubremetz, J.-F.  
Deposited on : 2008-10-21  
Resolution : 1.97 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	368	
1	B	368	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	1559	-	-	X	-

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ROP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	354	Total 2930	C 1857	N 543	O 522	S 8	0	10	0
1	B	358	Total 2956	C 1874	N 541	O 533	S 8	0	9	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	228	Total 228	O 228	0	0
3	B	296	Total 296	O 296	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ROP2



- Molecule 1: ROP2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	187.61Å 51.35Å 125.58Å 90.00° 128.65° 90.00°	Depositor
Resolution (Å)	73.32 – 1.97 73.32 – 1.97	Depositor EDS
% Data completeness (in resolution range)	82.3 (73.32-1.97) 82.3 (73.32-1.97)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 1.97Å)	Xtrriage
Refinement program	REFMAC 5.4.0062	Depositor
R, $R_{free}$	0.193 , 0.240 0.214 , 0.259	Depositor DCC
$R_{free}$ test set	2766 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtrriage
Anisotropy	0.379	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 58.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6430	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/3055	0.70	1/4143 (0.0%)
1	B	0.50	0/3078	0.69	2/4176 (0.0%)
All	All	0.48	0/6133	0.70	3/8319 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1
1	B	1	0
All	All	2	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	463	HIS	CB-CA-C	7.18	124.76	110.40
1	B	451	THR	OG1-CB-CG2	5.91	123.60	110.00
1	B	322	ARG	NE-CZ-NH1	-5.42	117.59	120.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	463	HIS	CA
1	B	451	THR	CB

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	276	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2930	0	2930	32	0
1	B	2956	0	2936	36	0
2	A	5	0	0	0	0
2	B	15	0	0	2	0
3	A	228	0	0	2	0
3	B	296	0	0	4	0
All	All	6430	0	5866	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:395[A]:ARG:HG2	1:B:535:MET:HE1	1.56	0.86
1:A:258:VAL:HG22	1:A:268:LEU:CD1	2.13	0.78
1:B:541:GLU:OE2	1:B:544[A]:ARG:NH1	2.16	0.77
1:B:395[A]:ARG:HG2	1:B:535:MET:CE	2.20	0.71
1:B:305:LEU:HD12	1:B:308:LEU:HD12	1.72	0.71
1:B:228:LEU:HD11	1:B:348:MET:HG2	1.72	0.70
1:A:343:LEU:O	1:A:343:LEU:HD23	1.93	0.69
1:A:515:ALA:HB3	1:A:539:GLU:HG2	1.77	0.67
1:A:214:VAL:HG23	1:A:216:ASN:ND2	2.12	0.65
1:B:511:GLN:OE1	3:B:2235:HOH:O	2.14	0.65
1:A:232:ALA:HB2	1:A:341:VAL:HG21	1.82	0.61
1:A:340:ARG:HD3	1:A:349:TRP:CZ2	2.37	0.60
1:B:393:LEU:HD22	1:B:483:ILE:HD11	1.87	0.56
1:B:305:LEU:CD1	1:B:308:LEU:HD12	2.35	0.56
1:B:232:ALA:HB2	1:B:341:VAL:HG21	1.88	0.54
1:B:232:ALA:HB2	1:B:341:VAL:CG2	2.37	0.54
1:A:303:LEU:HD23	1:A:306[A]:ARG:HH21	1.71	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:448:PRO:HB2	1:B:451:THR:HG23	1.88	0.54
1:B:522[A]:ARG:NH1	2:B:1559:SO4:S	2.81	0.53
1:A:470:PRO:HA	1:A:473:THR:HG22	1.90	0.53
1:B:254:VAL:HG22	1:B:273:GLN:HE21	1.73	0.53
1:B:322:ARG:HD2	1:B:424:PHE:CE1	2.44	0.53
1:B:205:LEU:HA	1:B:381:LEU:HD21	1.89	0.52
1:B:522[A]:ARG:NH1	2:B:1559:SO4:O2	2.43	0.52
1:A:305:LEU:HD12	1:A:308:LEU:HD12	1.91	0.52
1:B:462:HIS:N	1:B:462:HIS:CD2	2.78	0.52
1:A:314:GLN:HG2	1:A:326:PRO:O	2.11	0.51
1:B:392:GLN:HE22	1:B:395[B]:ARG:HH11	1.59	0.51
1:B:218:ASN:HD22	1:B:221:VAL:H	1.59	0.50
1:A:251:ARG:HH11	1:A:279:LEU:HD11	1.76	0.49
1:A:258:VAL:HG22	1:A:268:LEU:HD11	1.93	0.49
1:A:251:ARG:NH1	1:A:279:LEU:HD11	2.28	0.48
1:B:552:PRO:O	1:B:555:GLN:HG2	2.13	0.48
1:B:544[B]:ARG:NH1	3:B:2274:HOH:O	2.48	0.47
1:B:215:TRP:CD2	1:B:260:ARG:HD2	2.50	0.46
1:A:258:VAL:HG22	1:A:268:LEU:HD13	1.95	0.46
1:A:205:LEU:HD12	1:A:209:ILE:HD12	1.98	0.45
1:A:258:VAL:HA	1:A:268:LEU:HD13	1.98	0.45
1:A:463:HIS:HB2	1:A:464:PRO:HD2	1.98	0.45
1:A:290:PRO:HD3	1:A:349:TRP:CG	2.52	0.45
1:A:556:THR:C	3:A:2227:HOH:O	2.55	0.45
1:A:275:THR:O	1:A:277:GLU:N	2.50	0.44
1:B:466:LEU:HG	1:B:468:THR:HG23	1.99	0.44
1:B:438:ALA:HB3	1:B:467:MET:HG3	2.00	0.43
1:A:301:GLN:HG3	1:A:429:GLU:HG3	2.00	0.43
1:B:254:VAL:CG2	1:B:273:GLN:HE21	2.30	0.43
1:A:235:ARG:H	1:A:238:GLN:NE2	2.17	0.42
1:A:275:THR:O	1:A:277:GLU:CB	2.67	0.42
1:B:511:GLN:NE2	3:B:2234:HOH:O	2.49	0.42
1:B:395[A]:ARG:CG	1:B:535:MET:CE	2.94	0.42
1:A:361:GLN:HG3	1:A:419:GLN:HA	2.02	0.42
1:B:451:THR:HG21	1:B:471:PHE:CE1	2.54	0.42
1:B:451:THR:HA	1:B:454:GLU:HG2	2.02	0.41
1:B:305:LEU:HD21	1:B:313:ASN:C	2.40	0.41
1:A:205:LEU:HA	1:A:381:LEU:HD21	2.02	0.41
1:A:205:LEU:CD1	1:A:209:ILE:HD12	2.50	0.41
1:A:469:PHE:N	1:A:470:PRO:CD	2.83	0.41
1:A:492:GLU:HG2	1:A:493:ASP:OD1	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:547:LEU:O	1:B:551:LEU:HG	2.21	0.41
1:A:404:GLY:O	1:A:433[B]:ARG:HD3	2.21	0.41
1:B:555:GLN:HE21	1:B:557:ASP:HB3	1.86	0.41
1:B:556:THR:HG22	1:B:556:THR:O	2.21	0.41
1:A:309:ARG:HG3	3:A:2065:HOH:O	2.21	0.41
1:A:405:LEU:HD13	1:A:431:LEU:HD21	2.03	0.41
1:B:393:LEU:HD22	1:B:483:ILE:CD1	2.51	0.41
1:A:454:GLU:HG3	1:A:465:THR:HG22	2.02	0.40
1:B:322:ARG:HD3	3:B:2001:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/368 (97%)	352 (98%)	5 (1%)	1 (0%)	37	27
1	B	363/368 (99%)	357 (98%)	6 (2%)	0	100	100
All	All	721/736 (98%)	709 (98%)	11 (2%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/328 (99%)	313 (96%)	13 (4%)	27	16
1	B	328/328 (100%)	309 (94%)	19 (6%)	17	7
All	All	654/656 (100%)	622 (95%)	32 (5%)	23	10

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	234	TRP
1	A	268	LEU
1	A	301	GLN
1	A	306[A]	ARG
1	A	306[B]	ARG
1	A	322	ARG
1	A	340	ARG
1	A	343	LEU
1	A	349	TRP
1	A	356	LEU
1	A	463	HIS
1	A	500	GLU
1	A	547	LEU
1	B	234	TRP
1	B	248	GLU
1	B	260	ARG
1	B	288	GLU
1	B	293	ASN
1	B	315	ARG
1	B	322	ARG
1	B	346[A]	ARG
1	B	346[B]	ARG
1	B	356	LEU
1	B	442	ILE
1	B	451	THR
1	B	462	HIS
1	B	463	HIS
1	B	500	GLU
1	B	544[A]	ARG
1	B	544[B]	ARG
1	B	547	LEU
1	B	557	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	216	ASN
1	A	238	GLN
1	A	273	GLN
1	B	218	ASN
1	B	273	GLN
1	B	293	ASN
1	B	392	GLN
1	B	402	HIS
1	B	419	GLN
1	B	462	HIS
1	B	555	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	1558	-	4,4,4	0.22	0	6,6,6	0.53	0
2	SO4	B	1559	-	4,4,4	0.25	0	6,6,6	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	1560	-	4,4,4	0.22	0	6,6,6	0.16	0
2	SO4	A	1557	-	4,4,4	0.21	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1559	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

**Warning:** The R factor obtained from EDS is 0.2486, which does not match the depositor's R factor of 0.19339. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	354/368 (96%)	0.86	39 (11%) 12 17	18, 30, 44, 47	46 (12%)
1	B	358/368 (97%)	0.89	38 (10%) 13 18	17, 29, 42, 47	50 (13%)
All	All	712/736 (96%)	0.88	77 (10%) 12 18	17, 29, 43, 47	96 (13%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	455	ARG	6.0
1	A	556	THR	5.9
1	A	343	LEU	5.6
1	B	453	ALA	4.9
1	A	194	HIS	4.3
1	A	453	ALA	3.8
1	B	305	LEU	3.7
1	B	307	LEU	3.7
1	A	346	ARG	3.6
1	B	462	HIS	3.6
1	B	557	ASP	3.6
1	B	346[A]	ARG	3.6
1	B	321	LEU	3.5
1	B	323	PHE	3.5
1	B	195	THR	3.5
1	B	442	ILE	3.3
1	A	276	GLY	3.2
1	B	454	GLU	3.1
1	A	463	HIS	3.1
1	B	315	ARG	3.0
1	B	308	LEU	3.0
1	B	247	GLY	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	311	ILE	2.9
1	A	341	VAL	2.9
1	A	349	TRP	2.9
1	B	345[A]	GLU	2.9
1	A	221	VAL	2.9
1	B	322	ARG	2.9
1	B	341	VAL	2.9
1	B	310	GLY	2.8
1	A	275	THR	2.8
1	B	194	HIS	2.8
1	B	324	ILE	2.8
1	B	317	ALA	2.8
1	A	299	SER	2.8
1	A	454	GLU	2.8
1	B	222	LEU	2.7
1	B	215	TRP	2.7
1	B	325	PHE	2.7
1	B	508	ASN	2.7
1	A	320	TYR	2.7
1	A	311	ILE	2.6
1	A	310	GLY	2.6
1	B	279	LEU	2.5
1	B	343	LEU	2.5
1	A	303	LEU	2.5
1	B	463	HIS	2.4
1	B	246	LEU	2.4
1	A	216	ASN	2.4
1	A	249[A]	ARG	2.4
1	A	214	VAL	2.3
1	A	207[A]	ASN	2.3
1	A	442	ILE	2.3
1	B	439	VAL	2.3
1	A	309	ARG	2.2
1	B	342	ARG	2.2
1	B	303[A]	LEU	2.2
1	A	438	ALA	2.2
1	B	339	ILE	2.2
1	A	302	VAL	2.2
1	A	306[A]	ARG	2.2
1	A	295	ILE	2.2
1	B	452	THR	2.2
1	A	223	SER	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	376	SER	2.1
1	A	340	ARG	2.1
1	B	320	TYR	2.1
1	A	250	PRO	2.1
1	A	347	ASP	2.1
1	B	358	PRO	2.1
1	A	499	ILE	2.1
1	A	455	ARG	2.1
1	A	298	LEU	2.1
1	A	279	LEU	2.0
1	A	277	GLU	2.0
1	A	230	SER	2.0
1	A	286	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	1560	5/5	0.60	0.25	54,54,55,55	5
2	SO4	A	1557	5/5	0.86	0.17	43,44,44,44	5
2	SO4	B	1559	5/5	0.92	0.17	35,36,37,37	5
2	SO4	B	1558	5/5	0.96	0.18	28,28,31,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.