



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 03:55 pm GMT

PDB ID : 2WI2  
Title : Orally Active 2-Amino Thienopyrimidine Inhibitors of the Hsp90 Chaperone  
Authors : Brough, P.A.; Barril, X.; Borgognoni, J.; Chene, P.; Davies, N.G.M.; Davis, B.; Drysdale, M.J.; Dymock, B.; Eccles, S.A.; Garcia-Echeverria, C.; Fromont, C.; Hayes, A.; Hubbard, R.E.; Jordan, A.M.; Rugaard-Jensen, M.; Massey, A.; Merret, A.; Padfield, A.; Parsons, R.; Radimerski, T.; Raynaud, F.I.; Robertson, A.; Roughley, S.D.; Schoepfer, J.; Simmonite, H.; Surgenor, A.; Valenti, M.; Walls, S.; Webb, P.; Wood, M.; Workman, P.; Wright, L.M.  
Deposited on : 2009-05-08  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	
1	B	236	

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEAT SHOCK PROTEIN HSP 90-ALPHA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	209	Total 1634	C 1038	N 269	O 322	S 5	0	1	1
1	B	209	Total 1632	C 1036	N 270	O 321	S 5	0	0	1

- Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
2	A	1	Total 4	C 2	O 1	S 1	0	0

- Molecule 3 is 4-METHYL-6-(METHYLSULFANYL)-1,3,5-TRIAZIN-2-AMINE (three-letter code: ZZ3) (formula: C<sub>5</sub>H<sub>8</sub>N<sub>4</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
3	A	1	10	5	4	1	0	0
3	B	1	10	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	119	119	119	0	0
4	B	119	119	119	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEAT SHOCK PROTEIN HSP 90-ALPHA



- Molecule 1: HEAT SHOCK PROTEIN HSP 90-ALPHA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.27Å 88.44Å 98.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.09 29.26 – 2.09	Depositor EDS
% Data completeness (in resolution range)	92.3 (100.00-2.09) 92.2 (29.26-2.09)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.237 , 0.289 0.231 , 0.240	Depositor DCC
$R_{free}$ test set	1591 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtrriage
Anisotropy	0.385	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3528	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8787e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, ZZ3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.04	3/1663 (0.2%)	0.97	4/2244 (0.2%)
1	B	1.03	0/1658	1.00	3/2237 (0.1%)
All	All	1.03	3/3321 (0.1%)	0.99	7/4481 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	TYR	CD1-CE1	7.05	1.50	1.39
1	A	144	VAL	CB-CG1	5.23	1.63	1.52
1	A	223	GLU	C-N	-5.02	1.22	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	NE-CZ-NH1	9.00	124.80	120.30
1	A	103	LEU	CA-CB-CG	6.85	131.05	115.30
1	B	180	MET	C-N-CA	-6.43	108.81	122.30
1	B	107	LEU	CA-C-N	5.84	127.88	116.20
1	A	87	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	32	LEU	CB-CG-CD1	-5.07	102.39	111.00
1	B	107	LEU	C-N-CA	5.00	132.80	122.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	107	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1634	0	1630	20	0
1	B	1632	0	1630	42	0
2	A	4	0	6	0	1
3	A	10	0	8	0	0
3	B	10	0	8	0	0
4	A	119	0	0	3	1
4	B	119	0	0	5	0
All	All	3528	0	3282	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:LEU:HD13	1:A:119:MET:HE3	1.17	1.14
1:A:32:LEU:HD13	1:A:119:MET:CE	1.87	1.05
1:B:29:LEU:HB2	1:B:119:MET:HE2	1.47	0.96
1:B:154:HIS:CD2	1:B:156:ASP:H	1.87	0.93
1:A:32:LEU:CD1	1:A:119:MET:HE3	1.98	0.92
1:A:32:LEU:CD1	1:A:119:MET:CE	2.48	0.92
1:B:29:LEU:N	1:B:119:MET:HE1	1.85	0.91
1:B:154:HIS:HD2	1:B:156:ASP:H	0.96	0.91
1:B:29:LEU:HB2	1:B:119:MET:CE	2.07	0.84
1:A:189:HIS:HB3	4:A:2043:HOH:O	1.77	0.83
1:B:114:GLY:HA3	1:B:134:PHE:O	1.83	0.79
1:B:23:GLN:NE2	1:B:110:ILE:H	1.81	0.79
1:B:154:HIS:HD2	1:B:156:ASP:N	1.80	0.75

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:SER:HB2	4:B:2027:HOH:O	1.84	0.75
1:B:224:LYS:N	4:B:2119:HOH:O	2.19	0.75
1:B:29:LEU:CB	1:B:119:MET:HE2	2.16	0.75
1:A:32:LEU:HD12	1:A:119:MET:HE2	1.69	0.73
1:B:178:GLU:HG3	4:B:2096:HOH:O	1.89	0.72
1:A:32:LEU:CD1	1:A:119:MET:HE2	2.18	0.70
1:A:26:ILE:HD12	1:A:26:ILE:N	2.07	0.69
1:A:110:ILE:O	1:A:110:ILE:HG13	1.91	0.69
1:B:104:ILE:O	1:B:108:GLY:HA2	1.92	0.69
1:B:25:GLU:O	1:B:119:MET:HE3	1.92	0.68
1:B:25:GLU:O	1:B:119:MET:CE	2.44	0.66
1:A:26:ILE:N	1:A:26:ILE:CD1	2.62	0.61
1:A:127:ASP:OD1	4:A:2063:HOH:O	2.15	0.59
1:B:200:GLU:HG2	4:B:2116:HOH:O	2.02	0.59
1:B:91:ILE:HD12	1:B:91:ILE:N	2.19	0.58
1:B:29:LEU:CB	1:B:119:MET:CE	2.80	0.57
1:B:113:SER:HB2	1:B:136:VAL:HG13	1.85	0.57
1:B:29:LEU:CA	1:B:119:MET:CE	2.83	0.56
1:B:107:LEU:O	1:B:139:TYR:OH	2.17	0.56
1:B:23:GLN:HE22	1:B:110:ILE:H	1.54	0.56
1:B:104:ILE:O	1:B:108:GLY:CA	2.54	0.56
1:B:189:HIS:HB3	4:B:2054:HOH:O	2.06	0.56
1:A:26:ILE:CD1	1:A:26:ILE:H	2.21	0.54
1:B:29:LEU:H	1:B:119:MET:HE1	1.67	0.53
1:B:29:LEU:CA	1:B:119:MET:HE2	2.37	0.53
1:B:103:LEU:O	1:B:107:LEU:HB2	2.10	0.52
1:B:29:LEU:N	1:B:119:MET:CE	2.67	0.51
1:A:110:ILE:O	1:A:110:ILE:CG1	2.54	0.50
1:B:130:MET:CE	1:B:133:GLN:HE22	2.25	0.49
1:B:29:LEU:CA	1:B:119:MET:HE1	2.43	0.48
1:B:130:MET:HE3	1:B:133:GLN:HE22	1.78	0.48
1:B:23:GLN:NE2	1:B:109:THR:HB	2.27	0.48
1:A:84:LYS:HA	1:A:198:LEU:HD13	1.98	0.45
1:B:44:PHE:CG	1:B:45:LEU:N	2.84	0.45
1:A:213:PHE:HA	4:A:2107:HOH:O	2.16	0.44
1:B:43:ILE:C	1:B:43:ILE:HD12	2.38	0.44
1:A:203:ILE:HG21	1:A:220:LEU:HD13	2.00	0.43
1:B:25:GLU:O	1:B:119:MET:HE1	2.18	0.43
1:B:23:GLN:HE21	1:B:109:THR:HB	1.83	0.43
1:A:91:ILE:N	1:A:91:ILE:HD12	2.33	0.43
1:B:23:GLN:NE2	1:B:110:ILE:N	2.61	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:PHE:O	1:A:214:ILE:HD13	2.19	0.42
1:B:198:LEU:HD23	1:B:198:LEU:HA	1.72	0.42
1:A:66:ASP:HA	1:A:67:PRO:HD2	1.81	0.41
1:B:131:ILE:CG1	1:B:136:VAL:HG23	2.51	0.41
1:A:116:LYS:O	1:A:120:GLU:HG3	2.21	0.41
1:B:77:HIS:CD2	1:B:77:HIS:N	2.89	0.40
1:B:38:TYR:CZ	1:B:40:ASN:HB3	2.57	0.40
1:B:147:LYS:HE3	1:B:163:GLU:CD	2.42	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1225:DMS:C2	4:A:2118:HOH:O[2_755]	2.00	0.20

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

#### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

#### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry i

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DMS	A	1225	-	3,3,3	2.25	1 (33%)	3,3,3	0.23	0
3	ZZ3	A	1226	-	10,10,10	2.27	1 (10%)	13,13,13	4.79	7 (53%)
3	ZZ3	B	1225	-	10,10,10	2.25	1 (10%)	13,13,13	4.79	7 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ZZ3	A	1226	-	-	1/2/2/2	0/1/1/1
3	ZZ3	B	1225	-	-	2/2/2/2	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1226	ZZ3	C4-S1	-6.86	1.69	1.75
3	B	1225	ZZ3	C4-S1	-6.77	1.70	1.75
2	A	1225	DMS	O-S	3.81	1.76	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1226	ZZ3	C4-N4-C2	11.31	119.99	113.87
3	B	1225	ZZ3	C4-N4-C2	11.30	119.99	113.87
3	B	1225	ZZ3	C6-S1-C4	9.64	109.47	102.27
3	A	1226	ZZ3	C6-S1-C4	9.62	109.45	102.27
3	A	1226	ZZ3	N3-C4-N4	-3.79	120.00	126.95
3	B	1225	ZZ3	N3-C4-N4	-3.79	120.01	126.95
3	A	1226	ZZ3	C2-N1-C3	3.70	120.02	115.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1225	ZZ3	C2-N1-C3	3.69	120.01	115.25
3	A	1226	ZZ3	N1-C2-N4	-3.48	119.96	125.42
3	B	1225	ZZ3	N1-C2-N4	-3.45	120.01	125.42
3	A	1226	ZZ3	C3-N3-C4	3.45	120.02	113.63
3	B	1225	ZZ3	C3-N3-C4	3.44	120.01	113.63
3	B	1225	ZZ3	N3-C3-N1	-3.12	119.99	125.72
3	A	1226	ZZ3	N3-C3-N1	-3.10	120.01	125.72

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1225	ZZ3	N4-C4-S1-C6
3	B	1225	ZZ3	N3-C4-S1-C6
3	A	1226	ZZ3	N4-C4-S1-C6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1225	DMS	0	1

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/236 (88%)	0.49	12 (5%) 23 29	17, 25, 35, 44	0
1	B	209/236 (88%)	0.51	16 (7%) 13 17	16, 25, 38, 44	0
All	All	418/472 (88%)	0.50	28 (6%) 17 22	16, 25, 37, 44	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	GLY	6.0
1	A	176	THR	5.5
1	B	176	THR	5.2
1	B	125	GLY	3.7
1	B	64	LEU	3.7
1	B	66	ASP	3.5
1	A	156	ASP	3.4
1	B	16	GLU	3.3
1	B	123	GLN	3.1
1	A	124	ALA	3.1
1	A	120	GLU	3.1
1	B	68	SER	3.1
1	B	156	ASP	2.9
1	A	68	SER	2.8
1	A	125	GLY	2.8
1	B	91	ILE	2.8
1	A	107	LEU	2.7
1	A	192	GLU	2.5
1	B	65	THR	2.5
1	A	123	GLN	2.3
1	A	48	LEU	2.2
1	B	186	VAL	2.2
1	B	158	GLU	2.2
1	B	111	ALA	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	113	SER	2.1
1	A	174	THR	2.1
1	B	85	GLN	2.1
1	B	17	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
2	DMS	A	1225	4/4	0.72	0.37	55,55,57,58	4
3	ZZ3	B	1225	10/10	0.91	0.18	18,24,34,35	0
3	ZZ3	A	1226	10/10	0.93	0.18	19,23,26,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.