



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 07:36 pm BST

PDB ID : 2XHG  
Title : Crystal Structure of the Epimerization Domain from the Initiation Module of Tyrocidine Biosynthesis  
Authors : Samel, S.-A.; Heine, A.; Marahiel, M.A.; Essen, L.-O.  
Deposited on : 2010-06-15  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYROCIDINE SYNTHETASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	457	3703	2363	628	702	10	0	14	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	441	Total	O	0	0
			441	441		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TYROCIDINE SYNTHETASE A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.71Å 74.59Å 124.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.50 24.64 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.00-1.50) 99.8 (24.64-1.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.160 , 0.179 0.177 , 0.197	Depositor DCC
$R_{free}$ test set	1061 reflections (1.47%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.7	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4175	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/3834	0.59	0/5212

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3703	0	3604	29	0
2	A	30	0	40	2	0
3	A	1	0	0	0	0
4	A	441	0	0	10	0
All	All	4175	0	3644	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ARG:NH1	4:A:2263:HOH:O	2.03	0.92
1:A:332[A]:MET:HE1	1:A:335:ILE:HD11	1.69	0.74
1:A:260[B]:ASN:H	1:A:260[B]:ASN:HD22	1.40	0.67
1:A:368:ASN:OD1	4:A:2370:HOH:O	2.14	0.64
1:A:260[B]:ASN:H	1:A:260[B]:ASN:ND2	1.98	0.62
1:A:117:SER:O	2:A:1465:GOL:H31	2.03	0.58
1:A:409[A]:THR:HG21	4:A:2367:HOH:O	2.04	0.57
1:A:73:GLN:O	1:A:79:VAL:HA	2.05	0.57
1:A:224[A]:VAL:HG13	1:A:227[A]:CYS:SG	2.46	0.55
1:A:416:LEU:HD11	1:A:418:GLU:OE1	2.08	0.53
1:A:224[A]:VAL:CG1	1:A:227[A]:CYS:SG	2.97	0.53
1:A:317:GLN:HE22	1:A:327:LEU:HD13	1.74	0.52
1:A:318:THR:HA	1:A:324[B]:TYR:CD1	2.46	0.51
1:A:250:HIS:N	4:A:2269:HOH:O	2.44	0.51
1:A:73:GLN:NE2	4:A:2079:HOH:O	2.45	0.49
1:A:216:VAL:O	4:A:2240:HOH:O	2.20	0.48
1:A:213:ALA:HA	1:A:279:GLN:HE22	1.78	0.47
1:A:414[A]:THR:HG22	4:A:2405:HOH:O	2.14	0.47
1:A:105:GLN:HG2	1:A:387:TYR:CD1	2.51	0.45
1:A:283:HIS:CD2	1:A:366:THR:HG23	2.52	0.45
1:A:364:GLU:HG3	2:A:1468:GOL:O2	2.17	0.45
1:A:409[A]:THR:HG22	4:A:2401:HOH:O	2.17	0.45
1:A:408:TYR:CD1	1:A:409[A]:THR:HG23	2.53	0.45
1:A:414[A]:THR:HG23	4:A:2407:HOH:O	2.18	0.42
1:A:436:GLU:HA	1:A:439:GLN:HE21	1.85	0.42
1:A:141:PHE:CZ	1:A:143:ALA:HB2	2.54	0.42
1:A:166:ALA:HB2	1:A:381:LEU:HD11	2.02	0.41
1:A:414[A]:THR:CG2	4:A:2407:HOH:O	2.68	0.41
1:A:214:LYS:H	1:A:279:GLN:NE2	2.19	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	469/466 (101%)	460 (98%)	8 (2%)	1 (0%)	47 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	374	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	403/405 (100%)	389 (96%)	14 (4%)	36 9

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	TRP
1	A	83	ASN
1	A	89	GLN
1	A	109[A]	GLU
1	A	109[B]	GLU
1	A	122	GLN
1	A	169	LEU
1	A	200	LEU
1	A	238	ARG
1	A	283	HIS
1	A	317	GLN
1	A	343	ASP
1	A	362	GLN
1	A	380	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	73	GLN
1	A	76	ASN
1	A	89	GLN
1	A	122	GLN
1	A	256	GLN
1	A	279	GLN
1	A	283	HIS
1	A	308	GLN
1	A	317	GLN
1	A	353	ASN
1	A	372	GLN
1	A	439	GLN
1	A	440	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	1468	-	5,5,5	0.39	0	5,5,5	0.20	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	1464	-	5,5,5	0.35	0	5,5,5	0.62	0
2	GOL	A	1467	-	5,5,5	0.39	0	5,5,5	0.76	0
2	GOL	A	1466	-	5,5,5	0.35	0	5,5,5	0.49	0
2	GOL	A	1465	-	5,5,5	0.32	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1468	-	-	0/4/4/4	-
2	GOL	A	1464	-	-	0/4/4/4	-
2	GOL	A	1467	-	-	0/4/4/4	-
2	GOL	A	1466	-	-	0/4/4/4	-
2	GOL	A	1465	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1465	GOL	C1-C2-C3-O3
2	A	1465	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1468	GOL	1	0
2	A	1465	GOL	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	457/466 (98%)	0.33	34 (7%) <b>14</b> <b>15</b>	10, 20, 38, 68	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	373	PHE	16.3
1	A	216	VAL	7.8
1	A	215	ASN	7.7
1	A	463	HIS	6.6
1	A	374	ASP	6.1
1	A	375	ALA	5.3
1	A	317	GLN	4.8
1	A	461	HIS	4.4
1	A	462	HIS	4.2
1	A	295	GLN	4.0
1	A	217	SER	4.0
1	A	377	MET	3.2
1	A	74	HIS	3.2
1	A	7	LYS	3.1
1	A	171	GLY	3.0
1	A	144	ILE	2.9
1	A	362	GLN	2.8
1	A	135	LEU	2.8
1	A	225	THR	2.8
1	A	226	ASP	2.7
1	A	136	HIS	2.6
1	A	250	HIS	2.6
1	A	394	GLY	2.6
1	A	372	GLN	2.6
1	A	396	ASP	2.6
1	A	73	GLN	2.6
1	A	127	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	264[A]	LEU	2.4
1	A	87	GLY	2.4
1	A	376	ASP	2.3
1	A	61	ILE	2.3
1	A	267	LEU	2.2
1	A	68	LEU	2.1
1	A	161	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	A	1469	1/1	0.70	0.21	64,64,64,64	0
2	GOL	A	1468	6/6	0.86	0.21	32,35,35,39	0
2	GOL	A	1467	6/6	0.87	0.16	24,34,38,42	0
2	GOL	A	1465	6/6	0.88	0.23	29,34,38,40	0
2	GOL	A	1466	6/6	0.89	0.10	32,33,35,40	0
2	GOL	A	1464	6/6	0.92	0.10	21,30,31,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.