

# Full wwPDB X-ray Structure Validation Report (i)

#### Apr 29, 2024 – 06:26 pm BST

PDB ID : 2XIT

Title : Crystal structure of monomeric MipZ

Authors: Kiekebusch, D.; Michie, K.A.; Essen, L.O.; Lowe, J.; Thanbichler, M.

Deposited on : 2010-06-30

Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467 Xtriage (Phenix): 1.13

EDS : 2.36.2

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	294	82%	11%	• 6%
1	В	294	82%	12%	• 5%



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called MIPZ.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	277	Total 2114	C 1330	11	O 388	S 7	0	0	1
1	В	279	Total 2167	C 1363	N 398	O 398	S 8	0	4	1

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	ALA	-	expression tag	UNP B8GY04
A	280	VAL	_	expression tag	UNP B8GY04
A	281	ASP	-	expression tag	UNP B8GY04
A	282	LYS	-	expression tag	UNP B8GY04
A	283	LEU	-	expression tag	UNP B8GY04
A	284	ALA	-	expression tag	UNP B8GY04
A	285	ALA	-	expression tag	UNP B8GY04
A	286	ALA	-	expression tag	UNP B8GY04
A	287	LEU	-	expression tag	UNP B8GY04
A	288	GLU	-	expression tag	UNP B8GY04
A	289	HIS	-	expression tag	UNP B8GY04
A	290	HIS	-	expression tag	UNP B8GY04
A	291	HIS	-	expression tag	UNP B8GY04
A	292	HIS	-	expression tag	UNP B8GY04
A	293	HIS	-	expression tag	UNP B8GY04
A	294	HIS	-	expression tag	UNP B8GY04
В	279	ALA	-	expression tag	UNP B8GY04
В	280	VAL	-	expression tag	UNP B8GY04
В	281	ASP	-	expression tag	UNP B8GY04
В	282	LYS	-	expression tag	UNP B8GY04
В	283	LEU	-	expression tag	UNP B8GY04
В	284	ALA	-	expression tag	UNP B8GY04
В	285	ALA	-	expression tag	UNP B8GY04
В	286	ALA	-	expression tag	UNP B8GY04
В	287	LEU	-	expression tag	UNP B8GY04

Continued on next page...



### $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	288	GLU	-	expression tag	UNP B8GY04
В	289	HIS	-	expression tag	UNP B8GY04
В	290	HIS	-	expression tag	UNP B8GY04
В	291	HIS	-	expression tag	UNP B8GY04
В	292	HIS	-	expression tag	UNP B8GY04
В	293	HIS	-	expression tag	UNP B8GY04
В	294	HIS	-	expression tag	UNP B8GY04


## • Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	187	Total O 187 187	0	0
2	В	233	Total O 233 233	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	124.67Å 124.67Å 239.05Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	49.27 - 1.80	Depositor
Resolution (A)	49.20 - 1.80	EDS
% Data completeness	99.9 (49.27-1.80)	Depositor
(in resolution range)	99.8 (49.20-1.80)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.55 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
D D.	0.184 , 0.230	Depositor
$R, R_{free}$	0.209 , $0.250$	DCC
$R_{free}$ test set	1215  reflections  (1.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.0	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 45.7	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4701	wwPDB-VP
Average B, all atoms $(Å^2)$	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
WIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.52	0/2146	0.67	1/2906 (0.0%)	
1	В	0.56	0/2213	0.71	0/2995	
All	All	0.54	0/4359	0.69	1/5901 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	133	MET	CG-SD-CE	-5.46	91.47	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Group
1	A	272	GLU	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2135	29	1
1	В	2167	0	2211	44	0
2	A	187	0	0	1	0
2	В	233	0	0	8	0
All	All	4701	0	4346	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:2:ALA:HA	1:A:177:GLN:HE22	1.23	1.03
1:B:149:VAL:O	1:B:150:THR:HG23	1.64	0.96
1:A:272:GLU:HA	1:A:272:GLU:OE2	1.75	0.85
1:B:179[A]:MET:SD	2:B:2095:HOH:O	2.35	0.84
1:B:4:THR:N	2:B:2002:HOH:O	2.10	0.82
1:B:247:SER:H	1:B:250:HIS:HD2	1.32	0.75
1:A:146:VAL:HG12	1:A:146:VAL:O	1.87	0.74
1:B:273:THR:O	2:B:2230:HOH:O	2.08	0.72
1:A:2:ALA:HA	1:A:177:GLN:NE2	2.03	0.68
1:A:273:THR:HG23	2:A:2064:HOH:O	1.94	0.68
1:B:146:VAL:HG12	1:B:148:PRO:HD2	1.75	0.66
1:B:86:GLU:OE2	2:B:2071:HOH:O	2.14	0.66
1:A:12:GLU:CB	1:A:142:MET:HG3	2.25	0.65
1:B:247:SER:H	1:B:250:HIS:CD2	2.13	0.65
1:A:133:MET:HE2	1:A:143:LEU:HD11	1.78	0.64
1:B:146:VAL:HG12	1:B:148:PRO:CD	2.27	0.64
1:B:116:SER:OG	1:B:119[A]:THR:HG23	1.97	0.64
1:A:5:ARG:HD2	1:A:105:ASP:O	1.98	0.64
1:A:257:LEU:HB3	1:A:273:THR:HG21	1.80	0.63
1:A:133:MET:CE	1:A:143:LEU:HD11	2.29	0.62
1:B:146:VAL:O	1:B:148:PRO:HD2	1.99	0.62
1:B:255:GLN:HE21	1:B:258:ARG:HH11	1.48	0.62
1:B:125:ARG:NH1	2:B:2097:HOH:O	2.24	0.61
1:B:168:LYS:HG2	1:B:172:LEU:HD13	1.82	0.61
1:A:146:VAL:O	1:A:146:VAL:CG1	2.50	0.60
1:A:247:SER:H	1:A:250:HIS:HD2	1.51	0.58
1:B:147:ASP:CB	1:B:149:VAL:HG23	2.32	0.58
1:A:279:ALA:O	1:A:280:VAL:C	2.41	0.58
1:B:148:PRO:HD3	1:B:151:LEU:HA	1.86	0.57
1:A:133:MET:CE	1:A:143:LEU:CD1	2.83	0.56
		Continue	ed on nert nage

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap(Å)
1:B:273:THR:C	2:B:2230:HOH:O	2.45	0.55
1:B:146:VAL:C	1:B:148:PRO:HD2	2.28	0.54
1:B:258:ARG:HE	1:B:273:THR:HB	1.72	0.54
1:B:147:ASP:C	1:B:149:VAL:N	2.61	0.53
1:A:247:SER:H	1:A:250:HIS:CD2	2.26	0.52
1:B:147:ASP:CA	1:B:149:VAL:HG23	2.40	0.51
1:A:145:THR:HG22	1:A:146:VAL:HB	1.91	0.50
1:A:221:ARG:HH11	1:A:249:GLN:NE2	2.10	0.50
1:B:24:HIS:HE1	1:B:234:ILE:H	1.58	0.50
1:B:222:VAL:HG13	1:B:225:ARG:HE	1.77	0.50
1:B:86:GLU:HG2	2:B:2071:HOH:O	2.11	0.50
1:A:101:MET:SD	1:A:107:ILE:HD12	2.53	0.48
1:A:146:VAL:O	1:A:152:GLU:O	2.31	0.48
1:B:255:GLN:NE2	1:B:258:ARG:HH11	2.10	0.47
1:B:46:ARG:NH1	2:B:2030:HOH:O	2.47	0.47
1:A:31:TYR:HB3	1:A:271:GLY:HA3	1.97	0.47
1:A:5:ARG:CD	1:A:105:ASP:O	2.61	0.46
1:B:24:HIS:CE1	1:B:234:ILE:H	2.34	0.46
1:B:147:ASP:O	1:B:148:PRO:C	2.55	0.46
1:A:163:VAL:HG13	1:A:179:MET:HE3	1.97	0.45
1:B:9:VAL:HG12	1:B:18:LYS:HG3	1.99	0.44
1:B:147:ASP:O	1:B:149:VAL:HG23	2.18	0.44
1:B:146:VAL:HG12	1:B:148:PRO:HD3	1.99	0.44
1:B:149:VAL:O	1:B:150:THR:CG2	2.52	0.44
1:B:24:HIS:HE1	1:B:234:ILE:HG12	1.83	0.44
1:B:25:LEU:CD2	1:B:261:MET:SD	3.06	0.44
1:B:147:ASP:C	1:B:149:VAL:HG23	2.39	0.43
1:A:24:HIS:HE1	1:A:234:ILE:HG12	1.83	0.43
1:B:101:MET:SD	1:B:107:ILE:HD12	2.59	0.43
1:B:221:ARG:HH11	1:B:249:GLN:NE2	2.17	0.43
1:B:258:ARG:NE	1:B:273:THR:HB	2.32	0.43
1:B:146:VAL:HG23	1:B:206:LEU:HD23	2.02	0.42
1:B:258:ARG:HH21	1:B:273:THR:CG2	2.32	0.42
1:A:24:HIS:CE1	1:A:234:ILE:H	2.38	0.42
1:B:123:HIS:HE1	1:B:159:TYR:OH	2.03	0.42
1:A:221:ARG:HH11	1:A:249:GLN:HE22	1.68	0.42
1:A:20:THR:O	1:A:24:HIS:HD2	2.03	0.41
1:A:9:VAL:HG12	1:A:18:LYS:HG3	2.02	0.41
1:A:5:ARG:NH2	1:A:104:CYS:O	2.54	0.41
1:A:59:LEU:HD21	1:A:66:LEU:HD12	2.01	0.41
1:B:258:ARG:HH21	1:B:273:THR:HG21	1.86	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:B:147:ASP:CB	1:B:149:VAL:CG2	2.98	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:2:ALA:CB	1:A:2:ALA:CB[17_555]	1.96	0.24

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/294 (92%)	264 (97%)	5 (2%)	2 (1%)	22 10
1	В	279/294~(95%)	272 (98%)	4 (1%)	3 (1%)	14 4
All	All	550/588~(94%)	536 (98%)	9 (2%)	5 (1%)	17 6

#### All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	VAL
1	В	148	PRO
1	В	150	THR
1	A	272	GLU
1	В	272	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	216/238 (91%)	213 (99%)	3 (1%)	67 59
1	В	$227/238 \ (95\%)$	225 (99%)	2 (1%)	78 75
All	All	443/476 (93%)	438 (99%)	5 (1%)	73 68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	242	ARG
1	A	272	GLU
1	A	273	THR
1	В	5	ARG
1	В	272	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	177	GLN
1	A	249	GLN
1	A	250	HIS
1	В	24	HIS
1	В	123	HIS
1	В	195	ASN
1	В	249	GLN
1	В	250	HIS
1	В	255	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

There are no ligands in this entry.

# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	277/294 (94%)	0.23	14 (5%) 28 22	14, 25, 42, 57	4 (1%)
1	В	279/294~(94%)	0.06	8 (2%) 51 46	11, 20, 37, 55	0
All	All	556/588 (94%)	0.14	22 (3%) 38 32	11, 23, 40, 57	4 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	149	VAL	7.6
1	A	273	THR	6.5
1	A	193	ALA	5.5
1	A	194	ARG	5.2
1	A	2	ALA	5.0
1	A	192	GLU	4.8
1	A	279	ALA	4.4
1	A	280	VAL	4.2
1	В	273	THR	4.2
1	A	64	ILE	3.9
1	A	288	GLU	3.8
1	A	195	ASN	3.8
1	A	146	VAL	3.3
1	В	113	GLY	2.7
1	A	189	ALA	2.7
1	A	65	GLU	2.6
1	В	148	PRO	2.5
1	В	150	THR	2.3
1	В	288	GLU	2.3
1	В	114	GLY	2.2
1	В	272	GLU	2.2
1	A	137	PHE	2.1



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.4 Ligands (i)

There are no ligands in this entry.

### 6.5 Other polymers (i)

There are no such residues in this entry.

