



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:06 PM UTC

PDB ID : 2XXZ / pdb_00002xxz
Title : Crystal structure of the human JMJD3 jumonji domain
Authors : Che, K.H.; Yue, W.W.; Krojer, T.; Muniz, J.R.C.; Ng, S.S.; Tumber, A.; Daniel, M.; Burgess-Brown, N.; Savitsky, P.; Ugochukwu, E.; Filippakopoulos, P.; Arrowsmith, C.; Weigelt, J.; Edwards, A.; Bountra, C.; Oppermann, U.
Deposited on : 2010-11-12
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	 5% 78% 10% 12%
1	B	332	 7% 75% 10% 14%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSINE-SPECIFIC DEMETHYLASE 6B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	292	2400	1530	415	444	11	0	11	0
1	B	286	2311	1479	400	422	10	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1171	SER	-	expression tag	UNP O15054
A	1172	MET	-	expression tag	UNP O15054
B	1171	SER	-	expression tag	UNP O15054
B	1172	MET	-	expression tag	UNP O15054

- Molecule 2 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ni	0	0
			2	2		
2	B	1	Total	Ni	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is 8-hydroxyquinoline-5-carboxylic acid (CCD ID: 8XQ) (formula: C₁₀H₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	10	1	3		
5	B	1	Total	C	N	O	0	0
			14	10	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	164	Total	O	0	0
			164	164		
6	B	155	Total	O	0	0
			155	155		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LYSINE-SPECIFIC DEMETHYLASE 6B



- Molecule 1: LYSINE-SPECIFIC DEMETHYLASE 6B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.76Å 71.43Å 159.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.16 – 1.80 29.16 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.16-1.80) 99.7 (29.16-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.183 , 0.216 0.213 , 0.245	Depositor DCC
R_{free} test set	3058 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5082	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, NA, EDO, 8XQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	2/2480 (0.1%)	0.84	3/3386 (0.1%)
1	B	0.80	5/2390 (0.2%)	0.84	5/3264 (0.2%)
All	All	0.81	7/4870 (0.1%)	0.84	8/6650 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1272	THR	C-O	-7.01	1.19	1.24
1	B	1248	GLN	CD-OE1	5.67	1.34	1.23
1	A	1268	SER	C-N	-5.43	1.26	1.33
1	B	1248	GLN	CD-NE2	-5.34	1.22	1.33
1	B	1238	HIS	ND1-CE1	5.31	1.37	1.32
1	B	1271	HIS	ND1-CE1	5.29	1.37	1.32
1	A	1400	ILE	CA-CB	5.09	1.60	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1248	GLN	N-CA-C	7.47	114.72	108.07
1	A	1321	HIS	CB-CG-CD2	-6.67	122.53	131.20
1	B	1238	HIS	CB-CG-CD2	-6.61	122.61	131.20
1	B	1271	HIS	CB-CG-CD2	-6.42	122.85	131.20
1	B	1238	HIS	CB-CG-ND1	5.78	131.38	122.70
1	A	1321	HIS	CB-CG-ND1	5.67	131.20	122.70
1	A	1355	SER	CB-CA-C	-5.51	110.23	116.63
1	B	1271	HIS	CB-CG-ND1	5.50	130.96	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2400	0	2313	26	0
1	B	2311	0	2222	35	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	16	0	24	0	0
4	B	4	0	6	3	0
5	A	14	0	5	0	0
5	B	14	0	5	0	0
6	A	164	0	0	3	0
6	B	155	0	0	4	1
All	All	5082	0	4575	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1251:ASP:OD1	1:B:1438:LEU:HD12	1.73	0.89
1:A:1329:ILE:CD1	1:A:1377:MET:HE2	2.04	0.88
1:B:1191:ARG:HH11	4:B:2501:EDO:H12	1.40	0.87
1:B:1251:ASP:OD1	1:B:1438:LEU:CD1	2.26	0.82
1:A:1329:ILE:HD13	1:A:1377:MET:HE2	1.68	0.75
1:A:1241:GLU:OE1	6:A:2034:HOH:O	2.10	0.70
1:A:1243[B]:ARG:HD3	1:A:1269:ARG:HG2	1.74	0.70
1:B:1240:VAL:CG2	1:B:1274:ILE:HD13	2.23	0.69
1:B:1395:SER:HB3	1:B:1461:ILE:HD13	1.76	0.68
1:B:1329:ILE:CD1	1:B:1377[A]:MET:HE2	2.26	0.65
1:A:1329:ILE:HD11	1:A:1377:MET:HE2	1.77	0.64
1:B:1249:PRO:HG2	1:B:1252:GLU:OE1	1.97	0.64
1:B:1353:VAL:CG1	1:B:1367:ILE:HG23	2.29	0.63
1:A:1243[B]:ARG:CD	1:A:1269:ARG:HG2	2.31	0.60
1:B:1251:ASP:OD1	1:B:1438:LEU:HD11	2.02	0.59
1:A:1242:VAL:HG21	1:A:1272:THR:HG22	1.85	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1320:HIS:ND1	1:A:1320:HIS:C	2.61	0.58
1:B:1212:ILE:HD12	1:B:1458:LEU:HD22	1.86	0.57
1:B:1248:GLN:HB2	1:B:1249:PRO:HD2	1.87	0.56
1:B:1240:VAL:HG21	1:B:1274:ILE:HD13	1.87	0.56
1:A:1428:ASP:OD2	1:A:1431:THR:OG1	2.15	0.55
1:A:1242:VAL:CG2	1:A:1272:THR:HG22	2.36	0.55
1:B:1250:SER:HA	1:B:1470:GLN:OE1	2.07	0.54
1:B:1267:SER:HB3	6:B:2040:HOH:O	2.07	0.53
1:A:1243[A]:ARG:HD3	1:A:1269:ARG:HG2	1.89	0.53
1:B:1329:ILE:HD11	1:B:1377[A]:MET:HE2	1.89	0.53
1:B:1322:ILE:HD12	1:B:1322:ILE:O	2.09	0.53
1:B:1248:GLN:HB2	1:B:1249:PRO:CD	2.40	0.52
1:B:1191:ARG:NH1	4:B:2501:EDO:H12	2.18	0.52
1:A:1226:PHE:CE1	1:A:1478[B]:ILE:HD12	2.44	0.52
1:A:1243[B]:ARG:NH1	1:A:1269:ARG:HD2	2.25	0.51
1:B:1251:ASP:OD2	1:B:1450:ARG:NH1	2.45	0.50
1:A:1412:GLU:OE2	6:A:2120:HOH:O	2.19	0.48
1:B:1250:SER:HA	1:B:1383:ARG:HH21	1.76	0.48
1:B:1242:VAL:CG2	1:B:1272:THR:HG22	2.43	0.48
1:A:1353[A]:VAL:HG22	1:A:1480:TRP:CG	2.49	0.48
1:A:1344:LEU:HD11	1:A:1478[B]:ILE:HD11	1.96	0.48
1:B:1190:LYS:NZ	4:B:2501:EDO:O2	2.42	0.48
1:B:1233:GLU:HG3	6:B:2034:HOH:O	2.14	0.48
1:A:1353[A]:VAL:HG23	6:A:2069:HOH:O	2.13	0.47
1:B:1322:ILE:HD12	1:B:1322:ILE:C	2.39	0.46
1:B:1246:VAL:HG12	6:B:2048:HOH:O	2.15	0.46
1:B:1329:ILE:HD13	1:B:1377[A]:MET:HE2	1.97	0.46
1:A:1242:VAL:HG21	1:A:1272:THR:CG2	2.47	0.45
1:A:1290:GLN:HG2	1:A:1291:GLU:N	2.31	0.45
1:A:1320:HIS:ND1	1:A:1321:HIS:N	2.65	0.45
1:A:1244:THR:OG1	1:A:1324:LYS:HD3	2.16	0.45
1:B:1252:GLU:HB3	1:B:1254:TRP:NE1	2.32	0.44
1:A:1343:GLU:OE2	1:A:1346:LYS:NZ	2.45	0.44
1:B:1272:THR:OG1	1:B:1273:THR:N	2.51	0.44
1:B:1231:LEU:HD13	1:B:1375:LEU:HD21	1.99	0.43
1:B:1353:VAL:CG1	1:B:1367:ILE:CG2	2.95	0.43
1:B:1240:VAL:CG2	1:B:1274:ILE:CD1	2.96	0.42
1:A:1243[B]:ARG:NH1	1:A:1269:ARG:CG	2.83	0.41
1:A:1335[B]:LYS:HA	1:A:1335[B]:LYS:HD3	1.74	0.41
1:A:1219:LEU:CD1	1:A:1347:LEU:HD23	2.50	0.41
1:B:1242:VAL:HG23	1:B:1272:THR:HG22	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1353:VAL:HG11	1:B:1367:ILE:CG2	2.50	0.41
1:A:1362:HIS:C	1:A:1363:VAL:HG22	2.45	0.41
1:B:1267:SER:CB	6:B:2040:HOH:O	2.67	0.41
1:B:1353:VAL:HG12	1:B:1367:ILE:HG23	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:2058:HOH:O	6:B:2100:HOH:O[3_754]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/332 (90%)	293 (98%)	6 (2%)	0	100	100
1	B	284/332 (86%)	279 (98%)	5 (2%)	0	100	100
All	All	583/664 (88%)	572 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/295 (90%)	262 (99%)	2 (1%)	73	70
1	B	250/295 (85%)	246 (98%)	4 (2%)	55	47
All	All	514/590 (87%)	508 (99%)	6 (1%)	63	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1209	ILE
1	A	1229	LYS
1	B	1209	ILE
1	B	1251	ASP
1	B	1287	GLU
1	B	1482	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1425	HIS
1	A	1477	ASN
1	A	1491	GLN
1	B	1238	HIS
1	B	1279	GLN
1	B	1374	GLN
1	B	1388	GLN
1	B	1390	ASN
1	B	1477	ASN
1	B	1481	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	8XQ	B	3001	2	15,15,15	1.20	1 (6%)	21,21,21	1.16	2 (9%)
4	EDO	A	2496	-	3,3,3	0.51	0	2,2,2	0.49	0
4	EDO	B	2501	-	3,3,3	0.29	0	2,2,2	0.62	0
4	EDO	A	2498	-	3,3,3	0.50	0	2,2,2	0.23	0
4	EDO	A	2499	-	3,3,3	0.34	0	2,2,2	0.15	0
4	EDO	A	2497	-	3,3,3	0.34	0	2,2,2	0.40	0
5	8XQ	A	3001	2	15,15,15	1.21	1 (6%)	21,21,21	1.49	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	8XQ	B	3001	2	-	0/4/4/4	0/2/2/2
4	EDO	A	2496	-	-	0/1/1/1	-
4	EDO	B	2501	-	-	1/1/1/1	-
4	EDO	A	2498	-	-	1/1/1/1	-
4	EDO	A	2499	-	-	0/1/1/1	-
4	EDO	A	2497	-	-	0/1/1/1	-
5	8XQ	A	3001	2	-	0/4/4/4	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	3001	8XQ	CAN-NAI	-2.41	1.33	1.37
5	B	3001	8XQ	CAN-NAI	-2.19	1.34	1.37

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	3001	8XQ	CAE-NAI-CAN	4.42	122.52	117.31
5	B	3001	8XQ	CAE-NAI-CAN	3.13	121.00	117.31
5	B	3001	8XQ	OAB-CAJ-CAL	-2.43	116.00	122.31
5	A	3001	8XQ	OAB-CAJ-CAL	-2.30	116.35	122.31

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2501	EDO	O1-C1-C2-O2
4	A	2498	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2501	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/332 (87%)	0.22	15 (5%) 33 32	9, 24, 51, 69	11 (3%)
1	B	286/332 (86%)	0.46	22 (7%) 19 17	15, 31, 59, 74	4 (1%)
All	All	578/664 (87%)	0.34	37 (6%) 25 24	9, 28, 58, 74	15 (2%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1251	ASP	5.1
1	A	1256	LEU	5.0
1	B	1499	TRP	4.0
1	B	1320	HIS	3.5
1	B	1369	GLY	3.4
1	B	1250	SER	3.3
1	B	1249	PRO	3.1
1	B	1269	ARG	3.1
1	B	1356	THR	3.1
1	B	1368	LEU	2.9
1	B	1256	LEU	2.9
1	B	1371	ASN	2.7
1	B	1321	HIS	2.6
1	A	1251	ASP	2.6
1	B	1268	SER	2.5
1	A	1320	HIS	2.5
1	A	1321	HIS	2.5
1	B	1497	TYR	2.5
1	A	1493	ALA	2.5
1	A	1257	THR	2.4
1	A	1335[A]	LYS	2.4
1	B	1367	ILE	2.4
1	A	1358	ASN	2.4
1	B	1175	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1292	GLU	2.3
1	A	1249	PRO	2.2
1	B	1488	TYR	2.2
1	B	1289	LEU	2.2
1	A	1254	TRP	2.1
1	B	1290	GLN	2.1
1	A	1325	PHE	2.0
1	B	1370	MET	2.0
1	A	1430	LEU	2.0
1	A	1243[A]	ARG	2.0
1	B	1274	ILE	2.0
1	B	1322	ILE	2.0
1	A	1260	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	A	2498	4/4	0.83	0.19	29,45,50,53	0
4	EDO	A	2496	4/4	0.87	0.12	29,31,33,34	0
4	EDO	B	2501	4/4	0.88	0.15	25,26,29,30	0
3	NA	A	2495	1/1	0.91	0.16	37,37,37,37	0
2	NI	A	2500	1/1	0.91	0.06	33,33,33,33	1
4	EDO	A	2499	4/4	0.93	0.09	21,22,27,28	0
4	EDO	A	2497	4/4	0.95	0.07	25,26,27,28	0
5	8XQ	B	3001	14/14	0.95	0.06	18,21,26,28	0
5	8XQ	A	3001	14/14	0.97	0.05	12,18,20,21	0
2	NI	A	2494	1/1	0.98	0.04	20,20,20,20	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NI	B	2500	1/1	0.99	0.05	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.