



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2021 – 10:09 PM EDT

PDB ID : 2YXV
Title : The deletion mutant of Multicopper Oxidase CueO
Authors : Higuchi, Y.; Komori, H.
Deposited on : 2007-04-27
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	
1	B	446	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue copper oxidase cueO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	444	3404	2162	601	623	18	0	0	0
1	B	440	3364	2138	589	619	18	0	0	0

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	357	GLY	PRO	engineered mutation	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	LEU	deletion	UNP P36649
A	?	-	ASP	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	GLN	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	LEU	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	GLU	deletion	UNP P36649
A	?	-	LYS	deletion	UNP P36649
A	?	-	TYR	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	ASP	deletion	UNP P36649
A	?	-	GLN	deletion	UNP P36649
A	?	-	ALA	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	ALA	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	ASP	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP P36649
A	?	-	GLN	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	ASN	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	ASN	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649
A	?	-	MET	deletion	UNP P36649
A	?	-	ASN	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	GLY	deletion	UNP P36649
A	?	-	LYS	deletion	UNP P36649
A	?	-	PHE	deletion	UNP P36649
A	?	-	ASP	deletion	UNP P36649
A	?	-	PHE	deletion	UNP P36649
A	?	-	HIS	deletion	UNP P36649
A	406	GLY	HIS	engineered mutation	UNP P36649
A	517	GLY	-	expression tag	UNP P36649
A	518	HIS	-	expression tag	UNP P36649
A	519	HIS	-	expression tag	UNP P36649
A	520	HIS	-	expression tag	UNP P36649
A	521	HIS	-	expression tag	UNP P36649
A	522	HIS	-	expression tag	UNP P36649
B	357	GLY	PRO	engineered mutation	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	LEU	deletion	UNP P36649
B	?	-	ASP	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	GLN	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	LEU	deletion	UNP P36649

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P36649
B	?	-	GLU	deletion	UNP P36649
B	?	-	LYS	deletion	UNP P36649
B	?	-	TYR	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	ASP	deletion	UNP P36649
B	?	-	GLN	deletion	UNP P36649
B	?	-	ALA	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	ALA	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	ASP	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	?	-	SER	deletion	UNP P36649
B	?	-	GLN	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	ASN	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	ASN	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	?	-	MET	deletion	UNP P36649
B	?	-	ASN	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	GLY	deletion	UNP P36649
B	?	-	LYS	deletion	UNP P36649
B	?	-	PHE	deletion	UNP P36649
B	?	-	ASP	deletion	UNP P36649
B	?	-	PHE	deletion	UNP P36649
B	?	-	HIS	deletion	UNP P36649
B	406	GLY	HIS	engineered mutation	UNP P36649
B	517	GLY	-	expression tag	UNP P36649
B	518	HIS	-	expression tag	UNP P36649
B	519	HIS	-	expression tag	UNP P36649

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	520	HIS	-	expression tag	UNP P36649
B	521	HIS	-	expression tag	UNP P36649
B	522	HIS	-	expression tag	UNP P36649

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cu 2 2	0	0
2	B	2	Total Cu 2 2	0	0

- Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N O 4 1 3	0	0

- Molecule 4 is CU-O-CU LINKAGE (three-letter code: C2O) (formula: Cu₂O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Cu	O	0	0
			3	2	1		
4	B	1	Total	Cu	O	0	0
			3	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	178	Total	O	0	0
			178	178		
5	B	217	Total	O	0	0
			217	217		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blue copper oxidase cueO



- Molecule 1: Blue copper oxidase cueO



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.98Å 50.95Å 86.05Å 83.20° 90.83° 66.66°	Depositor
Resolution (Å)	24.40 – 1.81 24.24 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.7 (24.40-1.81) 92.3 (24.24-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.171 , 0.208 0.180 , 0.214	Depositor DCC
R_{free} test set	3348 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtrriage
Anisotropy	0.589	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7177	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, C2O, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3492	0.61	0/4753
1	B	0.41	0/3448	0.63	0/4693
All	All	0.41	0/6940	0.62	0/9446

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	30	GLU	CA

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3404	0	3379	13	0
1	B	3364	0	3351	13	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	4	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	A	178	0	0	2	0
5	B	217	0	0	0	0
All	All	7177	0	6730	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:519:HIS:CE1	5:A:884:HOH:O	2.41	0.71
1:B:175:GLN:HG3	1:B:178:ILE:HD12	1.77	0.66
1:A:519:HIS:NE2	1:A:521:HIS:ND1	2.50	0.60
1:A:141:HIS:HB2	1:A:142:PRO:HD2	1.84	0.60
1:B:48:ILE:HD12	1:B:86:VAL:HG11	1.84	0.59
1:B:175:GLN:CG	1:B:178:ILE:HD12	2.39	0.53
1:A:454:ILE:HG21	1:A:457:GLU:HB2	1.93	0.50
1:B:40:LEU:HD21	1:B:48:ILE:HG23	1.94	0.49
1:B:141:HIS:HB2	1:B:142:PRO:HD2	1.96	0.48
1:A:204:THR:HG22	1:A:504:GLU:HG2	1.96	0.48
1:B:410:ILE:HG21	1:B:512:LEU:HD23	1.96	0.47
1:A:289:ASP:O	1:A:321:PRO:HB2	2.15	0.47
1:A:64:ALA:HB2	1:A:194:GLY:O	2.16	0.45
1:B:204:THR:HG22	1:B:504:GLU:HG2	1.98	0.45
1:A:119:ILE:HG12	1:A:125:ARG:HG3	1.99	0.45
1:A:337:LEU:HD21	1:A:468:GLY:HA2	1.99	0.43
1:A:512:LEU:HD12	1:A:512:LEU:C	2.40	0.42
1:A:350:LYS:C	1:A:351:LEU:HD12	2.39	0.42
1:A:519:HIS:HE1	5:A:884:HOH:O	1.94	0.42
1:B:100:LEU:C	1:B:100:LEU:HD23	2.41	0.41
1:B:80:LEU:HB3	1:B:131:VAL:HG21	2.01	0.41
1:B:64:ALA:HB2	1:B:194:GLY:O	2.20	0.41
1:B:343:LEU:N	1:B:343:LEU:HD12	2.36	0.41
1:B:264:LEU:O	1:B:334:LEU:HB2	2.21	0.40
1:A:351:LEU:HD12	1:A:351:LEU:N	2.37	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:PHE:HB3	1:B:484:VAL:HG12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/446 (99%)	428 (97%)	14 (3%)	0	100	100
1	B	438/446 (98%)	423 (97%)	15 (3%)	0	100	100
All	All	880/892 (99%)	851 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/365 (100%)	359 (99%)	5 (1%)	67	58
1	B	360/365 (99%)	355 (99%)	5 (1%)	67	58
All	All	724/730 (99%)	714 (99%)	10 (1%)	67	58

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	HIS
1	A	280	ARG
1	A	325	SER
1	A	419	LYS
1	A	492	LYS
1	B	30	GLU
1	B	110	GLU
1	B	143	HIS
1	B	169	LYS
1	B	280	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	B	81	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	C2O	B	702	1	0,2,2	-	-	-	-	-
4	C2O	A	702	1	0,2,2	-	-	-	-	-
3	NO3	A	706	-	1,3,3	3.13	1 (100%)	0,3,3	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	706	NO3	O1-N	3.13	1.38	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/446 (99%)	0.51	35 (7%) 12 9	20, 25, 33, 37	1 (0%)
1	B	440/446 (98%)	0.37	28 (6%) 19 15	21, 25, 32, 35	1 (0%)
All	All	884/892 (99%)	0.44	63 (7%) 16 12	20, 25, 33, 37	2 (0%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	44	ALA	7.9
1	B	406	GLY	7.3
1	A	357	GLY	7.2
1	B	290	ASN	7.0
1	A	289	ASP	6.5
1	B	356	ASP	6.1
1	A	356	ASP	6.0
1	A	518	HIS	5.8
1	B	357	GLY	5.3
1	A	45	ARG	5.2
1	A	520	HIS	5.0
1	A	406	GLY	5.0
1	A	171	MET	4.7
1	B	407	ALA	4.3
1	B	437	VAL	4.2
1	B	44	ALA	4.1
1	B	289	ASP	4.0
1	A	407	ALA	3.9
1	A	521	HIS	3.7
1	A	519	HIS	3.7
1	A	421	MET	3.6
1	B	288	ASN	3.4
1	B	291	LYS	3.4
1	A	413	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	327	SER	3.3
1	A	418	ASN	3.3
1	B	440	MET	3.2
1	A	199	GLN	3.1
1	A	416	ASP	3.1
1	B	199	GLN	3.0
1	A	290	ASN	3.0
1	A	336	SER	2.8
1	A	291	LYS	2.8
1	A	447	ILE	2.8
1	A	337	LEU	2.8
1	B	418	ASN	2.8
1	B	30	GLU	2.7
1	B	326	ALA	2.7
1	B	61	GLY	2.6
1	B	231	LEU	2.6
1	B	233	LEU	2.6
1	A	145	HIS	2.6
1	A	95	THR	2.6
1	A	417	MET	2.5
1	A	303	MET	2.5
1	B	303	MET	2.4
1	A	333	THR	2.4
1	B	171	MET	2.4
1	A	445	PHE	2.3
1	B	502	LEU	2.3
1	B	498	ALA	2.3
1	A	344	GLU	2.3
1	A	47	ARG	2.2
1	B	145	HIS	2.2
1	A	43	ASP	2.2
1	B	63	THR	2.2
1	B	192	ALA	2.2
1	A	287	VAL	2.1
1	B	239	CYS	2.1
1	B	344	GLU	2.0
1	A	49	GLN	2.0
1	A	110	GLU	2.0
1	B	447	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NO3	A	706	4/4	0.95	0.15	26,27,27,27	0
2	CU	B	701	1/1	0.99	0.08	25,25,25,25	0
2	CU	B	703	1/1	0.99	0.02	38,38,38,38	0
2	CU	A	701	1/1	0.99	0.08	26,26,26,26	0
4	C2O	A	702	3/3	0.99	0.06	29,29,30,33	0
4	C2O	B	702	3/3	0.99	0.07	29,29,32,33	0
2	CU	A	703	1/1	1.00	0.05	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.