



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 12:06 PM EDT

PDB ID : 2ZIE
Title : Crystal Structure of TTHA0409, Putative DNA Modification Methylase from *Thermus thermophilus* HB8- Selenomethionine derivative
Authors : Morita, R.; Ishikawa, H.; Nakagawa, N.; Masui, R.; Yokoyama, S.; Kuramitsu, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2008-02-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	 5% 51% 24% 22%
1	B	297	 2% 53% 22% 22%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative modification methylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	233	1910	1239	345	323	1	2	0	0	0
1	B	232	1894	1227	340	324	1	2	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	31	Total	O	0	0
			31	31		
2	B	43	Total	O	0	0
			43	43		


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative modification methylase



- Molecule 1: Putative modification methylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.58Å 58.25Å 80.91Å 90.00° 105.52° 90.00°	Depositor
Resolution (Å)	19.92 – 2.50 19.92 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.2 (19.92-2.50) 97.0 (19.92-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	18.04 (at 2.50Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.222 , 0.254 0.215 , 0.251	Depositor DCC
R_{free} test set	1909 reflections (9.84%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3878	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1959	0.66	1/2650 (0.0%)
1	B	0.40	0/1943	0.65	0/2630
All	All	0.40	0/3902	0.66	1/5280 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	LEU	CA-CB-CG	5.88	128.81	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1910	0	1919	76	0
1	B	1894	0	1891	75	0
2	A	31	0	0	0	0
2	B	43	0	0	1	0
All	All	3878	0	3810	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (141) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:248:THR:HG22	1:B:276:ARG:HH22	0.92	1.02
1:B:248:THR:HG22	1:B:276:ARG:NH2	1.76	1.01
1:B:228:LEU:HA	1:B:231:MSE:HE2	1.42	0.99
1:B:22:His:HD2	1:B:260:ALA:H	1.18	0.90
1:A:42:His:HD2	1:A:238:VAL:H	1.16	0.89
1:B:42:His:HD2	1:B:238:VAL:H	1.16	0.88
1:A:103:ALA:HB2	1:A:110:His:ND1	1.91	0.85
1:B:228:LEU:HA	1:B:231:MSE:CE	2.06	0.85
1:B:228:LEU:HD23	1:B:231:MSE:CE	2.08	0.84
1:B:66:ILE:HD12	1:B:66:ILE:H	1.43	0.81
1:B:96:VAL:HG21	1:B:220:PHE:CD2	2.16	0.80
1:A:42:His:CD2	1:A:238:VAL:H	2.02	0.78
1:B:228:LEU:HD23	1:B:231:MSE:HE1	1.67	0.77
1:B:248:THR:CG2	1:B:276:ARG:HH22	1.86	0.77
1:B:172:MSE:HE1	1:B:232:PHE:CZ	2.24	0.73
1:A:180:ARG:HG2	1:A:180:ARG:HH21	1.52	0.73
1:A:83:VAL:HA	1:A:86:LEU:HD11	1.71	0.72
1:A:248:THR:HB	1:A:276:ARG:HH12	1.54	0.71
1:A:95:ILE:HG22	1:A:97:VAL:HG22	1.74	0.70
1:B:42:His:CD2	1:B:238:VAL:H	2.05	0.69
1:A:22:His:HD2	1:A:260:ALA:H	1.40	0.69
1:A:180:ARG:HG2	1:A:180:ARG:NH2	2.08	0.67
1:B:95:ILE:HG22	1:B:97:VAL:HG22	1.76	0.66
1:A:83:VAL:HA	1:A:86:LEU:CD1	2.25	0.66
1:B:227:ARG:O	1:B:231:MSE:HG3	1.96	0.66
1:B:180:ARG:HH22	1:B:237:ASP:CG	1.99	0.66
1:B:30:ARG:HH11	1:B:30:ARG:HG3	1.61	0.64
1:B:228:LEU:CA	1:B:231:MSE:HE2	2.23	0.64
1:A:129:ASP:OD2	1:B:109:ARG:HD2	1.98	0.63
1:A:178:GLY:O	1:A:179:TYR:HB3	1.96	0.63
1:B:96:VAL:HG21	1:B:220:PHE:HD2	1.63	0.62
1:B:96:VAL:O	1:B:96:VAL:HG23	2.00	0.61
1:A:118:ASP:OD2	1:B:124:ARG:NH2	2.34	0.61
1:A:188:GLU:HA	1:A:191:ARG:HH11	1.66	0.60
1:B:136:TRP:CH2	1:B:138:LYS:HD3	2.36	0.60
1:A:207:ASP:CG	1:A:227:ARG:HH22	2.04	0.60
1:B:66:ILE:O	1:B:70:GLU:HG2	2.01	0.59
1:B:266:VAL:HB	1:B:269:TYR:CD1	2.38	0.59
1:A:22:His:HB3	1:A:260:ALA:HB3	1.85	0.59
1:A:219:PRO:HG2	1:A:220:PHE:H	1.68	0.58
1:B:180:ARG:NH2	1:B:237:ASP:OD1	2.36	0.58
1:A:62:GLN:HG3	1:A:63:LEU:CD1	2.33	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:PHE:CD1	1:B:157:TYR:HA	2.39	0.58
1:A:36:PHE:HB2	1:A:86:LEU:HD23	1.87	0.57
1:A:176:PRO:HB3	1:B:109:ARG:HH11	1.70	0.57
1:A:29:ALA:HA	1:A:263:VAL:HG11	1.87	0.57
1:A:265:LEU:C	1:A:265:LEU:HD23	2.25	0.57
1:A:271:GLN:O	1:A:275:GLU:HG3	2.05	0.56
1:B:25:HIS:CE1	1:B:289:LEU:HD12	2.40	0.56
1:B:102:VAL:HG13	1:B:102:VAL:O	2.05	0.56
1:B:115:LEU:O	1:B:119:ILE:HG12	2.06	0.55
1:A:65:HIS:H	1:A:65:HIS:CD2	2.22	0.55
1:B:136:TRP:CE2	1:B:138:LYS:HG2	2.42	0.55
1:A:73:LEU:HB3	1:A:122:ARG:NE	2.22	0.54
1:A:34:ALA:HA	1:A:82:GLU:OE2	2.07	0.54
1:B:194:LYS:HB2	1:B:194:LYS:NZ	2.22	0.54
1:A:62:GLN:O	1:A:63:LEU:HD12	2.08	0.53
1:A:122:ARG:O	1:A:126:LEU:HD13	2.09	0.53
1:B:102:VAL:HG11	1:B:113:PHE:CE2	2.44	0.53
1:A:255:ARG:NH1	1:A:280:GLU:HG3	2.24	0.52
1:B:21:VAL:HG11	1:B:23:ARG:NH2	2.23	0.52
1:A:188:GLU:HA	1:A:191:ARG:NH1	2.24	0.52
1:A:30:ARG:HH11	1:A:30:ARG:HG3	1.75	0.52
1:A:135:ILE:HB	1:A:204:ILE:HD13	1.91	0.51
1:A:266:VAL:HB	1:A:269:TYR:CD2	2.46	0.51
1:A:180:ARG:HH21	1:A:180:ARG:CG	2.24	0.50
1:A:264:GLU:HA	1:A:264:GLU:OE1	2.12	0.50
1:A:227:ARG:O	1:A:231:MSE:HG3	2.13	0.49
1:A:22:HIS:CD2	1:A:260:ALA:H	2.26	0.49
1:A:92:ARG:NH1	1:A:172:MSE:HE3	2.28	0.49
1:A:66:ILE:O	1:A:70:GLU:HG3	2.13	0.48
1:B:66:ILE:H	1:B:66:ILE:CD1	2.18	0.48
1:A:226:GLU:OE1	1:A:255:ARG:HD2	2.13	0.48
1:A:245:GLY:H	1:A:264:GLU:HG2	1.79	0.48
1:A:62:GLN:HG3	1:A:63:LEU:HD13	1.94	0.48
1:B:120:GLN:HG3	1:B:171:LEU:HD22	1.96	0.48
1:A:86:LEU:H	1:A:86:LEU:HD12	1.79	0.48
1:B:83:VAL:HG11	1:B:93:LEU:HD13	1.96	0.48
1:B:96:VAL:HG12	1:B:170:ILE:CD1	2.44	0.48
1:B:134:ILE:HB	1:B:170:ILE:HB	1.96	0.48
1:B:210:GLY:O	1:B:211:GLU:HB2	2.14	0.48
1:B:66:ILE:HD12	1:B:66:ILE:N	2.21	0.47
1:A:240:LEU:HA	1:A:261:LEU:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:VAL:CG1	1:A:115:LEU:HD13	2.45	0.47
1:A:274:LYS:HD3	1:A:288:VAL:HG23	1.97	0.47
1:A:92:ARG:HD3	1:A:172:MSE:CG	2.44	0.47
1:B:117:ALA:O	1:B:121:VAL:HG23	2.15	0.46
1:A:118:ASP:CG	1:B:124:ARG:HH22	2.18	0.46
1:A:251:ILE:HD11	1:A:276:ARG:HG2	1.98	0.46
1:B:42:HIS:ND1	1:B:180:ARG:NH2	2.64	0.46
1:B:96:VAL:HG12	1:B:170:ILE:HD13	1.97	0.46
1:A:65:HIS:H	1:A:65:HIS:HD2	1.64	0.46
1:B:67:GLU:OE2	1:B:67:GLU:HA	2.16	0.46
1:B:102:VAL:HG13	1:B:111:LEU:HD21	1.98	0.46
1:A:103:ALA:HA	1:A:110:HIS:HA	1.97	0.46
1:B:23:ARG:HH11	1:B:289:LEU:HD11	1.81	0.46
1:A:92:ARG:HH11	1:A:172:MSE:HE3	1.81	0.45
1:A:43:LEU:CD2	1:A:44:VAL:N	2.80	0.45
1:A:157:TYR:OH	1:B:194:LYS:HE3	2.16	0.45
1:B:198:HIS:HB3	2:B:326:HOH:O	2.16	0.45
1:A:111:LEU:HD13	1:B:129:ASP:HB3	1.99	0.45
1:B:194:LYS:HB2	1:B:194:LYS:HZ2	1.81	0.44
1:B:98:GLY:CA	1:B:168:GLU:HG2	2.47	0.44
1:B:281:VAL:HG12	1:B:284:PHE:HB2	1.98	0.44
1:A:201:PHE:CE2	1:B:161:ALA:HB3	2.53	0.44
1:B:240:LEU:HA	1:B:261:LEU:O	2.18	0.44
1:B:271:GLN:NE2	1:B:274:LYS:HD2	2.33	0.44
1:B:179:TYR:CD1	1:B:179:TYR:N	2.79	0.44
1:A:115:LEU:HD22	1:A:119:ILE:HD11	2.00	0.43
1:A:124:ARG:HG2	1:A:124:ARG:HH11	1.83	0.43
1:A:23:ARG:HH11	1:A:25:HIS:CE1	2.37	0.43
1:A:49:PRO:HD3	1:A:243:PHE:CD2	2.53	0.43
1:A:62:GLN:HG3	1:A:63:LEU:HD12	2.00	0.43
1:A:255:ARG:HH12	1:A:280:GLU:HG3	1.83	0.43
1:A:287:GLU:OE2	1:A:289:LEU:HD11	2.18	0.43
1:B:271:GLN:HE22	1:B:274:LYS:HD2	1.83	0.43
1:B:96:VAL:O	1:B:96:VAL:CG2	2.64	0.43
1:A:179:TYR:HD1	1:A:180:ARG:O	2.02	0.43
1:A:245:GLY:N	1:A:264:GLU:HG2	2.34	0.43
1:B:50:TYR:CE2	1:B:100:VAL:HG11	2.54	0.43
1:A:48:PRO:CG	1:A:97:VAL:HG13	2.48	0.42
1:A:197:PHE:CE1	1:B:157:TYR:HA	2.54	0.42
1:A:251:ILE:CD1	1:A:276:ARG:HG2	2.49	0.42
1:B:38:GLU:HG3	1:B:85:ARG:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:38:GLU:CG	1:B:39:ALA:N	2.82	0.42
1:A:62:GLN:C	1:A:63:LEU:HD12	2.39	0.42
1:B:179:TYR:H	1:B:179:TYR:HD1	1.63	0.42
1:B:231:MSE:HE3	1:B:232:PHE:CE2	2.55	0.42
1:B:208:ILE:HD11	1:B:227:ARG:CZ	2.50	0.41
1:A:97:VAL:HG11	1:A:115:LEU:HD13	2.01	0.41
1:A:65:HIS:CD2	1:A:65:HIS:N	2.89	0.41
1:B:180:ARG:NH1	1:B:180:ARG:HG2	2.36	0.41
1:B:138:LYS:HE2	1:B:168:GLU:OE2	2.20	0.41
1:A:182:PRO:HA	1:A:186:GLN:OE1	2.21	0.41
1:B:138:LYS:HD2	1:B:210:GLY:HA3	2.02	0.41
1:A:28:ASP:OD2	1:A:31:GLU:HG2	2.21	0.41
1:A:53:LEU:HD23	1:A:53:LEU:HA	1.92	0.41
1:A:118:ASP:CG	1:B:124:ARG:NH2	2.74	0.41
1:B:164:LYS:HE2	1:B:164:LYS:HB3	1.92	0.40
1:B:169:TYR:CD2	1:B:169:TYR:N	2.90	0.40
1:B:266:VAL:HA	1:B:267:PRO:HD2	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	223/297 (75%)	214 (96%)	8 (4%)	1 (0%)	30 49
1	B	222/297 (75%)	213 (96%)	8 (4%)	1 (0%)	25 44
All	All	445/594 (75%)	427 (96%)	16 (4%)	2 (0%)	30 49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	179	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	179	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/248 (81%)	186 (92%)	15 (8%)	11	23
1	B	199/248 (80%)	184 (92%)	15 (8%)	11	23
All	All	400/496 (81%)	370 (92%)	30 (8%)	11	23

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LEU
1	A	53	LEU
1	A	65	HIS
1	A	66	ILE
1	A	86	LEU
1	A	96	VAL
1	A	97	VAL
1	A	115	LEU
1	A	172	MSE
1	A	222	LEU
1	A	224	LEU
1	A	248	THR
1	A	264	GLU
1	A	280	GLU
1	A	286	LEU
1	B	23	ARG
1	B	24	LEU
1	B	30	ARG
1	B	43	LEU
1	B	52	THR
1	B	65	HIS
1	B	97	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	109	ARG
1	B	111	LEU
1	B	115	LEU
1	B	131	LEU
1	B	138	LYS
1	B	222	LEU
1	B	248	THR
1	B	271	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	42	HIS
1	A	62	GLN
1	A	65	HIS
1	A	173	GLN
1	B	22	HIS
1	B	25	HIS
1	B	42	HIS
1	B	139	HIS
1	B	173	GLN
1	B	271	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/297 (77%)	0.14	14 (6%) 28 26	16, 33, 64, 82	0
1	B	230/297 (77%)	-0.05	6 (2%) 57 54	16, 34, 59, 82	0
All	All	461/594 (77%)	0.04	20 (4%) 40 37	16, 34, 63, 82	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	TYR	4.4
1	A	53	LEU	3.7
1	B	156	PRO	3.3
1	A	219	PRO	3.1
1	A	180	ARG	3.0
1	A	52	THR	3.0
1	B	108	GLY	3.0
1	A	288	VAL	2.9
1	A	103	ALA	2.9
1	B	290	ASP	2.8
1	A	102	VAL	2.8
1	B	50	TYR	2.7
1	B	102	VAL	2.5
1	B	179	TYR	2.4
1	A	50	TYR	2.4
1	A	289	LEU	2.3
1	A	104	ARG	2.3
1	A	285	SER	2.2
1	A	178	GLY	2.1
1	A	156	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.