



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2024 – 05:53 PM EDT

PDB ID : 2ZXK  
Title : Crystal structure of SeMet-Red chlorophyll catabolite reductase  
Authors : Sugishima, M.; Kitamori, Y.; Fukuyama, K.  
Deposited on : 2008-12-29  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	 64% 25% 5% 7%
1	B	285	 55% 29% 11%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Red chlorophyll catabolite reductase, chloroplastic.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	266	2098	1327	361	401	3	6	0	0	0
1	B	253	1967	1247	334	378	3	5	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	GLY	-	expression tag	UNP Q8LDU4
A	36	PRO	-	expression tag	UNP Q8LDU4
A	37	LEU	-	expression tag	UNP Q8LDU4
A	38	GLY	-	expression tag	UNP Q8LDU4
A	39	SER	-	expression tag	UNP Q8LDU4
B	35	GLY	-	expression tag	UNP Q8LDU4
B	36	PRO	-	expression tag	UNP Q8LDU4
B	37	LEU	-	expression tag	UNP Q8LDU4
B	38	GLY	-	expression tag	UNP Q8LDU4
B	39	SER	-	expression tag	UNP Q8LDU4

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total	O	0	0
			33	33		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	11	Total	O	0	0
			11	11		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Red chlorophyll catabolite reductase, chloroplastic



- Molecule 1: Red chlorophyll catabolite reductase, chloroplastic



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.98Å 83.54Å 120.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.48 – 2.50 19.48 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.48-2.50) 95.4 (19.48-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	20.98 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.215 , 0.291 0.228 , 0.303	Depositor DCC
$R_{free}$ test set	963 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.5	Xtrriage
Anisotropy	0.085	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 26.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4111	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	1/2134 (0.0%)	0.87	0/2878
1	B	0.68	0/2001	0.81	1/2706 (0.0%)
All	All	0.77	1/4135 (0.0%)	0.84	1/5584 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	CYS	CB-SG	5.62	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	LEU	CA-CB-CG	5.19	127.23	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	229	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2098	0	2070	52	0
1	B	1967	0	1908	56	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	33	0	0	3	0
3	B	11	0	0	0	0
All	All	4111	0	3978	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:LYS:HA	1:B:187:GLN:HG3	1.52	0.92
1:B:197:SER:HA	1:B:200:LYS:HD2	1.64	0.80
1:A:162:SER:HB2	1:A:228:ASP:HB3	1.63	0.79
1:B:183:LYS:HA	1:B:187:GLN:CG	2.15	0.76
1:A:162:SER:HA	1:A:227:ILE:O	1.86	0.75
1:A:273:ARG:HG2	1:A:273:ARG:HH11	1.53	0.73
1:A:66:MSE:HE3	1:A:155:LEU:HD12	1.72	0.71
1:A:161:LYS:O	1:A:228:ASP:HA	1.94	0.68
1:B:124:HIS:HA	1:B:133:LEU:O	1.94	0.68
1:B:62:ARG:HH21	1:B:136:THR:HG23	1.57	0.68
1:A:55:PHE:HB2	1:A:63:LYS:HG2	1.76	0.68
1:A:76:ARG:HG2	1:A:76:ARG:NH2	2.08	0.67
1:A:70:MSE:HE2	1:A:138:ILE:HG12	1.76	0.67
1:A:189:THR:CG2	1:A:191:LEU:HG	2.27	0.65
1:B:62:ARG:NH2	1:B:136:THR:HG23	2.11	0.65
1:A:168:ASP:CG	1:A:169:LEU:H	2.01	0.65
1:A:189:THR:HG21	1:A:191:LEU:HG	1.78	0.64
1:A:280:ASP:O	1:A:284:ARG:HG3	1.98	0.63
1:A:50:ARG:HA	1:A:50:ARG:NE	2.14	0.62
1:A:76:ARG:HH21	1:A:76:ARG:CG	2.13	0.61
1:A:149:PRO:HD2	1:A:258:CYS:SG	2.42	0.60

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:GLN:NE2	1:A:114:PRO:HD2	2.17	0.59
1:B:66:MSE:HE1	1:B:138:ILE:HG13	1.84	0.59
1:A:244:SER:HB2	1:A:245:PRO:HD3	1.84	0.59
1:B:80:GLN:O	1:B:109:GLY:HA2	2.03	0.58
1:B:82:LEU:HD11	1:B:110:ASP:HB3	1.84	0.58
1:B:119:ILE:CG2	1:B:139:SER:HB2	2.33	0.58
1:B:244:SER:HB2	1:B:245:PRO:HD3	1.86	0.58
1:B:122:TRP:HA	1:B:135:ILE:O	2.05	0.56
1:B:171:HIS:H	1:B:171:HIS:CD2	2.23	0.56
1:A:76:ARG:NH2	1:A:76:ARG:CG	2.69	0.56
1:A:76:ARG:HG2	1:A:76:ARG:HH21	1.68	0.56
1:B:166:ILE:HD12	1:B:218:PHE:CZ	2.40	0.56
1:B:245:PRO:O	1:B:249:GLU:HG3	2.05	0.56
1:B:162:SER:HA	1:B:227:ILE:O	2.06	0.56
1:A:273:ARG:HH11	1:A:273:ARG:CG	2.18	0.56
1:B:70:MSE:HE2	1:B:74:GLU:HG3	1.89	0.55
1:B:69:LEU:O	1:B:73:VAL:HG23	2.06	0.55
1:A:288:ILE:HG23	1:A:292:LEU:HD12	1.89	0.55
1:B:240:ARG:NH1	1:B:241:ASP:OD1	2.40	0.54
1:B:62:ARG:HH21	1:B:136:THR:CG2	2.20	0.54
1:B:62:ARG:HH11	1:B:157:GLN:HE21	1.54	0.54
1:B:150:ASN:O	1:B:169:LEU:HA	2.07	0.54
1:A:219:SER:HB2	1:A:220:PRO:HD2	1.89	0.54
1:A:168:ASP:CG	1:A:169:LEU:N	2.61	0.53
1:A:92:ASN:HB3	3:A:322:HOH:O	2.07	0.53
1:A:171:HIS:H	1:A:171:HIS:CD2	2.26	0.53
1:A:143:ASN:C	1:A:143:ASN:HD22	2.11	0.53
1:A:180:ASP:O	1:A:184:GLU:HG2	2.07	0.53
1:A:143:ASN:HD22	1:A:145:SER:H	1.57	0.52
1:B:166:ILE:HD12	1:B:218:PHE:HZ	1.75	0.52
1:A:169:LEU:HD11	1:A:254:TRP:HB2	1.92	0.51
1:B:124:HIS:ND1	1:B:134:ASN:ND2	2.59	0.51
1:A:159:SER:HB3	1:A:162:SER:HB3	1.92	0.50
1:B:80:GLN:NE2	1:B:114:PRO:HD2	2.26	0.50
1:A:187:GLN:C	1:A:189:THR:H	2.13	0.50
1:B:305:SER:O	1:B:309:VAL:HG12	2.12	0.50
1:B:57:TYR:OH	1:B:101:GLU:OE1	2.14	0.50
1:B:168:ASP:OD2	1:B:218:PHE:HD2	1.95	0.49
1:B:291:ASP:HB3	1:B:292:LEU:HD23	1.95	0.49
1:A:70:MSE:CE	1:A:138:ILE:HG12	2.40	0.48
1:A:161:LYS:O	1:A:161:LYS:HD2	2.13	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:MSE:HE1	1:B:118:VAL:HG11	1.94	0.48
1:B:77:LEU:O	1:B:79:SER:N	2.46	0.48
1:B:139:SER:HB3	1:B:141:PHE:CE2	2.48	0.48
1:A:203:GLU:OE2	1:A:234:LYS:HE2	2.13	0.47
1:A:155:LEU:CD2	1:A:163:LEU:HD21	2.45	0.47
1:B:119:ILE:HG23	1:B:139:SER:HB2	1.96	0.47
1:B:114:PRO:HG2	1:B:115:ILE:HD12	1.97	0.47
1:A:142:LEU:HD21	1:A:254:TRP:CZ2	2.50	0.46
1:A:202:PRO:HD2	1:A:203:GLU:OE1	2.15	0.46
1:B:255:LEU:C	1:B:257:ARG:H	2.19	0.46
1:A:192:ASP:O	1:A:196:GLN:HG3	2.14	0.46
1:A:161:LYS:O	1:A:228:ASP:CA	2.64	0.45
1:B:82:LEU:CD1	1:B:110:ASP:HB3	2.46	0.45
1:A:144:SER:C	1:A:146:THR:H	2.20	0.45
1:B:144:SER:O	1:B:146:THR:N	2.50	0.45
1:B:192:ASP:HB2	1:B:221:THR:HG22	1.99	0.45
1:A:253:VAL:O	1:A:257:ARG:HG2	2.16	0.45
1:A:293:ASP:O	1:A:297:PRO:HG2	2.16	0.45
1:A:84:CYS:HB2	1:A:107:ARG:HD2	1.98	0.45
1:B:66:MSE:HE2	1:B:136:THR:HB	2.00	0.44
1:B:95:ASN:HA	1:B:96:PRO:HD2	1.75	0.44
1:A:51:LYS:O	1:A:51:LYS:HD2	2.17	0.44
1:B:288:ILE:HD12	1:B:288:ILE:HA	1.75	0.44
1:A:189:THR:HG22	1:A:191:LEU:HG	2.00	0.44
1:B:66:MSE:CE	1:B:138:ILE:HG13	2.48	0.44
1:B:307:ARG:O	1:B:310:HIS:HB3	2.18	0.43
1:A:51:LYS:HE3	1:A:56:PRO:HG3	2.01	0.43
1:A:298:ARG:HD3	1:A:299:MSE:H	1.83	0.43
1:B:213:PHE:O	1:B:216:SER:HB2	2.18	0.43
1:A:76:ARG:NH1	1:A:252:GLU:OE2	2.52	0.43
1:B:194:HIS:O	1:B:198:LEU:HD22	2.19	0.42
3:A:324:HOH:O	1:B:208:VAL:CG2	2.67	0.42
1:B:81:LEU:HG	1:B:107:ARG:HB3	2.01	0.42
1:A:69:LEU:HB3	1:A:138:ILE:HD13	2.01	0.42
1:A:187:GLN:C	1:A:189:THR:N	2.73	0.42
1:B:144:SER:C	1:B:146:THR:H	2.22	0.42
1:B:209:SER:CB	1:B:214:VAL:CG2	2.97	0.42
1:B:187:GLN:HE21	1:B:187:GLN:HB3	1.69	0.42
1:A:72:THR:O	1:A:76:ARG:HB2	2.19	0.41
1:B:308:VAL:O	1:B:312:ILE:HG12	2.20	0.41
3:A:324:HOH:O	1:B:208:VAL:HG21	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:LYS:HA	1:B:187:GLN:HG2	2.01	0.41
1:B:113:SER:HB2	1:B:114:PRO:HD2	2.02	0.41
1:B:248:LYS:O	1:B:252:GLU:HB2	2.19	0.40
1:A:95:ASN:OD1	1:A:97:ASN:CG	2.60	0.40
1:B:185:TYR:HE2	1:B:257:ARG:HD2	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/285 (92%)	241 (92%)	18 (7%)	3 (1%)	12	23
1	B	247/285 (87%)	223 (90%)	16 (6%)	8 (3%)	3	4
All	All	509/570 (89%)	464 (91%)	34 (7%)	11 (2%)	5	9

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	78	GLN
1	B	111	LYS
1	B	290	ASP
1	B	299	MSE
1	A	130	GLY
1	A	269	GLY
1	B	145	SER
1	B	159	SER
1	B	96	PRO
1	B	99	SER
1	A	259	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/257 (92%)	208 (88%)	29 (12%)	4	8
1	B	220/257 (86%)	195 (89%)	25 (11%)	4	9
All	All	457/514 (89%)	403 (88%)	54 (12%)	4	9

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	53	MSE
1	A	63	LYS
1	A	69	LEU
1	A	71	SER
1	A	76	ARG
1	A	81	LEU
1	A	82	LEU
1	A	94	ASN
1	A	143	ASN
1	A	161	LYS
1	A	162	SER
1	A	169	LEU
1	A	172	ARG
1	A	175	LEU
1	A	177	LEU
1	A	198	LEU
1	A	226	LYS
1	A	232	GLU
1	A	253	VAL
1	A	260	LYS
1	A	273	ARG
1	A	281	LYS
1	A	286	LYS
1	A	295	GLN
1	A	297	PRO
1	A	298	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	302	GLU
1	A	304	VAL
1	B	51	LYS
1	B	65	LEU
1	B	69	LEU
1	B	78	GLN
1	B	81	LEU
1	B	82	LEU
1	B	94	ASN
1	B	95	ASN
1	B	108	SER
1	B	156	ILE
1	B	166	ILE
1	B	167	LEU
1	B	169	LEU
1	B	177	LEU
1	B	187	GLN
1	B	198	LEU
1	B	214	VAL
1	B	228	ASP
1	B	232	GLU
1	B	235	LEU
1	B	237	GLU
1	B	239	LEU
1	B	257	ARG
1	B	277	GLU
1	B	288	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	A	92	ASN
1	A	143	ASN
1	A	171	HIS
1	A	194	HIS
1	A	310	HIS
1	B	75	ASN
1	B	80	GLN
1	B	94	ASN
1	B	95	ASN
1	B	134	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	157	GLN
1	B	171	HIS
1	B	187	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	260/285 (91%)	-0.28	0 <b>100</b>   <b>100</b>	12, 24, 37, 40	0
1	B	247/285 (86%)	-0.23	1 (0%) <b>89</b>   <b>86</b>	17, 31, 47, 52	0
All	All	507/570 (88%)	-0.26	1 (0%) <b>92</b>   <b>90</b>	12, 28, 43, 52	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	160	SER	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	A	1	1/1	0.81	0.14	39,39,39,39	0
2	NA	B	2	1/1	0.87	0.19	40,40,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.