



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 10, 2024 – 06:44 AM EST

PDB ID : 3AU3  
Title : Crystal structure of armadillo repeat domain of APC  
Authors : Murayama, K.; Kato-Murayama, M.; Terada, T.; Shirouzu, M.; Yokoyama, S.;  
RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2011-01-28  
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenomatous polyposis coli protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	301	2327	1443	417	444	12	11	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	GLY	-	expression tag	UNP P25054
A	390	SER	-	expression tag	UNP P25054
A	391	SER	-	expression tag	UNP P25054
A	392	GLY	-	expression tag	UNP P25054
A	393	SER	-	expression tag	UNP P25054
A	394	SER	-	expression tag	UNP P25054
A	395	GLY	-	expression tag	UNP P25054


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	117	Total	O	0	0
			117	117		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenomatous polyposis coli protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.33Å 114.33Å 61.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.51 – 2.10 49.51 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.51-2.10) 99.9 (49.51-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.88 (at 2.10Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.204 , 0.232 0.209 , 0.237	Depositor DCC
$R_{free}$ test set	1316 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.7	Xtrriage
Anisotropy	0.130	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 37.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2444	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2347	0.48	0/3153

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2327	0	2340	56	0
2	A	117	0	0	4	0
All	All	2444	0	2340	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:518:THR:HG22	1:A:522:MSE:HE2	1.30	1.13
1:A:519:LEU:HA	1:A:522:MSE:HE3	1.45	0.95
1:A:435:LYS:HG3	1:A:436:ASN:OD1	1.70	0.91
1:A:432:ASP:HB2	1:A:435:LYS:HE2	1.65	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:703:ALA:HA	1:A:706:MSE:HE2	1.66	0.76
1:A:628:THR:HG23	2:A:116:HOH:O	1.89	0.72
1:A:423:TRP:O	1:A:427:HIS:HB3	1.92	0.69
1:A:518:THR:HG22	1:A:522:MSE:CE	2.17	0.68
1:A:649:ASN:HD22	1:A:649:ASN:C	1.98	0.67
1:A:473:GLN:O	1:A:477:GLU:HG3	1.98	0.64
1:A:414:ARG:NH2	1:A:469:LEU:O	2.32	0.62
1:A:564:ARG:HG3	1:A:602:ASN:HD21	1.63	0.62
1:A:582:GLU:OE2	1:A:626:THR:HG22	2.00	0.62
1:A:538:GLU:HG2	1:A:541:GLN:NE2	2.15	0.60
1:A:654:GLN:O	1:A:658:GLU:HG3	2.01	0.60
1:A:570:LYS:NZ	1:A:608:ALA:HB3	2.16	0.59
1:A:532:GLN:HB2	1:A:544:ILE:HG12	1.84	0.59
1:A:626:THR:HG23	1:A:628:THR:HG22	1.85	0.59
1:A:561:LYS:O	1:A:565:GLU:HG3	2.02	0.58
1:A:432:ASP:O	1:A:435:LYS:HG2	2.07	0.54
1:A:407:LEU:O	1:A:411:GLU:HG3	2.09	0.53
1:A:495:ILE:O	1:A:499:ARG:HG3	2.09	0.53
1:A:542:GLN:HE22	1:A:583:SER:HB2	1.72	0.52
1:A:472:LEU:HD23	1:A:522:MSE:HE1	1.92	0.52
1:A:649:ASN:HD21	1:A:651:ASP:HB2	1.74	0.51
1:A:555:ALA:O	1:A:560:LYS:HE3	2.11	0.51
1:A:427:HIS:CD2	1:A:436:ASN:HD22	2.30	0.50
1:A:654:GLN:HE21	1:A:658:GLU:CG	2.25	0.49
1:A:564:ARG:HG3	1:A:602:ASN:ND2	2.27	0.49
1:A:464:HIS:O	1:A:468:GLU:HG2	2.13	0.49
1:A:431:MSE:HE3	1:A:431:MSE:HA	1.95	0.48
1:A:626:THR:CG2	1:A:628:THR:HG22	2.44	0.47
1:A:432:ASP:HB3	1:A:435:LYS:HG2	1.97	0.47
1:A:704:VAL:HG23	1:A:705:SER:N	2.29	0.47
1:A:575:CYS:O	1:A:579:VAL:HG13	2.14	0.47
1:A:445:GLN:HA	2:A:7:HOH:O	2.14	0.47
1:A:467:ASN:HD21	1:A:515:ASN:HD21	1.62	0.47
1:A:570:LYS:HZ1	1:A:608:ALA:HB3	1.79	0.46
1:A:427:HIS:CD2	1:A:436:ASN:ND2	2.84	0.46
1:A:624:SER:OG	1:A:628:THR:HG22	2.17	0.45
1:A:397:ASP:O	1:A:401:ARG:HG2	2.17	0.45
1:A:688:SER:HB3	2:A:99:HOH:O	2.17	0.43
1:A:432:ASP:CB	1:A:435:LYS:HG2	2.48	0.43
1:A:570:LYS:HZ3	1:A:608:ALA:HB3	1.83	0.43
1:A:427:HIS:HD2	1:A:436:ASN:HB2	1.83	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:602:ASN:ND2	2:A:39:HOH:O	2.52	0.43
1:A:472:LEU:CD2	1:A:522:MSE:HE1	2.49	0.43
1:A:451:CYS:SG	1:A:455:LYS:HE3	2.59	0.42
1:A:519:LEU:CA	1:A:522:MSE:HE3	2.32	0.42
1:A:629:LEU:O	1:A:633:GLU:HG3	2.19	0.42
1:A:649:ASN:C	1:A:649:ASN:ND2	2.70	0.42
1:A:441:PRO:HB2	1:A:493:TYR:HE2	1.84	0.42
1:A:472:LEU:HD13	1:A:515:ASN:HB3	2.01	0.42
1:A:400:GLY:O	1:A:404:ILE:HG13	2.21	0.41
1:A:555:ALA:HB1	1:A:559:SER:HB2	2.03	0.41
1:A:423:TRP:CZ2	1:A:438:MSE:HB2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	297/344 (86%)	291 (98%)	5 (2%)	1 (0%)	37 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	429	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	257/276 (93%)	255 (99%)	2 (1%)	79 84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	436	ASN
1	A	649	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	467	ASN
1	A	541	GLN
1	A	542	GLN
1	A	602	ASN
1	A	649	ASN
1	A	654	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/344 (84%)	0.57	27 (9%) <b>16</b> <b>17</b>	23, 36, 78, 95	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	707	LEU	8.2
1	A	689	ALA	6.5
1	A	704	VAL	6.1
1	A	700	ASP	6.1
1	A	703	ALA	5.4
1	A	702	GLY	5.2
1	A	430	GLY	4.7
1	A	429	PRO	4.5
1	A	626	THR	4.0
1	A	435	LYS	4.0
1	A	625	GLN	3.8
1	A	436	ASN	3.4
1	A	627	ASN	3.0
1	A	427	HIS	3.0
1	A	433	GLN	2.9
1	A	397	ASP	2.8
1	A	628	THR	2.7
1	A	705	SER	2.6
1	A	688	SER	2.5
1	A	426	ALA	2.5
1	A	399	ARG	2.4
1	A	434	ASP	2.4
1	A	622	TYR	2.3
1	A	660	ASN	2.2
1	A	672	HIS	2.2
1	A	608	ALA	2.1
1	A	432	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.