



Full wwPDB X-ray Structure Validation Report i

Oct 23, 2024 – 02:32 AM EDT

PDB ID : 3AVF
Title : Crystal structures of novel allosteric peptide inhibitors of HIV integrase in the LEDGF binding site
Authors : Peat, T.S.; Deadman, J.J.; Newman, J.; Rhodes, D.I.
Deposited on : 2011-03-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACY	A	210	-	-	X	-
5	ACY	A	211	-	-	X	-
5	ACY	B	210	-	-	X	-

2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 2922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	1281	823	215	238	5	0	18	0
1	B	150	1280	822	217	236	5	0	17	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	expression tag	UNP P12497
A	31	GLY	-	expression tag	UNP P12497
A	32	SER	-	expression tag	UNP P12497
A	33	SER	-	expression tag	UNP P12497
A	34	HIS	-	expression tag	UNP P12497
A	35	HIS	-	expression tag	UNP P12497
A	36	HIS	-	expression tag	UNP P12497
A	37	HIS	-	expression tag	UNP P12497
A	38	HIS	-	expression tag	UNP P12497
A	39	HIS	-	expression tag	UNP P12497
A	40	SER	-	expression tag	UNP P12497
A	41	SER	-	expression tag	UNP P12497
A	42	GLY	-	expression tag	UNP P12497
A	43	LEU	-	expression tag	UNP P12497
A	44	VAL	-	expression tag	UNP P12497
A	45	PRO	-	expression tag	UNP P12497
A	46	ARG	-	expression tag	UNP P12497
A	47	GLY	-	expression tag	UNP P12497
A	48	SER	-	expression tag	UNP P12497
A	49	HIS	-	expression tag	UNP P12497
A	56	SER	CYS	engineered mutation	UNP P12497
A	139	ASP	PHE	engineered mutation	UNP P12497
A	185	HIS	PHE	engineered mutation	UNP P12497
B	30	MET	-	expression tag	UNP P12497
B	31	GLY	-	expression tag	UNP P12497

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	SER	-	expression tag	UNP P12497
B	33	SER	-	expression tag	UNP P12497
B	34	HIS	-	expression tag	UNP P12497
B	35	HIS	-	expression tag	UNP P12497
B	36	HIS	-	expression tag	UNP P12497
B	37	HIS	-	expression tag	UNP P12497
B	38	HIS	-	expression tag	UNP P12497
B	39	HIS	-	expression tag	UNP P12497
B	40	SER	-	expression tag	UNP P12497
B	41	SER	-	expression tag	UNP P12497
B	42	GLY	-	expression tag	UNP P12497
B	43	LEU	-	expression tag	UNP P12497
B	44	VAL	-	expression tag	UNP P12497
B	45	PRO	-	expression tag	UNP P12497
B	46	ARG	-	expression tag	UNP P12497
B	47	GLY	-	expression tag	UNP P12497
B	48	SER	-	expression tag	UNP P12497
B	49	HIS	-	expression tag	UNP P12497
B	56	SER	CYS	engineered mutation	UNP P12497
B	139	ASP	PHE	engineered mutation	UNP P12497
B	185	HIS	PHE	engineered mutation	UNP P12497

- Molecule 2 is a protein called LEDGF peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	8	Total	C	N	O	0	1	0
			73	46	11	16			
2	F	8	Total	C	N	O	0	0	0
			65	40	10	15			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	84	Total O 84 84	0	0
6	B	82	Total O 82 82	0	0
6	D	1	Total O 1 1	0	0
6	F	2	Total O 2 2	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrase




- Molecule 1: Integrase



- Molecule 2: LEDGF peptide



- Molecule 2: LEDGF peptide



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	70.65Å 70.65Å 66.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.19 – 1.70 61.19 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.19-1.70) 99.9 (61.19-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.70 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.6.0062	Depositor
R , R_{free}	0.161 , 0.199 0.159 , 0.197	Depositor DCC
R_{free} test set	2064 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 35.7	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.020 for -h,-k,l 0.477 for h,-h-k,-l 0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2922	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	116	ASP	CB-CG-OD1	7.23	124.81	118.30
1	B	102	LEU	CB-CG-CD2	-6.58	99.81	111.00
1	A	57	SER	CB-CA-C	5.96	121.42	110.10
1	B	107	ARG	NE-CZ-NH1	-5.77	117.41	120.30
1	A	102	LEU	CB-CG-CD2	-5.77	101.19	111.00
1	A	107	ARG	NE-CZ-NH1	-5.70	117.45	120.30
2	F	4	ILE	CG1-CB-CG2	-5.29	99.75	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1281	0	1310	51	0
1	B	1280	0	1309	42	0
2	D	73	0	74	5	0
2	F	65	0	64	0	0
3	A	20	0	0	2	0
3	B	20	0	0	1	1
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	8	0	6	12	0
5	B	4	0	3	5	0
6	A	84	0	0	11	1
6	B	82	0	0	10	2
6	D	1	0	0	0	0
6	F	2	0	0	0	0
All	All	2922	0	2766	98	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171:HIS:HB2	6:B:245:HOH:O	2.15	0.46
1:A:141[B]:ILE:CD1	1:A:142:PRO:HD2	2.28	0.46
1:A:139:ASP:OD1	1:A:141[B]:ILE:CG2	2.54	0.45
5:A:210:ACY:H2	5:A:211:ACY:H2	1.97	0.45
1:A:201:VAL:HG12	1:B:205:ALA:HB2	1.99	0.45
1:A:162:ILE:O	1:A:166:ARG:HB2	2.18	0.44
1:A:136[A]:LYS:NZ	6:A:283:HOH:O	1.94	0.44
1:B:141[B]:ILE:HG23	1:B:141[B]:ILE:O	2.17	0.44
1:A:139:ASP:CG	1:A:141[B]:ILE:HG22	2.35	0.44
5:A:211:ACY:C	1:B:109:PRO:HD3	2.48	0.43
1:A:205:ALA:HB2	1:B:201:VAL:HG12	2.00	0.43
1:B:185:HIS:ND1	5:B:210:ACY:H2	2.35	0.42
1:B:141[B]:ILE:HD12	1:B:141[B]:ILE:HA	1.77	0.42
1:B:152:GLU:HG2	1:B:156[B]:LYS:HE3	2.00	0.41
1:A:127:LYS:HZ2	1:A:137:GLN:HE22	1.68	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1:SO4:O3	6:B:280:HOH:O[2_565]	1.97	0.23
6:A:14:HOH:O	6:A:283:HOH:O[3_564]	2.13	0.07
6:B:234:HOH:O	6:B:280:HOH:O[2_565]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	163/180 (91%)	162 (99%)	1 (1%)	0	100 100
1	B	163/180 (91%)	161 (99%)	2 (1%)	0	100 100
2	D	7/8 (88%)	6 (86%)	1 (14%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	339/376 (90%)	334 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/147 (96%)	137 (97%)	4 (3%)	38	21
1	B	140/147 (95%)	138 (99%)	2 (1%)	62	49
2	D	9/8 (112%)	8 (89%)	1 (11%)	5	1
2	F	8/8 (100%)	8 (100%)	0	100	100
All	All	298/310 (96%)	291 (98%)	7 (2%)	45	29

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	SER
1	A	151[A]	ILE
1	A	151[B]	ILE
1	A	167	ASP
1	B	152	GLU
1	B	188	LYS
2	D	2	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
1	A	155	ASN
1	B	155	ASN
1	B	185	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	209	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	7	-	4,4,4	1.45	1 (25%)	6,6,6	1.23	1 (16%)
3	SO4	B	3	-	4,4,4	0.49	0	6,6,6	0.84	0
3	SO4	B	8	-	4,4,4	1.20	0	6,6,6	1.70	1 (16%)
5	ACY	B	210	-	3,3,3	0.67	0	3,3,3	2.57	2 (66%)
3	SO4	A	2	-	4,4,4	0.40	0	6,6,6	1.19	1 (16%)
3	SO4	B	5	-	4,4,4	0.27	0	6,6,6	2.25	3 (50%)
3	SO4	A	6	-	4,4,4	0.35	0	6,6,6	2.28	2 (33%)
3	SO4	B	1	-	4,4,4	0.35	0	6,6,6	1.11	0
5	ACY	A	210	-	3,3,3	1.06	0	3,3,3	1.83	1 (33%)
5	ACY	A	211	-	3,3,3	0.82	0	3,3,3	1.13	0
3	SO4	A	4	-	4,4,4	0.39	0	6,6,6	0.87	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	7	SO4	O1-S	2.39	1.59	1.44

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	6	SO4	O3-S-O1	-4.15	87.87	109.56
3	B	5	SO4	O2-S-O1	-3.59	83.80	109.06
5	B	210	ACY	O-C-CH3	-3.29	109.05	122.53
5	B	210	ACY	OXT-C-O	2.93	132.89	122.03
3	B	5	SO4	O4-S-O2	2.81	124.25	109.56
5	A	210	ACY	OXT-C-O	2.60	131.66	122.03
3	A	7	SO4	O3-S-O1	2.39	122.06	109.56
3	B	5	SO4	O3-S-O2	2.30	121.59	109.56
3	A	6	SO4	O3-S-O2	2.27	121.44	109.56
3	A	2	SO4	O4-S-O2	-2.20	98.04	109.56
3	B	8	SO4	O4-S-O1	-2.14	98.37	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	210	ACY	5	0
3	A	2	SO4	1	0
3	B	5	SO4	1	0
3	A	6	SO4	1	0
3	B	1	SO4	0	1
5	A	210	ACY	11	0
5	A	211	ACY	10	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	149/180 (82%)	-1.14	0 100 100	6, 15, 34, 45	36 (24%)
1	B	150/180 (83%)	-1.13	0 100 100	7, 15, 34, 59	34 (22%)
2	D	8/8 (100%)	-0.44	0 100 100	15, 31, 54, 57	1 (12%)
2	F	8/8 (100%)	-0.54	0 100 100	27, 34, 50, 58	0
All	All	315/376 (83%)	-1.10	0 100 100	6, 16, 38, 59	71 (22%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	7	5/5	0.96	0.08	27,28,34,42	5
3	SO4	B	8	5/5	0.97	0.07	23,29,41,56	5
4	CL	A	1	1/1	0.97	0.07	48,48,48,48	0
4	CL	B	2	1/1	0.97	0.07	48,48,48,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACY	A	210	4/4	0.98	0.08	31,33,40,40	4
5	ACY	A	211	4/4	0.98	0.06	31,33,33,34	4
5	ACY	B	210	4/4	0.98	0.06	40,42,44,49	0
3	SO4	A	4	5/5	0.99	0.05	25,32,35,36	5
3	SO4	A	6	5/5	0.99	0.04	23,28,33,36	5
3	SO4	A	2	5/5	0.99	0.03	27,33,39,41	5
3	SO4	B	1	5/5	0.99	0.04	27,33,43,45	5
3	SO4	B	3	5/5	0.99	0.04	26,35,36,36	5
3	SO4	B	5	5/5	0.99	0.04	21,26,33,35	5

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.