



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 07:51 AM EDT

PDB ID : 3B7D  
Title : Crystal structure of the GLUR2 ligand binding core (HS1S2J) in complex with CNQX at 2.5 Å resolution  
Authors : Hays, F.A.; Center for Structures of Membrane Proteins (CSMP)  
Deposited on : 2007-10-30  
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	 4% 90% 7% ..
1	B	261	 % 92% 7% .
1	C	261	 2% 90% 8% .
1	D	261	 % 93% 7% .
1	E	261	 4% 92% 7% .

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	261	 5% 94% 5%
1	G	261	 5% 91% 7%
1	H	261	 3% 93% 6%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2006	1275	334	383	14	0	0	0
1	B	258	2023	1287	337	385	14	0	2	0
1	C	257	1990	1269	328	379	14	0	0	0
1	D	259	2011	1280	333	384	14	0	0	0
1	E	258	1981	1262	329	377	13	0	0	0
1	F	258	2006	1277	333	383	13	0	0	0
1	G	257	1985	1267	329	375	14	0	0	0
1	H	258	2002	1278	330	380	14	0	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P19491
A	2	ALA	-	expression tag	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
B	1	GLY	-	expression tag	UNP P19491
B	2	ALA	-	expression tag	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
C	1	GLY	-	expression tag	UNP P19491
C	2	ALA	-	expression tag	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491
D	1	GLY	-	expression tag	UNP P19491

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	2	ALA	-	expression tag	UNP P19491
D	118	GLY	-	linker	UNP P19491
D	119	THR	-	linker	UNP P19491
E	1	GLY	-	expression tag	UNP P19491
E	2	ALA	-	expression tag	UNP P19491
E	118	GLY	-	linker	UNP P19491
E	119	THR	-	linker	UNP P19491
F	1	GLY	-	expression tag	UNP P19491
F	2	ALA	-	expression tag	UNP P19491
F	118	GLY	-	linker	UNP P19491
F	119	THR	-	linker	UNP P19491
G	1	GLY	-	expression tag	UNP P19491
G	2	ALA	-	expression tag	UNP P19491
G	118	GLY	-	linker	UNP P19491
G	119	THR	-	linker	UNP P19491
H	1	GLY	-	expression tag	UNP P19491
H	2	ALA	-	expression tag	UNP P19491
H	118	GLY	-	linker	UNP P19491
H	119	THR	-	linker	UNP P19491

- Molecule 2 is 7-nitro-2,3-dioxo-2,3-dihydroquinoxaline-6-carbonitrile (three-letter code: CNI) (formula: C<sub>9</sub>H<sub>2</sub>N<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	17	9	4	4	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			17	9	4	4		
2	C	1	Total	C	N	O	0	0
			17	9	4	4		
2	D	1	Total	C	N	O	0	0
			17	9	4	4		
2	E	1	Total	C	N	O	0	0
			17	9	4	4		
2	F	1	Total	C	N	O	0	0
			17	9	4	4		
2	G	1	Total	C	N	O	0	0
			17	9	4	4		
2	H	1	Total	C	N	O	0	0
			17	9	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	77	Total	O	0	0
			77	77		
3	B	94	Total	O	0	0
			94	94		
3	C	69	Total	O	0	0
			69	69		
3	D	100	Total	O	0	0
			100	100		
3	E	50	Total	O	0	0
			50	50		
3	F	57	Total	O	0	0
			57	57		
3	G	44	Total	O	0	0
			44	44		
3	H	76	Total	O	0	0
			76	76		


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor 2




- Molecule 1: Glutamate receptor 2




- Molecule 1: Glutamate receptor 2




- Molecule 1: Glutamate receptor 2






- Molecule 1: Glutamate receptor 2





- Molecule 1: Glutamate receptor 2



- Molecule 1: Glutamate receptor 2



- Molecule 1: Glutamate receptor 2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.58Å 97.63Å 100.97Å 81.85° 89.08° 77.89°	Depositor
Resolution (Å)	39.90 – 2.50 39.87 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.8 (39.90-2.50) 96.3 (39.87-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.255 , 0.283 0.253 , 0.285	Depositor DCC
$R_{free}$ test set	4640 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.2	Xtrriage
Anisotropy	0.461	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	16707	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4373e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
CNI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/2041	0.50	1/2746 (0.0%)
1	B	0.40	0/2063	0.49	0/2777
1	C	0.34	0/2026	0.47	0/2728
1	D	0.33	0/2047	0.47	0/2756
1	E	0.31	0/2017	0.46	0/2719
1	F	0.30	0/2042	0.46	0/2748
1	G	0.31	0/2021	0.47	0/2723
1	H	0.30	0/2038	0.46	0/2744
All	All	0.34	0/16295	0.47	1/21941 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	G	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	109	LEU	CA-CB-CG	5.99	129.08	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	153	ALA	Peptide
1	G	163	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2006	0	2023	12	0
1	B	2023	0	2031	8	0
1	C	1990	0	1993	12	0
1	D	2011	0	2014	6	0
1	E	1981	0	1967	9	0
1	F	2006	0	2012	3	0
1	G	1985	0	1995	8	0
1	H	2002	0	2014	5	0
2	A	17	0	2	0	0
2	B	17	0	2	0	0
2	C	17	0	2	0	0
2	D	17	0	2	0	0
2	E	17	0	2	0	0
2	F	17	0	2	0	0
2	G	17	0	2	0	0
2	H	17	0	2	0	0
3	A	77	0	0	2	0
3	B	94	0	0	0	0
3	C	69	0	0	1	0
3	D	100	0	0	0	0
3	E	50	0	0	1	0
3	F	57	0	0	0	0
3	G	44	0	0	0	0
3	H	76	0	0	0	0
All	All	16707	0	16065	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:100:ILE:HD12	1:C:223:ALA:HB1	1.78	0.65
1:E:100:ILE:HD12	1:E:223:ALA:HB1	1.77	0.65
1:A:100:ILE:HD12	1:A:223:ALA:HB1	1.79	0.64
1:G:100:ILE:HD12	1:G:223:ALA:HB1	1.79	0.64
1:H:137:THR:HG22	1:H:191:LEU:HB2	1.80	0.63
1:F:137:THR:HG22	1:F:191:LEU:HB2	1.80	0.62
1:D:137:THR:HG22	1:D:191:LEU:HB2	1.82	0.61
1:G:159:TRP:O	1:G:163:ARG:HG3	2.01	0.60
1:A:225:PRO:HB3	1:D:118:GLY:HA2	1.84	0.59
1:B:137:THR:HG22	1:B:191:LEU:HB2	1.83	0.59
1:B:242:ASN:HB2	1:B:247:LEU:HD12	1.86	0.58
1:C:153:ALA:HB3	3:C:268:HOH:O	2.06	0.55
1:G:216:ASP:OD2	1:G:218:LYS:HE3	2.07	0.54
1:A:216:ASP:OD2	1:A:218:LYS:HE3	2.08	0.53
1:E:216:ASP:OD2	1:E:218:LYS:HE3	2.10	0.52
1:A:153:ALA:HB3	3:A:331:HOH:O	2.09	0.52
1:E:86:ALA:HB3	1:E:223:ALA:HB3	1.90	0.52
1:D:121:ILE:HG21	1:D:127:LEU:HD13	1.92	0.51
1:B:121:ILE:HG21	1:B:127:LEU:HD13	1.93	0.51
1:E:225:PRO:HB3	1:H:118:GLY:HA2	1.93	0.51
1:H:121:ILE:HG21	1:H:127:LEU:HD13	1.93	0.50
1:C:152:ILE:HD13	1:C:155:PHE:CE2	2.47	0.50
1:A:86:ALA:HB3	1:A:223:ALA:HB3	1.95	0.49
1:C:152:ILE:O	1:C:153:ALA:C	2.51	0.49
1:C:88:ALA:HB1	1:C:89:PRO:HD2	1.94	0.49
1:C:216:ASP:OD2	1:C:218:LYS:HE3	2.12	0.48
1:F:121:ILE:HG21	1:F:127:LEU:HD13	1.94	0.48
1:A:88:ALA:HB1	1:A:89:PRO:HD2	1.96	0.48
1:A:131:THR:HA	3:A:323:HOH:O	2.14	0.47
1:G:86:ALA:HB3	1:G:223:ALA:HB3	1.97	0.47
1:G:88:ALA:HB1	1:G:89:PRO:HD2	1.97	0.46
1:E:88:ALA:HB1	1:E:89:PRO:HD2	1.97	0.46
1:A:121:ILE:HD13	1:A:126:ASP:HB3	1.98	0.46
1:F:88:ALA:HB1	1:F:89:PRO:HD2	1.99	0.45
1:B:68:THR:O	1:B:69:LYS:CB	2.63	0.44
1:D:88:ALA:HB1	1:D:89:PRO:HD2	1.99	0.44
1:B:88:ALA:HB1	1:B:89:PRO:HD2	2.00	0.44
1:C:86:ALA:HB3	1:C:223:ALA:HB3	2.00	0.43
1:A:171:VAL:HG11	1:A:177:GLY:HA2	2.00	0.43
1:C:105:PRO:HA	1:C:220:TYR:O	2.19	0.43
1:A:246:LEU:O	1:A:250:LEU:HB2	2.19	0.43
1:G:171:VAL:HG11	1:G:177:GLY:HA2	2.00	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:130:GLN:HE22	1:E:133:ILE:HB	1.83	0.43
1:C:152:ILE:HD13	1:C:155:PHE:HE2	1.84	0.43
1:H:88:ALA:HB1	1:H:89:PRO:HD2	2.00	0.43
1:C:171:VAL:HG11	1:C:177:GLY:HA2	1.99	0.43
1:C:246:LEU:O	1:C:250:LEU:HB2	2.19	0.43
1:G:246:LEU:O	1:G:250:LEU:HB2	2.19	0.42
1:H:6:VAL:HG13	1:H:84:ASP:HB2	2.01	0.42
1:G:130:GLN:HE22	1:G:133:ILE:HB	1.85	0.42
1:E:171:VAL:HG11	1:E:177:GLY:HA2	2.01	0.42
1:E:246:LEU:O	1:E:250:LEU:HB2	2.19	0.42
1:A:159:TRP:O	1:A:163:ARG:HG3	2.20	0.41
1:B:242:ASN:CB	1:B:247:LEU:HD12	2.51	0.41
1:E:130:GLN:NE2	3:E:309:HOH:O	2.53	0.41
1:C:130:GLN:HE22	1:C:133:ILE:HB	1.86	0.41
1:D:127:LEU:HG	1:D:135:TYR:CD1	2.56	0.41
1:A:105:PRO:HA	1:A:220:TYR:O	2.20	0.41
1:B:127:LEU:HG	1:B:135:TYR:CD1	2.57	0.40
1:B:6:VAL:HG13	1:B:84:ASP:HB2	2.02	0.40
1:D:6:VAL:HG13	1:D:84:ASP:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/261 (98%)	249 (97%)	7 (3%)	0	100	100
1	B	258/261 (99%)	256 (99%)	2 (1%)	0	100	100
1	C	255/261 (98%)	250 (98%)	4 (2%)	1 (0%)	34	54
1	D	257/261 (98%)	255 (99%)	2 (1%)	0	100	100
1	E	256/261 (98%)	251 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	256/261 (98%)	253 (99%)	3 (1%)	0	100	100
1	G	255/261 (98%)	248 (97%)	7 (3%)	0	100	100
1	H	256/261 (98%)	254 (99%)	2 (1%)	0	100	100
All	All	2049/2088 (98%)	2016 (98%)	32 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	153	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/218 (98%)	210 (98%)	4 (2%)	57	80
1	B	216/218 (99%)	212 (98%)	4 (2%)	57	80
1	C	210/218 (96%)	204 (97%)	6 (3%)	42	69
1	D	213/218 (98%)	206 (97%)	7 (3%)	38	64
1	E	207/218 (95%)	201 (97%)	6 (3%)	42	69
1	F	213/218 (98%)	206 (97%)	7 (3%)	38	64
1	G	209/218 (96%)	204 (98%)	5 (2%)	49	74
1	H	212/218 (97%)	205 (97%)	7 (3%)	38	64
All	All	1694/1744 (97%)	1648 (97%)	46 (3%)	44	71

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	LEU
1	A	130	GLN
1	A	131	THR
1	A	163	ARG
1	B	109	LEU

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	193	GLU
1	B	249	LYS
1	B	250	LEU
1	C	109	LEU
1	C	122	GLU
1	C	130	GLN
1	C	131	THR
1	C	163	ARG
1	C	203	ARG
1	D	69	LYS
1	D	109	LEU
1	D	163	ARG
1	D	193	GLU
1	D	246	LEU
1	D	249	LYS
1	D	250	LEU
1	E	109	LEU
1	E	122	GLU
1	E	130	GLN
1	E	131	THR
1	E	163	ARG
1	E	203	ARG
1	F	69	LYS
1	F	109	LEU
1	F	163	ARG
1	F	193	GLU
1	F	246	LEU
1	F	249	LYS
1	F	250	LEU
1	G	109	LEU
1	G	122	GLU
1	G	130	GLN
1	G	131	THR
1	G	203	ARG
1	H	69	LYS
1	H	109	LEU
1	H	163	ARG
1	H	193	GLU
1	H	246	LEU
1	H	249	LYS
1	H	250	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are

no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CNI	E	262	-	16,18,18	2.20	1 (6%)	15,26,26	0.93	0
2	CNI	A	262	-	16,18,18	2.26	1 (6%)	15,26,26	0.99	0
2	CNI	F	262	-	16,18,18	2.24	1 (6%)	15,26,26	0.88	0
2	CNI	H	262	-	16,18,18	2.24	1 (6%)	15,26,26	1.03	0
2	CNI	C	262	-	16,18,18	2.22	1 (6%)	15,26,26	1.08	2 (13%)
2	CNI	G	262	-	16,18,18	2.20	1 (6%)	15,26,26	0.93	0
2	CNI	B	262	-	16,18,18	2.26	1 (6%)	15,26,26	1.11	1 (6%)
2	CNI	D	262	-	16,18,18	2.25	1 (6%)	15,26,26	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CNI	E	262	-	-	2/2/18/18	0/2/2/2
2	CNI	A	262	-	-	2/2/18/18	0/2/2/2
2	CNI	F	262	-	-	2/2/18/18	0/2/2/2
2	CNI	H	262	-	-	2/2/18/18	0/2/2/2
2	CNI	C	262	-	-	0/2/18/18	0/2/2/2
2	CNI	G	262	-	-	0/2/18/18	0/2/2/2
2	CNI	B	262	-	-	2/2/18/18	0/2/2/2
2	CNI	D	262	-	-	2/2/18/18	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	262	CNI	C-C8	-7.74	1.28	1.43
2	B	262	CNI	C-C8	-7.64	1.28	1.43
2	H	262	CNI	C-C8	-7.61	1.28	1.43
2	F	262	CNI	C-C8	-7.58	1.28	1.43
2	C	262	CNI	C-C8	-7.56	1.28	1.43
2	D	262	CNI	C-C8	-7.55	1.28	1.43
2	E	262	CNI	C-C8	-7.47	1.29	1.43
2	G	262	CNI	C-C8	-7.39	1.29	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	262	CNI	C8-C-N17	-2.54	173.23	177.72
2	C	262	CNI	C5-C7-C8	-2.27	119.12	122.15
2	C	262	CNI	C8-C-N17	-2.05	174.10	177.72

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	262	CNI	C5-C7-N3-O3
2	B	262	CNI	C8-C7-N3-O3
2	D	262	CNI	C8-C7-N3-O3
2	F	262	CNI	C5-C7-N3-O3
2	F	262	CNI	C8-C7-N3-O3
2	H	262	CNI	C5-C7-N3-O3
2	H	262	CNI	C8-C7-N3-O3
2	D	262	CNI	C5-C7-N3-O3
2	A	262	CNI	C5-C7-N3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	262	CNI	C8-C7-N3-O3
2	E	262	CNI	C5-C7-N3-O3
2	E	262	CNI	C8-C7-N3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/261 (98%)	0.36	10 (3%) 39 42	16, 23, 28, 29	0
1	B	258/261 (98%)	0.20	3 (1%) 79 80	14, 19, 23, 25	0
1	C	257/261 (98%)	0.29	6 (2%) 60 63	16, 23, 28, 29	0
1	D	259/261 (99%)	0.20	3 (1%) 79 80	14, 19, 23, 26	0
1	E	258/261 (98%)	0.44	11 (4%) 35 38	16, 23, 28, 28	0
1	F	258/261 (98%)	0.42	12 (4%) 31 33	14, 19, 23, 25	0
1	G	257/261 (98%)	0.44	14 (5%) 25 27	16, 23, 28, 29	0
1	H	258/261 (98%)	0.33	7 (2%) 54 58	14, 19, 23, 25	0
All	All	2063/2088 (98%)	0.33	66 (3%) 47 51	14, 20, 27, 29	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	24	GLU	4.8
1	E	129	LYS	3.5
1	C	259	GLY	3.4
1	H	219	GLY	3.3
1	H	24	GLU	3.3
1	F	48	GLY	3.2
1	C	125	GLU	3.2
1	F	131	THR	3.1
1	C	25	MET	2.9
1	F	259	GLY	2.9
1	G	46	HIS	2.8
1	H	25	MET	2.8
1	B	66	ALA	2.8
1	F	153	ALA	2.7
1	G	221	GLY	2.7
1	F	66	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	131	THR	2.6
1	E	125	GLU	2.6
1	A	66	ALA	2.6
1	G	186	GLY	2.6
1	H	206	CYS	2.6
1	B	131	THR	2.6
1	E	66	ALA	2.6
1	G	24	GLU	2.6
1	A	131	THR	2.5
1	G	105	PRO	2.5
1	F	219	GLY	2.5
1	A	5	THR	2.4
1	D	219	GLY	2.4
1	E	25	MET	2.4
1	A	261	CYS	2.4
1	H	125	GLU	2.4
1	F	205	PRO	2.4
1	H	131	THR	2.3
1	A	67	ASP	2.3
1	C	24	GLU	2.3
1	A	186	GLY	2.3
1	G	26	LEU	2.3
1	F	46	HIS	2.3
1	G	6	VAL	2.3
1	G	129	LYS	2.3
1	A	4	LYS	2.3
1	H	105	PRO	2.2
1	E	156	ASP	2.2
1	G	131	THR	2.2
1	E	23	HIS	2.2
1	E	130	GLN	2.2
1	F	107	MET	2.2
1	A	69	LYS	2.2
1	G	50	LYS	2.2
1	E	105	PRO	2.1
1	B	261	CYS	2.1
1	E	221	GLY	2.1
1	F	108	SER	2.1
1	C	22	ASN	2.1
1	G	259	GLY	2.1
1	E	6	VAL	2.1
1	F	105	PRO	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	3	ASN	2.1
1	F	25	MET	2.1
1	G	7	VAL	2.1
1	G	230	LEU	2.1
1	A	133	ILE	2.0
1	C	21	LYS	2.0
1	A	142	SER	2.0
1	G	49	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CNI	B	262	17/17	0.84	0.22	43,43,45,45	0
2	CNI	E	262	17/17	0.84	0.18	48,48,49,50	0
2	CNI	F	262	17/17	0.86	0.20	43,43,44,44	0
2	CNI	A	262	17/17	0.88	0.17	38,38,39,39	0
2	CNI	H	262	17/17	0.90	0.19	49,49,50,50	0
2	CNI	C	262	17/17	0.91	0.13	35,35,36,37	0
2	CNI	D	262	17/17	0.92	0.16	31,32,34,34	0
2	CNI	G	262	17/17	0.94	0.13	43,43,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.