



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2023 – 10:08 PM EDT

PDB ID : 3D74
Title : Crystal structure of a pheromone binding protein mutant D35A, from *Apis mellifera*, soaked at pH 5.5
Authors : Pesenti, M.E.; Spinelli, S.; Bezirard, V.; Briand, L.; Pernollet, J.C.; Tegoni, M.; Cambillau, C.
Deposited on : 2008-05-20
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	119	 81% 18% .
1	B	119	 89% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pheromone-binding protein ASP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	119	Total 929	C 582	N 146	O 191	S 10	0	3	0
1	B	119	Total 916	C 572	N 146	O 188	S 10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	ALA	ASP	engineered mutation	UNP Q9U9J6
B	35	ALA	ASP	engineered mutation	UNP Q9U9J6

- Molecule 2 is N-BUTYL-BENZENESULFONAMIDE (three-letter code: NBB) (formula: $C_{10}H_{15}NO_2S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total 14	C 10	N 1	O 2	S 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			14	10	1	2	1		
2	B	1	Total	C	N	O	S	0	0
			14	10	1	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		
3	B	63	Total	O	0	0
			63	63		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pheromone-binding protein ASP1

Chain A:  81% 18%



- Molecule 1: Pheromone-binding protein ASP1

Chain B:  89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	61.13Å 60.77Å 56.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.10 30.38 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (15.00-2.10) 99.7 (30.38-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.175 , 0.229 0.221 , 0.255	Depositor DCC
R_{free} test set	956 reflections (7.52%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.713	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2009	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.4636e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NBB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/955	0.66	0/1300
1	B	0.65	1/933 (0.1%)	0.69	0/1270
All	All	0.61	1/1888 (0.1%)	0.67	0/2570

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	37	GLY	C-N	5.91	1.47	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	929	0	892	21	0
1	B	916	0	872	11	0
2	A	14	0	15	2	0
2	B	28	0	30	3	0
3	A	59	0	0	3	0
3	B	63	0	0	1	0
All	All	2009	0	1809	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:MET:HG2	3:A:168:HOH:O	1.83	0.79
1:A:73:LEU:O	1:A:73:LEU:HD23	1.88	0.73
1:B:1:ALA:HB1	3:B:170:HOH:O	1.90	0.72
1:B:13:VAL:HG12	1:B:56:PHE:CE2	2.28	0.69
1:A:41:ASN:HD21	1:A:99:ASN:HD21	1.42	0.66
1:A:41:ASN:HD21	1:A:99:ASN:ND2	1.95	0.65
1:A:116:TRP:CE3	2:A:120:NBB:H142	2.35	0.62
1:B:13:VAL:HG12	1:B:56:PHE:CZ	2.37	0.60
1:A:58:LEU:CD2	1:A:70:MET:HA	2.33	0.58
1:A:73:LEU:HD22	3:A:157:HOH:O	2.05	0.57
1:A:58:LEU:HD21	1:A:70:MET:HA	1.87	0.56
1:A:67:GLU:HG2	1:A:86:MET:HE2	1.86	0.56
1:A:85[A]:VAL:HG11	1:A:112:ALA:HB3	1.87	0.56
1:A:85[A]:VAL:HA	1:A:88:LYS:HE2	1.89	0.55
1:A:85[B]:VAL:HA	1:A:88:LYS:HE2	1.89	0.55
1:B:13:VAL:HG12	1:B:56:PHE:HE2	1.73	0.52
1:B:1:ALA:HB3	1:B:2:PRO:HD3	1.93	0.50
1:A:85[A]:VAL:HG11	1:A:112:ALA:CB	2.43	0.48
1:B:12:LEU:HD22	2:B:121:NBB:H141	1.96	0.47
1:A:109:GLN:HA	1:A:113:PRO:HA	1.97	0.47
1:A:118:VAL:HG13	2:A:120:NBB:O1S	2.14	0.47
1:A:34:VAL:HG11	1:A:118:VAL:CG2	2.45	0.47
1:A:82:ALA:O	1:A:86:MET:HG3	2.16	0.46
1:A:9:VAL:HG12	1:A:119:ILE:HD11	1.98	0.45
1:A:70:MET:HB3	1:A:86:MET:HE1	1.98	0.44
1:B:112:ALA:HB1	1:B:115:VAL:CG2	2.47	0.44
1:A:8[A]:GLU:HG3	3:A:131:HOH:O	2.20	0.42
1:A:19:ARG:O	1:A:23:GLU:HG3	2.19	0.42
1:B:41:ASN:HD21	1:B:99:ASN:HD21	1.67	0.42
1:B:56:PHE:HE2	2:B:120:NBB:H1	1.84	0.42
1:B:48:TYR:CZ	1:B:52:LEU:HD22	2.55	0.41
1:B:12:LEU:HD22	2:B:121:NBB:C14	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/119 (101%)	120 (100%)	0	0	100	100
1	B	117/119 (98%)	117 (100%)	0	0	100	100
All	All	237/238 (100%)	237 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/104 (103%)	104 (97%)	3 (3%)	43	47
1	B	104/104 (100%)	103 (99%)	1 (1%)	76	82
All	All	211/208 (101%)	207 (98%)	4 (2%)	57	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	THR
1	A	76	ASP
1	A	88	LYS
1	B	111	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	79	GLN
1	A	99	ASN
1	A	103	ASN
1	B	38	ASN
1	B	77	GLN
1	B	99	ASN
1	B	103	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NBB	A	120	-	14,14,14	3.05	4 (28%)	18,18,18	2.42	4 (22%)
2	NBB	B	121	-	14,14,14	3.41	4 (28%)	18,18,18	2.58	6 (33%)
2	NBB	B	120	-	14,14,14	3.55	4 (28%)	18,18,18	2.72	4 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NBB	A	120	-	-	1/12/12/12	0/1/1/1
2	NBB	B	121	-	-	0/12/12/12	0/1/1/1
2	NBB	B	120	-	-	2/12/12/12	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	120	NBB	C4-S	-9.86	1.61	1.76
2	A	120	NBB	C4-S	-8.99	1.62	1.76
2	B	121	NBB	C4-S	-8.79	1.62	1.76
2	B	121	NBB	S-N10	-5.68	1.52	1.61
2	B	121	NBB	O1S-S	5.34	1.49	1.43
2	B	120	NBB	S-N10	-5.26	1.53	1.61
2	A	120	NBB	S-N10	-5.20	1.53	1.61
2	B	120	NBB	O2S-S	4.94	1.49	1.43
2	B	120	NBB	O1S-S	4.92	1.49	1.43
2	B	121	NBB	O2S-S	4.00	1.48	1.43
2	A	120	NBB	O2S-S	3.61	1.47	1.43
2	A	120	NBB	O1S-S	2.59	1.46	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	120	NBB	O2S-S-O1S	-9.46	107.92	119.55
2	B	121	NBB	O2S-S-O1S	-8.37	109.27	119.55
2	A	120	NBB	O2S-S-O1S	-7.23	110.67	119.55
2	A	120	NBB	O1S-S-N10	4.93	114.74	107.04
2	B	121	NBB	C4-S-N10	4.72	114.10	107.55
2	B	120	NBB	O1S-S-N10	3.88	113.10	107.04
2	B	120	NBB	O1S-S-C4	3.19	111.90	107.97
2	A	120	NBB	O2S-S-C4	3.13	111.82	107.97
2	B	120	NBB	C2-C3-C4	2.57	121.62	118.95
2	B	121	NBB	C12-C11-N10	-2.34	105.91	111.03
2	B	121	NBB	C2-C3-C4	2.29	121.34	118.95
2	B	121	NBB	O1S-S-N10	2.20	110.47	107.04
2	B	121	NBB	O1S-S-C4	-2.16	105.31	107.97
2	A	120	NBB	C2-C3-C4	2.05	121.08	118.95

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	120	NBB	C11-C12-C13-C14
2	B	120	NBB	C5-C4-S-O1S
2	B	120	NBB	C3-C4-S-O1S

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	120	NBB	2	0
2	B	121	NBB	2	0
2	B	120	NBB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.