



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 10:41 AM EDT

PDB ID : 3HPO
Title : Crystal structure of fragment DNA polymerase I from *Bacillus stearothermophilus* Y714S mutant bound to G:T mismatch
Authors : Wu, E.Y.; Beese, L.S.
Deposited on : 2009-06-04
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	580	
2	B	9	
3	C	10	
4	D	2	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I, large fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	580	4679	2973	812	876	18	0	6	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	329	ALA	ASP	engineered mutation	PDB 3HPO
A	714	SER	TYR	engineered mutation	PDB 3HPO


- Molecule 2 is a DNA chain called 5'-D(*CP*GP*AP*TP*CP*AP*CP*GP*(DOC))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	9	178	86	34	50	8	0	0	0

- Molecule 3 is a DNA chain called 5'-D(P*GP*GP*CP*GP*TP*GP*AP*TP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	10	209	98	40	61	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
4	D	2	23	12	11	0	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0

- Molecule 6 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: C₁₀H₁₇N₂O₁₄P₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O P 29 10 2 14 3	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	652	Total O 652 652	0	0
8	B	29	Total O 29 29	0	0
8	C	43	Total O 43 43	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I, large fragment



- Molecule 2: 5'-D(*CP*GP*AP*TP*CP*AP*CP*GP*(DOC))-3'



- Molecule 3: 5'-D(P*GP*GP*CP*GP*TP*GP*AP*TP*CP*G)-3'



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.21Å 93.73Å 105.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.98 – 1.75 45.97 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.98-1.75) 92.1 (45.97-1.75)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 1.75Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.201 , 0.241 0.200 , 0.240	Depositor DCC
R_{free} test set	4178 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtrriage
Anisotropy	0.186	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.021 for k,h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5848	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CME, MG, GLC, DOC, FRU, TTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/4763	0.54	0/6434
2	B	0.82	0/179	1.61	4/274 (1.5%)
3	C	0.75	0/234	1.42	4/360 (1.1%)
All	All	0.44	0/5176	0.68	8/7068 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	27	DC	C4'-C3'-C2'	-9.47	94.58	103.10
2	B	27	DC	O4'-C4'-C3'	-8.10	101.14	106.00
3	C	10	DT	C1'-O4'-C4'	-7.49	102.61	110.10
3	C	8	DG	O4'-C1'-N9	7.04	112.93	108.00
3	C	7	DT	C5-C4-O4	-5.98	120.72	124.90
2	B	27	DC	O5'-P-OP1	5.95	117.84	110.70
3	C	7	DT	N3-C4-O4	5.61	123.27	119.90
2	B	27	DC	O5'-P-OP2	-5.52	100.73	105.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4679	0	4746	50	0
2	B	178	0	102	4	0
3	C	209	0	113	0	0
4	D	23	0	21	1	0
5	A	1	0	0	0	0
6	A	29	0	13	2	0
7	A	5	0	0	0	0
8	A	652	0	0	2	0
8	B	29	0	0	0	0
8	C	43	0	0	0	0
All	All	5848	0	4995	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:704:GLN:CG	1:A:725:LEU:HD22	1.70	1.21
1:A:704:GLN:HG3	1:A:725:LEU:CD2	1.72	1.20
1:A:704:GLN:HG3	1:A:725:LEU:HD22	1.05	1.01
1:A:719:TYR:HB2	1:A:729:ARG:HE	1.36	0.88
1:A:692:VAL:HB	1:A:696:GLU:HG3	1.56	0.87
1:A:629:ARG:HH22	2:B:29:DOC:H5	1.45	0.81
1:A:629:ARG:NH2	2:B:29:DOC:H5	1.99	0.77
1:A:551:LYS:HD3	1:A:551:LYS:C	2.09	0.73
1:A:408:ASP:HB2	4:D:2:FRU:H11	1.72	0.71
1:A:728:SER:OG	1:A:731:GLU:HG3	1.91	0.70
1:A:713:VAL:HG13	8:A:107:HOH:O	1.90	0.70
1:A:670[A]:ASN:OD1	8:A:897:HOH:O	2.10	0.67
1:A:591:LEU:O	1:A:595:VAL:HG23	1.95	0.66
1:A:719:TYR:HB2	1:A:729:ARG:NE	2.12	0.60
1:A:662:LEU:CD2	1:A:713:VAL:HG11	2.32	0.59
1:A:692:VAL:HB	1:A:696:GLU:CG	2.32	0.57
1:A:629:ARG:NH1	2:B:28:DG:OP2	2.37	0.57
1:A:565:ALA:HB3	1:A:566:PRO:HD3	1.88	0.56
1:A:782:ASN:O	1:A:786:PHE:HD2	1.89	0.56
1:A:584:GLN:O	1:A:589:GLU:HG3	2.06	0.56
1:A:510:GLN:O	1:A:514:VAL:HG23	2.05	0.55
1:A:654:TYR:HB3	1:A:657:ILE:HB	1.88	0.54
1:A:719:TYR:O	1:A:723:GLN:HG2	2.07	0.54
1:A:656[A]:GLN:OE1	6:A:201:TTP:O1B	2.28	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:662:LEU:HD22	1:A:713:VAL:HG11	1.92	0.51
1:A:515:GLU:HG2	1:A:519:TYR:CE2	2.46	0.50
1:A:349:GLU:H	1:A:349:GLU:CD	2.15	0.49
1:A:575:LEU:O	1:A:579:GLN:HG3	2.13	0.48
1:A:552:THR:HG22	2:B:24:DT:OP1	2.13	0.48
1:A:715:GLY:HA3	1:A:789:ARG:HG2	1.96	0.48
1:A:794:THR:N	1:A:795:PRO:CD	2.77	0.47
1:A:495[A]:VAL:HG11	1:A:500:LEU:HD11	1.96	0.47
1:A:329:ALA:HB3	1:A:596:ARG:HH22	1.79	0.47
1:A:719:TYR:HB2	1:A:729:ARG:HH21	1.79	0.47
1:A:494:LYS:HD2	1:A:600:LYS:HB2	1.98	0.45
1:A:682:HIS:CD2	1:A:706:LYS:HG2	2.51	0.45
1:A:704:GLN:CD	1:A:725:LEU:HD22	2.33	0.45
1:A:671:LEU:HD12	1:A:681:ILE:HD11	1.97	0.45
1:A:656[A]:GLN:NE2	6:A:201:TTP:O3G	2.50	0.45
1:A:534:LEU:HD11	1:A:574:ILE:HD13	1.99	0.44
1:A:656[A]:GLN:HE22	1:A:682:HIS:CD2	2.35	0.44
1:A:820:LEU:HD21	1:A:843:ARG:CZ	2.48	0.44
1:A:459:ARG:N	1:A:460:PRO:HD2	2.34	0.43
1:A:565:ALA:N	1:A:566:PRO:CD	2.82	0.43
1:A:662:LEU:HD21	1:A:713:VAL:HG11	1.99	0.42
1:A:828:VAL:HB	1:A:831:GLU:HG2	2.02	0.42
1:A:704:GLN:CG	1:A:725:LEU:CD2	2.55	0.41
1:A:682:HIS:CG	1:A:706:LYS:HG2	2.56	0.41
1:A:551:LYS:C	1:A:551:LYS:CD	2.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	583/580 (100%)	567 (97%)	14 (2%)	2 (0%)	41 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	LYS
1	A	628	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	500/494 (101%)	490 (98%)	10 (2%)	55 34

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	356	GLN
1	A	551	LYS
1	A	591	LEU
1	A	623	LEU
1	A	695	ASP
1	A	704	GLN
1	A	724	ASN
1	A	779	ARG
1	A	844	LEU
1	A	862	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	704	GLN
1	A	782	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DOC	B	29	3,2	16,19,20	0.41	0	20,26,29	0.70	0
1	CME	A	388	1	8,9,10	0.91	0	5,9,11	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	B	29	3,2	-	0/7/18/19	0/2/2/2
1	CME	A	388	1	-	1/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	388	CME	SD-CE-CZ-OH

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	29	DOC	2	0

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	D	1	4	11,11,12	0.60	0	15,15,17	0.95	1 (6%)
4	FRU	D	2	4	11,12,12	0.54	0	10,18,18	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	D	1	4	-	0/2/19/22	0/1/1/1
4	FRU	D	2	4	-	0/5/24/24	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	GLC	C1-O5-C5	2.71	115.86	112.19

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	2	FRU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	SO4	A	912	-	4,4,4	0.13	0	6,6,6	0.24	0
6	TTP	A	201	5	26,30,30	1.23	3 (11%)	39,47,47	2.11	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TTP	A	201	5	-	3/22/34/34	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	201	TTP	C6-C5	2.59	1.38	1.34
6	A	201	TTP	C4-C5	2.41	1.48	1.44
6	A	201	TTP	C6-N1	-2.26	1.34	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	201	TTP	C4-N3-C2	-5.43	120.32	127.35
6	A	201	TTP	C5-C4-N3	5.14	119.70	115.31
6	A	201	TTP	O4-C4-C5	-4.82	119.31	124.90
6	A	201	TTP	N3-C2-N1	4.14	120.39	114.89
6	A	201	TTP	C5-C6-N1	-4.06	119.16	123.34
6	A	201	TTP	O2-C2-N1	-3.43	118.23	122.79
6	A	201	TTP	C2'-C1'-N1	-3.40	105.95	113.77
6	A	201	TTP	C5M-C5-C4	2.81	121.86	118.77
6	A	201	TTP	O3G-PG-O3B	2.35	112.51	104.64
6	A	201	TTP	PB-O3B-PG	-2.08	125.67	132.83
6	A	201	TTP	C5M-C5-C6	-2.02	120.15	122.85

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	201	TTP	PB-O3B-PG-O1G
6	A	201	TTP	PB-O3A-PA-O2A
6	A	201	TTP	PB-O3A-PA-O1A


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	201	TTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	579/580 (99%)	0.41	37 (6%) 19 25	11, 22, 40, 49	0
2	B	8/9 (88%)	0.07	0 100 100	15, 25, 47, 56	0
3	C	10/10 (100%)	-0.24	0 100 100	14, 20, 33, 42	0
All	All	597/599 (99%)	0.39	37 (6%) 20 26	11, 22, 41, 56	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	ALA	8.1
1	A	719	TYR	6.8
1	A	433	ALA	6.5
1	A	519	TYR	5.4
1	A	552	THR	5.0
1	A	434	LYS	4.8
1	A	551	LYS	4.5
1	A	554	TYR	4.4
1	A	516	GLN	3.9
1	A	511	LEU	3.9
1	A	431	LYS	3.6
1	A	531	PRO	3.3
1	A	512	GLY	3.2
1	A	298	LYS	3.0
1	A	513	THR	3.0
1	A	729	ARG	3.0
1	A	723	GLN	2.9
1	A	525	GLU	2.8
1	A	505	LYS	2.8
1	A	730	LYS	2.6
1	A	508	ALA	2.5
1	A	520	GLU	2.5
1	A	523	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	781	PHE	2.5
1	A	524	GLN	2.4
1	A	695	ASP	2.4
1	A	517	ARG	2.3
1	A	786	PHE	2.3
1	A	521	LEU	2.2
1	A	509	GLU	2.2
1	A	712	ILE	2.2
1	A	514	VAL	2.1
1	A	532	LYS	2.1
1	A	549	LYS	2.1
1	A	697	VAL	2.0
1	A	703	ARG	2.0
1	A	713	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	388	10/11	0.97	0.10	19,23,39,41	0
2	DOC	B	29	18/19	0.98	0.10	13,14,17,21	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FRU	D	2	12/12	0.92	0.10	22,25,25,26	0
4	GLC	D	1	11/12	0.95	0.09	24,25,27,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SO4	A	912	5/5	0.96	0.17	45,46,47,47	0
6	TTP	A	201	29/29	0.97	0.12	14,17,20,26	29
5	MG	A	200	1/1	0.99	0.05	18,18,18,18	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.