



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 08:56 PM EDT

PDB ID : 3L2B
Title : Crystal structure of the CBS and DRTGG domains of the regulatory region of *Clostridium perfringens* pyrophosphatase complexed with activator, diadenosine tetraphosphate
Authors : Tuominen, H.; Salminen, A.; Oksanen, E.; Jamsen, J.; Heikkila, O.; Lehtio, L.; Magretova, N.N.; Goldman, A.; Baykov, A.A.; Lahti, R.
Deposited on : 2009-12-15
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	 9% 87% 7% • 5%
1	B	245	 16% 85% 9% 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Probable manganese-dependent inorganic pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	233	Total	C	N	O	S	1	1	0
			1796	1134	294	359	9			
1	B	230	Total	C	N	O	S	0	1	0
			1773	1118	291	356	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	GLY	-	expression tag	UNP Q8XIQ9
A	63	SER	-	expression tag	UNP Q8XIQ9
A	64	HIS	-	expression tag	UNP Q8XIQ9
A	65	MET	-	expression tag	UNP Q8XIQ9
B	62	GLY	-	expression tag	UNP Q8XIQ9
B	63	SER	-	expression tag	UNP Q8XIQ9
B	64	HIS	-	expression tag	UNP Q8XIQ9
B	65	MET	-	expression tag	UNP Q8XIQ9

- Molecule 2 is BIS(ADENOSINE)-5'-TETRAPHOSPHATE (three-letter code: B4P) (formula: C₂₀H₂₈N₁₀O₁₉P₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	20	10	19	4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	41	Total O 41 41	0	0
3	B	33	Total O 33 33	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable manganese-dependent inorganic pyrophosphatase



- Molecule 1: Probable manganese-dependent inorganic pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.85Å 71.76Å 116.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 2.27 19.99 – 2.27	Depositor EDS
% Data completeness (in resolution range)	97.2 (19.99-2.27) 97.3 (19.99-2.27)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 2.28Å)	Xtrriage
Refinement program	REFMAC 5.4.0078	Depositor
R, R_{free}	0.209 , 0.259 0.213 , 0.255	Depositor DCC
R_{free} test set	1298 reflections (5.29%)	wwPDB-VP
Wilson B-factor (Å ²)	51.2	Xtrriage
Anisotropy	0.192	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3696	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: B4P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.81	2/1822 (0.1%)	0.89	4/2474 (0.2%)
1	B	0.63	0/1798	0.69	0/2441
All	All	0.72	2/3620 (0.1%)	0.80	4/4915 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124[A]	TYR	CE2-CZ	13.97	1.56	1.38
1	A	124[B]	TYR	CE2-CZ	13.97	1.56	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124[A]	TYR	CZ-CE2-CD2	-16.75	104.72	119.80
1	A	124[B]	TYR	CZ-CE2-CD2	-16.75	104.72	119.80
1	A	124[A]	TYR	CG-CD2-CE2	8.08	127.76	121.30
1	A	124[B]	TYR	CG-CD2-CE2	8.08	127.76	121.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1796	0	1832	20	0
1	B	1773	0	1812	15	0
2	A	53	0	24	1	0
3	A	41	0	0	0	0
3	B	33	0	0	0	0
All	All	3696	0	3668	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124[B]:TYR:CD2	1:B:124[B]:TYR:CD2	2.65	0.84
1:A:67:LYS:HE2	1:A:265:LEU:HD13	1.62	0.79
1:A:174:SER:O	1:A:177:GLU:HG2	1.92	0.68
1:A:129:ASP:OD1	1:A:131:ASN:HB3	1.94	0.68
1:A:73:LEU:HD11	1:A:266:VAL:HG22	1.78	0.66
1:A:124[B]:TYR:CD2	1:B:124[B]:TYR:CE2	2.88	0.61
1:A:291:VAL:O	1:A:291:VAL:CG1	2.48	0.60
1:A:132:ILE:HD12	1:A:133:LEU:N	2.18	0.58
1:A:129:ASP:HB3	1:A:132:ILE:HG23	1.89	0.55
1:A:132:ILE:HD12	1:A:133:LEU:H	1.74	0.52
1:A:129:ASP:OD1	1:A:131:ASN:CB	2.61	0.49
1:A:291:VAL:O	1:A:291:VAL:HG13	2.12	0.48
1:B:119:ASN:HB3	1:B:251:VAL:CG1	2.43	0.48
1:A:124[B]:TYR:CE2	1:B:124[B]:TYR:CD2	3.02	0.47
1:A:175:LEU:HD11	1:A:193:ILE:HD12	1.97	0.47
1:A:73:LEU:HD11	1:A:266:VAL:CG2	2.45	0.45
1:B:184:ILE:HD12	1:B:184:ILE:N	2.32	0.45
1:A:73:LEU:HD13	1:A:294:ILE:HD11	2.00	0.44
1:A:184:ILE:N	1:A:184:ILE:HD12	2.33	0.44
1:B:85:VAL:HG13	1:B:89:MET:HB2	1.98	0.44
1:B:258:VAL:HG21	1:B:276:THR:HG21	2.00	0.43
1:A:236:PHE:HB2	1:B:167:VAL:HG23	2.00	0.43
2:A:1:B4P:H2B	1:B:257:LEU:HA	2.01	0.43
1:A:124[A]:TYR:CD2	1:B:124[A]:TYR:CD2	3.07	0.43
1:B:114:MET:HE3	1:B:282:PRO:HG3	2.02	0.42
1:B:128:TRP:CZ3	1:B:130:SER:HA	2.55	0.42
1:B:104:VAL:HG21	1:B:248:VAL:CG1	2.50	0.41
1:B:78:ILE:HG23	1:B:78:ILE:O	2.21	0.41
1:B:276:THR:O	1:B:277:ARG:CB	2.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:VAL:O	1:A:226:ILE:HD12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/245 (95%)	226 (97%)	6 (3%)	0	100	100
1	B	229/245 (94%)	222 (97%)	6 (3%)	1 (0%)	34	40
All	All	461/490 (94%)	448 (97%)	12 (3%)	1 (0%)	47	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	127	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/216 (95%)	204 (100%)	1 (0%)	88	94
1	B	202/216 (94%)	199 (98%)	3 (2%)	65	77
All	All	407/432 (94%)	403 (99%)	4 (1%)	76	86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	ILE
1	B	77	LYS
1	B	170	MET
1	B	296	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	131	ASN
1	B	225	ASN
1	B	280	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	B4P	A	1	-	46,58,58	1.11	3 (6%)	49,91,91	1.46	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B4P	A	1	-	-	2/30/70/70	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	B4P	O4E-C1E	3.39	1.45	1.41
2	A	1	B4P	C5A-C4A	2.69	1.48	1.40
2	A	1	B4P	C5B-C4B	2.15	1.46	1.40

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	B4P	PD-O3G-PG	-3.84	119.66	132.83
2	A	1	B4P	N3A-C2A-N1A	-3.50	123.21	128.68
2	A	1	B4P	N3B-C2B-N1B	-3.39	123.39	128.68
2	A	1	B4P	PG-O3B-PB	-2.98	122.59	132.83
2	A	1	B4P	C4B-C5B-N7B	-2.93	106.35	109.40
2	A	1	B4P	C4A-C5A-N7A	-2.82	106.46	109.40
2	A	1	B4P	C3E-C2E-C1E	2.58	104.87	100.98
2	A	1	B4P	C3F-C2F-C1F	2.27	104.39	100.98

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	B4P	C5E-O5E-PA-O3A
2	A	1	B4P	C5E-O5E-PA-O1A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	B4P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/245 (95%)	0.48	22 (9%) 8 10	38, 53, 71, 80	0
1	B	230/245 (93%)	0.73	40 (17%) 1 1	34, 53, 84, 103	0
All	All	463/490 (94%)	0.60	62 (13%) 3 4	34, 53, 79, 103	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	127	ILE	5.1
1	B	204	LEU	4.8
1	B	256	ASN	4.3
1	A	124[A]	TYR	3.9
1	B	276	THR	3.8
1	B	68	LEU	3.7
1	A	204	LEU	3.6
1	A	130	SER	3.4
1	A	271	VAL	3.4
1	B	185	ALA	3.3
1	A	205	LEU	3.2
1	B	205	LEU	3.2
1	A	185	ALA	3.1
1	B	69	LYS	3.1
1	A	243	VAL	3.1
1	B	255	ASP	3.0
1	B	186	ILE	3.0
1	B	184	ILE	3.0
1	A	277	ARG	3.0
1	B	282	PRO	2.9
1	B	227	THR	2.8
1	B	274	SER	2.8
1	B	287	ASN	2.8
1	A	275	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	87	LEU	2.6
1	B	226	ILE	2.6
1	B	263	ASP	2.6
1	A	274	SER	2.6
1	B	96	ASP	2.6
1	A	282	PRO	2.6
1	A	260	VAL	2.5
1	B	177	GLU	2.5
1	B	168	ALA	2.5
1	A	186	ILE	2.5
1	A	206	ILE	2.5
1	A	246	LEU	2.5
1	B	228	VAL	2.4
1	A	184	ILE	2.4
1	A	242	ILE	2.4
1	B	206	ILE	2.3
1	B	72	ASP	2.3
1	B	167	VAL	2.2
1	B	103	PRO	2.2
1	B	76	ASP	2.2
1	B	173	GLU	2.2
1	B	288	ASN	2.2
1	B	198	ILE	2.2
1	A	286	GLU	2.2
1	A	178	PHE	2.1
1	A	87	LEU	2.1
1	B	115	LEU	2.1
1	A	245	SER	2.1
1	B	197	LEU	2.1
1	B	128	TRP	2.1
1	B	77	LYS	2.1
1	B	159	LYS	2.1
1	B	172	ALA	2.1
1	A	256	ASN	2.1
1	B	124[A]	TYR	2.1
1	B	248	VAL	2.0
1	B	277	ARG	2.0
1	B	215	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	B4P	A	1	53/53	0.96	0.11	43,51,57,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.