



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 08:02 PM EDT

PDB ID : 3LQ1
Title : Crystal structure of 2-succinyl-6-hydroxy-2,4-cyclohexadiene 1-carboxylic acid synthase/2-oxoglutarate decarboxylase FROM *Listeria monocytogenes* str. 4b F2365
Authors : Patskovsky, Y.; Toro, R.; Freeman, J.; Hu, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York Structural GenomiX Research Consortium (NYS-GXRC); New York SGX Research Center for Structural Genomics (NYS-GXRC)
Deposited on : 2010-02-08
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	 4% 78% 10% • 11%
1	B	578	 4% 77% 11% • 10%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 8131 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-succinyl-5-enolpyruvyl-6-hydroxy-3-cyclohexene-1-carboxylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	513	4020	2581	665	752	22	0	0	0
1	B	518	4057	2605	670	760	22	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q71YZ2
A	0	SER	-	expression tag	UNP Q71YZ2
A	1	LEU	-	expression tag	UNP Q71YZ2
A	569	GLU	-	expression tag	UNP Q71YZ2
A	570	GLY	-	expression tag	UNP Q71YZ2
A	571	HIS	-	expression tag	UNP Q71YZ2
A	572	HIS	-	expression tag	UNP Q71YZ2
A	573	HIS	-	expression tag	UNP Q71YZ2
A	574	HIS	-	expression tag	UNP Q71YZ2
A	575	HIS	-	expression tag	UNP Q71YZ2
A	576	HIS	-	expression tag	UNP Q71YZ2
B	-1	MET	-	expression tag	UNP Q71YZ2
B	0	SER	-	expression tag	UNP Q71YZ2
B	1	LEU	-	expression tag	UNP Q71YZ2
B	569	GLU	-	expression tag	UNP Q71YZ2
B	570	GLY	-	expression tag	UNP Q71YZ2
B	571	HIS	-	expression tag	UNP Q71YZ2
B	572	HIS	-	expression tag	UNP Q71YZ2
B	573	HIS	-	expression tag	UNP Q71YZ2
B	574	HIS	-	expression tag	UNP Q71YZ2
B	575	HIS	-	expression tag	UNP Q71YZ2
B	576	HIS	-	expression tag	UNP Q71YZ2

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	32	Total 32	O 32	0	0
2	B	22	Total 22	O 22	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.69Å 99.69Å 259.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 39.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-2.60) 99.5 (39.74-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.222 , 0.265 0.222 , 0.267	Depositor DCC
R_{free} test set	1283 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å ²)	71.4	Xtrriage
Anisotropy	0.008	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8131	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/4106	0.55	0/5566
1	B	0.34	0/4144	0.53	0/5621
All	All	0.35	0/8250	0.54	0/11187

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4020	0	4030	20	0
1	B	4057	0	4074	25	0
2	A	32	0	0	0	0
2	B	22	0	0	0	0
All	All	8131	0	8104	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:VAL:HG12	1:B:314:ARG:HB2	1.83	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:ILE:HD11	1:B:276:PHE:HZ	1.70	0.57
1:B:230:PRO:HG2	1:B:296:SER:HB2	1.88	0.55
1:A:328:ILE:HG22	1:A:330:ALA:HB2	1.88	0.54
1:B:266:ASP:HB2	1:B:420:GLN:HE22	1.73	0.53
1:A:447:SER:HA	1:A:453:MET:HG3	1.90	0.52
1:B:141:GLU:HB2	1:B:146:MET:HG2	1.91	0.52
1:B:421:ILE:HG12	1:B:423:LYS:HE2	1.91	0.52
1:A:53:ASP:HB3	1:A:59:ALA:HB2	1.91	0.51
1:B:219:CYS:HB3	1:B:290:VAL:HG11	1.92	0.51
1:A:421:ILE:HG12	1:A:423:LYS:HE2	1.92	0.50
1:B:270:ILE:HD11	1:B:276:PHE:CZ	2.47	0.50
1:A:230:PRO:HG2	1:A:296:SER:HB2	1.94	0.49
1:A:389:ILE:HD13	1:A:484:VAL:HG22	1.94	0.49
1:B:227:VAL:HB	1:B:293:ARG:HG2	1.95	0.48
1:A:342:LEU:HD23	1:A:345:ILE:HD11	1.96	0.47
1:A:293:ARG:NH2	1:A:325:LYS:O	2.47	0.47
1:B:137:MET:CE	1:B:171:LEU:HD23	2.45	0.47
1:B:268:VAL:HG12	1:B:360:LEU:HD11	1.97	0.47
1:A:297:MET:HG3	1:A:327:PRO:HD2	1.96	0.47
1:B:378:MET:HG3	1:B:388:LYS:HG2	1.96	0.46
1:A:444:LEU:HD13	1:A:479:LEU:HD22	1.99	0.45
1:A:260:ARG:NH2	1:A:415:ASP:O	2.50	0.45
1:B:421:ILE:HG23	1:B:423:LYS:H	1.82	0.45
1:A:408:SER:HB2	1:A:409:MET:H	1.53	0.44
1:A:237:GLU:H	1:A:237:GLU:HG2	1.52	0.43
1:A:40:LEU:HB3	1:A:183:LEU:HD21	2.00	0.43
1:B:396:LEU:HB3	1:B:540:ILE:HG21	2.00	0.43
1:B:447:SER:HA	1:B:453:MET:HG3	1.99	0.43
1:A:431:ARG:CG	1:A:431:ARG:HH11	2.31	0.42
1:B:119:GLN:HE21	1:B:119:GLN:HB3	1.66	0.42
1:B:229:GLY:HA2	1:B:254:ASP:HB2	2.01	0.42
1:A:253:ALA:O	1:A:260:ARG:NH1	2.43	0.42
1:B:270:ILE:HD12	1:B:367:ASN:HB2	2.02	0.42
1:B:469:LEU:HD22	1:B:549:LEU:HD21	2.02	0.42
1:A:6:GLN:HG2	1:A:182:ILE:HD12	2.02	0.42
1:B:9:THR:HA	1:B:41:MET:HE1	2.02	0.41
1:B:237:GLU:H	1:B:237:GLU:HG2	1.60	0.41
1:A:141:GLU:HB2	1:A:146:MET:HG2	2.02	0.41
1:A:396:LEU:HD11	1:A:537:GLU:HG2	2.03	0.41
1:B:60:GLY:HA2	1:B:76:LEU:HD11	2.03	0.41
1:B:405:ILE:HD13	1:B:414:VAL:HG11	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:LEU:HA	1:A:347:GLN:HE21	1.85	0.40
1:B:253:ALA:O	1:B:260:ARG:NH1	2.53	0.40
1:B:137:MET:HE2	1:B:171:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	505/578 (87%)	489 (97%)	15 (3%)	1 (0%)	47 71
1	B	512/578 (89%)	498 (97%)	13 (2%)	1 (0%)	47 71
All	All	1017/1156 (88%)	987 (97%)	28 (3%)	2 (0%)	47 71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	408	SER
1	A	408	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	434/489 (89%)	401 (92%)	33 (8%)	13 26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	439/489 (90%)	399 (91%)	40 (9%)	9	18
All	All	873/978 (89%)	800 (92%)	73 (8%)	11	21

All (73) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ARG
1	A	88	PHE
1	A	103	VAL
1	A	119	GLN
1	A	126	LEU
1	A	144	GLU
1	A	145	GLU
1	A	148	ARG
1	A	164	THR
1	A	183	LEU
1	A	218	GLU
1	A	237	GLU
1	A	240	MET
1	A	260	ARG
1	A	265	LEU
1	A	267	GLU
1	A	309	GLN
1	A	310	LEU
1	A	312	ASP
1	A	313	ILE
1	A	314	ARG
1	A	317	VAL
1	A	329	LYS
1	A	342	LEU
1	A	352	ASP
1	A	377	GLU
1	A	408	SER
1	A	424	LYS
1	A	426	LYS
1	A	431	ARG
1	A	444	LEU
1	A	524	ASP
1	A	551	ILE
1	B	4	HIS
1	B	34	ARG
1	B	36	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	49	LYS
1	B	54	VAL
1	B	88	PHE
1	B	98	GLN
1	B	103	VAL
1	B	126	LEU
1	B	145	GLU
1	B	148	ARG
1	B	213	GLN
1	B	236	LEU
1	B	237	GLU
1	B	260	ARG
1	B	266	ASP
1	B	270	ILE
1	B	290	VAL
1	B	309	GLN
1	B	310	LEU
1	B	313	ILE
1	B	317	VAL
1	B	329	LYS
1	B	332	THR
1	B	342	LEU
1	B	377	GLU
1	B	378	MET
1	B	380	ASN
1	B	396	LEU
1	B	411	ILE
1	B	414	VAL
1	B	421	ILE
1	B	424	LYS
1	B	426	LYS
1	B	444	LEU
1	B	467	ASN
1	B	479	LEU
1	B	486	ASN
1	B	546	HIS
1	B	554	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	309	GLN
1	A	347	GLN
1	B	98	GLN
1	B	119	GLN
1	B	309	GLN
1	B	347	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	513/578 (88%)	0.22	25 (4%) 29 23	55, 80, 119, 151	0
1	B	518/578 (89%)	0.13	21 (4%) 37 30	48, 85, 126, 157	0
All	All	1031/1156 (89%)	0.17	46 (4%) 33 26	48, 82, 123, 157	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	353	ASP	4.8
1	A	2	THR	4.5
1	A	379	ALA	4.2
1	B	382	THR	4.1
1	B	399	ASP	4.1
1	A	352	ASP	3.8
1	B	383	ILE	3.7
1	B	380	ASN	3.7
1	A	183	LEU	3.7
1	B	184	GLU	3.6
1	A	202	TYR	3.5
1	A	40	LEU	3.3
1	B	363	TRP	3.2
1	B	201	TYR	3.2
1	B	381	THR	3.1
1	B	183	LEU	3.0
1	A	378	MET	3.0
1	A	380	ASN	3.0
1	A	37	PRO	3.0
1	A	531	LYS	2.9
1	A	486	ASN	2.8
1	A	184	GLU	2.7
1	A	349	ASN	2.7
1	A	351	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	220	THR	2.6
1	A	182	ILE	2.5
1	B	384	LEU	2.5
1	B	220	THR	2.5
1	A	545	TYR	2.5
1	B	179	LEU	2.5
1	B	400	LYS	2.4
1	A	532	SER	2.4
1	A	38	LEU	2.4
1	A	348	GLN	2.3
1	B	353	ASP	2.3
1	A	188	PHE	2.3
1	A	179	LEU	2.3
1	B	202	TYR	2.3
1	B	486	ASN	2.3
1	B	46	PRO	2.2
1	B	217	THR	2.2
1	A	218	GLU	2.2
1	B	47	ILE	2.2
1	B	189	THR	2.2
1	A	180	VAL	2.1
1	B	286	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.