



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 10:19 PM EDT

PDB ID : 3MAE
Title : CRYSTAL STRUCTURE OF PROBABLE DIHYDROLIPOAMIDE ACETYLTRANSFERASE FROM LISTERIA MONOCYTOGENES 4b F2365
Authors : Patskovsky, Y.; Toro, R.; Gilmore, M.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-03-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	 10% 84% 7% 9%
1	B	256	 18% 88% 5% • 6%
1	C	256	 16% 86% 7% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	429	-	-	-	X
4	GOL	C	425	-	-	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5666 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-oxoisovalerate dehydrogenase E2 component, dihydrolipoamide acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	Total 1801	C 1147	N 311	O 335	S 8	0	2	0
1	B	240	Total 1843	C 1172	N 319	O 344	S 8	0	2	0
1	C	239	Total 1842	C 1173	N 320	O 341	S 8	0	2	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	169	MET	-	expression tag	UNP Q71ZU8
A	170	SER	-	expression tag	UNP Q71ZU8
A	171	LEU	-	expression tag	UNP Q71ZU8
A	417	GLU	-	expression tag	UNP Q71ZU8
A	418	GLY	-	expression tag	UNP Q71ZU8
A	419	HIS	-	expression tag	UNP Q71ZU8
A	420	HIS	-	expression tag	UNP Q71ZU8
A	421	HIS	-	expression tag	UNP Q71ZU8
A	422	HIS	-	expression tag	UNP Q71ZU8
A	423	HIS	-	expression tag	UNP Q71ZU8
A	424	HIS	-	expression tag	UNP Q71ZU8
B	169	MET	-	expression tag	UNP Q71ZU8
B	170	SER	-	expression tag	UNP Q71ZU8
B	171	LEU	-	expression tag	UNP Q71ZU8
B	417	GLU	-	expression tag	UNP Q71ZU8
B	418	GLY	-	expression tag	UNP Q71ZU8
B	419	HIS	-	expression tag	UNP Q71ZU8
B	420	HIS	-	expression tag	UNP Q71ZU8
B	421	HIS	-	expression tag	UNP Q71ZU8
B	422	HIS	-	expression tag	UNP Q71ZU8
B	423	HIS	-	expression tag	UNP Q71ZU8
B	424	HIS	-	expression tag	UNP Q71ZU8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	169	MET	-	expression tag	UNP Q71ZU8
C	170	SER	-	expression tag	UNP Q71ZU8
C	171	LEU	-	expression tag	UNP Q71ZU8
C	417	GLU	-	expression tag	UNP Q71ZU8
C	418	GLY	-	expression tag	UNP Q71ZU8
C	419	HIS	-	expression tag	UNP Q71ZU8
C	420	HIS	-	expression tag	UNP Q71ZU8
C	421	HIS	-	expression tag	UNP Q71ZU8
C	422	HIS	-	expression tag	UNP Q71ZU8
C	423	HIS	-	expression tag	UNP Q71ZU8
C	424	HIS	-	expression tag	UNP Q71ZU8

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Cl 3 3	0	0
2	B	4	Total Cl 4 4	0	0
2	C	2	Total Cl 2 2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	24	Total O 24 24	0	0
5	B	21	Total O 21 21	0	0
5	C	23	Total O 23 23	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-oxoisovalerate dehydrogenase E2 component, dihydrolipoamide acetyltransferase



- Molecule 1: 2-oxoisovalerate dehydrogenase E2 component, dihydrolipoamide acetyltransferase



- Molecule 1: 2-oxoisovalerate dehydrogenase E2 component, dihydrolipoamide acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	110.22Å 173.01Å 131.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 39.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-2.50) 99.4 (39.54-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.192 , 0.234 0.193 , 0.232	Depositor DCC
R_{free} test set	1342 reflections (3.10%)	wwPDB-VP
Wilson B-factor (Å ²)	43.9	Xtrriage
Anisotropy	0.321	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5666	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1834	0.53	1/2471 (0.0%)
1	B	0.37	0/1878	0.51	0/2534
1	C	0.37	0/1877	0.54	1/2530 (0.0%)
All	All	0.37	0/5589	0.53	2/7535 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	223	LEU	CA-CB-CG	5.88	128.83	115.30
1	C	223	LEU	CA-CB-CG	5.04	126.88	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1801	0	1863	9	0
1	B	1843	0	1903	10	0
1	C	1842	0	1908	10	0
2	A	3	0	0	0	0
2	B	4	0	0	0	0
2	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	20	0	0	0	0
3	B	10	0	0	0	0
3	C	25	0	0	0	0
4	A	30	0	40	2	0
4	B	6	0	8	0	0
4	C	12	0	16	0	0
5	A	24	0	0	0	0
5	B	21	0	0	0	0
5	C	23	0	0	0	0
All	All	5666	0	5738	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:374:VAL:HG13	1:C:367:ILE:HD13	1.85	0.58
1:A:343:GLN:HE22	1:B:343:GLN:NE2	2.02	0.58
1:A:367:ILE:HD13	1:C:374:VAL:HG13	1.87	0.56
1:A:295:LYS:NZ	1:A:324:ALA:O	2.39	0.56
1:B:269:ASP:HB3	1:B:270:LYS:HD3	1.87	0.55
1:C:235:LYS:HE2	1:C:241:SER:HB3	1.93	0.50
1:B:223:LEU:HD23	1:B:223:LEU:O	2.11	0.49
1:A:271:ILE:HG13	1:C:191:ILE:HD11	1.94	0.48
1:B:223:LEU:HD23	1:B:223:LEU:C	2.35	0.47
1:B:215:MET:HG3	1:B:381:CYS:SG	2.55	0.46
1:B:410:LYS:HD3	1:B:410:LYS:H	1.81	0.45
1:A:213:TRP:HE1	1:A:381:CYS:HB3	1.81	0.45
1:A:345:MET:HE2	1:A:345:MET:HB2	1.68	0.44
1:A:373:ALA:HB1	4:A:431:GOL:H12	2.00	0.43
4:A:425:GOL:H32	1:C:182:SER:HB3	2.00	0.43
1:C:288:LEU:HB2	1:C:289:LEU:H	1.71	0.42
1:C:215:MET:HG3	1:C:381:CYS:SG	2.59	0.42
1:B:191:ILE:HD11	1:C:271:ILE:HG13	2.00	0.42
1:A:343:GLN:HE22	1:B:343:GLN:HE22	1.68	0.41
1:A:365:PRO:HD3	1:B:367:ILE:HG12	2.03	0.41
1:C:261[B]:GLN:HG3	1:C:393:LEU:HD13	2.03	0.41
1:C:195:ARG:HG2	1:C:288:LEU:HD23	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/256 (91%)	226 (97%)	8 (3%)	0	100	100
1	B	240/256 (94%)	234 (98%)	6 (2%)	0	100	100
1	C	239/256 (93%)	233 (98%)	6 (2%)	0	100	100
All	All	713/768 (93%)	693 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/209 (92%)	184 (96%)	8 (4%)	30	54
1	B	198/209 (95%)	191 (96%)	7 (4%)	36	62
1	C	197/209 (94%)	190 (96%)	7 (4%)	35	61
All	All	587/627 (94%)	565 (96%)	22 (4%)	33	60

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	LYS
1	A	214	MET
1	A	218	VAL
1	A	223	LEU
1	A	230	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	357	GLN
1	A	366	VAL
1	A	399	GLN
1	B	214	MET
1	B	236	LYS
1	B	270	LYS
1	B	288	LEU
1	B	291	VAL
1	B	410	LYS
1	B	416	TYR
1	C	195	ARG
1	C	216	VAL
1	C	262	LEU
1	C	288	LEU
1	C	317	ARG
1	C	320	LYS
1	C	416	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	263	ASN
1	A	343	GLN
1	A	352	GLN
1	B	343	GLN
1	B	352	GLN
1	C	323	GLN
1	C	343	GLN
1	C	352	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 9 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	427	-	5,5,5	0.34	0	5,5,5	0.39	0
3	PO4	A	10	-	4,4,4	0.96	0	6,6,6	0.50	0
4	GOL	A	431	-	5,5,5	0.36	0	5,5,5	0.36	0
3	PO4	C	5	-	4,4,4	0.93	0	6,6,6	0.46	0
3	PO4	A	11	-	4,4,4	0.98	0	6,6,6	0.49	0
3	PO4	B	3	-	4,4,4	0.98	0	6,6,6	0.48	0
4	GOL	A	426	-	5,5,5	0.37	0	5,5,5	0.16	0
4	GOL	C	425	-	5,5,5	0.38	0	5,5,5	0.32	0
4	GOL	A	425	-	5,5,5	0.39	0	5,5,5	0.39	0
3	PO4	C	4	-	4,4,4	0.98	0	6,6,6	0.42	0
3	PO4	A	2	-	4,4,4	0.96	0	6,6,6	0.51	0
4	GOL	B	425	-	5,5,5	0.37	0	5,5,5	0.31	0
3	PO4	C	6	-	4,4,4	0.92	0	6,6,6	0.50	0
3	PO4	A	9	-	4,4,4	0.91	0	6,6,6	0.50	0
3	PO4	B	12	-	4,4,4	0.96	0	6,6,6	0.45	0
4	GOL	A	429	-	5,5,5	0.40	0	5,5,5	0.17	0
3	PO4	C	7	-	4,4,4	0.95	0	6,6,6	0.43	0
4	GOL	C	426	-	5,5,5	0.37	0	5,5,5	0.32	0
3	PO4	C	8	-	4,4,4	0.89	0	6,6,6	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	427	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	429	-	-	0/4/4/4	-
4	GOL	A	426	-	-	2/4/4/4	-
4	GOL	B	425	-	-	3/4/4/4	-
4	GOL	A	431	-	-	2/4/4/4	-
4	GOL	C	425	-	-	2/4/4/4	-
4	GOL	C	426	-	-	4/4/4/4	-
4	GOL	A	425	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	427	GOL	O1-C1-C2-O2
4	A	427	GOL	O1-C1-C2-C3
4	A	431	GOL	O1-C1-C2-O2
4	A	431	GOL	O1-C1-C2-C3
4	B	425	GOL	C1-C2-C3-O3
4	B	425	GOL	O2-C2-C3-O3
4	C	425	GOL	O1-C1-C2-C3
4	C	426	GOL	O1-C1-C2-O2
4	A	427	GOL	C1-C2-C3-O3
4	B	425	GOL	O1-C1-C2-C3
4	C	426	GOL	O1-C1-C2-C3
4	C	426	GOL	C1-C2-C3-O3
4	C	425	GOL	O1-C1-C2-O2
4	A	427	GOL	O2-C2-C3-O3
4	C	426	GOL	O2-C2-C3-O3
4	A	426	GOL	O1-C1-C2-O2
4	A	426	GOL	O1-C1-C2-C3
4	A	425	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	431	GOL	1	0
4	A	425	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/256 (91%)	0.49	25 (10%) 6 5	27, 43, 89, 109	0
1	B	240/256 (93%)	0.89	46 (19%) 1 1	29, 51, 102, 109	0
1	C	239/256 (93%)	0.85	42 (17%) 1 1	29, 44, 96, 110	0
All	All	713/768 (92%)	0.75	113 (15%) 2 1	27, 46, 98, 110	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	287	ASP	9.4
1	B	288	LEU	8.6
1	C	288	LEU	7.3
1	C	416	TYR	7.3
1	B	416	TYR	7.1
1	A	414	ALA	6.8
1	C	409	SER	6.8
1	A	416	TYR	6.5
1	B	287	ASP	6.3
1	C	410	LYS	6.0
1	C	413	THR	5.9
1	C	414	ALA	5.5
1	B	236	LYS	5.4
1	C	415	LEU	5.3
1	A	287	ASP	5.3
1	B	322	SER	5.3
1	A	236	LYS	5.2
1	A	412	ASN	5.2
1	C	412	ASN	5.1
1	C	286	GLY	5.1
1	C	236	LYS	5.1
1	B	323	GLN	5.0
1	B	177	THR	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	288	LEU	4.9
1	A	410	LYS	4.9
1	B	412	ASN	4.5
1	B	316	ALA	4.3
1	B	413	THR	4.3
1	B	317	ARG	4.3
1	B	303	LYS	4.2
1	B	240	TYR	4.2
1	B	320	LYS	4.1
1	A	413	THR	4.1
1	C	191	ILE	4.0
1	B	345	MET	4.0
1	A	183	ALA	4.0
1	A	285	ALA	4.0
1	A	415	LEU	4.0
1	B	411	GLU	4.0
1	A	191	ILE	4.0
1	B	284	ALA	4.0
1	B	290	TYR	3.9
1	B	324	ALA	3.9
1	C	233	SER	3.8
1	C	411	GLU	3.8
1	B	192	ASN	3.8
1	A	232	ASP	3.7
1	C	194	VAL	3.7
1	B	237	GLU	3.6
1	C	240	TYR	3.6
1	A	237	GLU	3.6
1	B	286	GLY	3.6
1	B	314	GLY	3.6
1	B	313	ALA	3.6
1	B	414	ALA	3.5
1	A	269	ASP	3.5
1	C	234	PHE	3.4
1	C	232	ASP	3.4
1	A	289	LEU	3.4
1	C	317	ARG	3.4
1	B	193	GLY	3.3
1	B	415	LEU	3.3
1	B	234	PHE	3.2
1	C	193	GLY	3.2
1	A	411	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	237	GLU	3.1
1	C	289	LEU	3.1
1	C	241	SER	3.1
1	B	319	GLY	3.1
1	C	318	ASN	3.1
1	B	321	LEU	3.1
1	B	318	ASN	3.0
1	C	316	ALA	3.0
1	B	194	VAL	2.9
1	C	213	TRP	2.9
1	B	215	MET	2.9
1	A	345	MET	2.8
1	B	344[A]	SER	2.8
1	C	242	LEU	2.8
1	B	409	SER	2.8
1	C	345	MET	2.7
1	A	409	SER	2.7
1	C	226	TYR	2.7
1	B	178	PRO	2.7
1	B	312	LEU	2.6
1	A	286	GLY	2.6
1	B	410	LYS	2.6
1	B	381	CYS	2.6
1	B	285	ALA	2.6
1	A	290	TYR	2.6
1	A	284	ALA	2.5
1	B	213	TRP	2.5
1	C	215	MET	2.5
1	C	192	ASN	2.5
1	C	231	LYS	2.4
1	B	325	ASP	2.4
1	C	290	TYR	2.4
1	C	319	GLY	2.4
1	B	191	ILE	2.4
1	C	383	SER	2.4
1	A	194	VAL	2.4
1	C	239	GLY	2.3
1	C	284	ALA	2.3
1	B	307	ARG	2.2
1	B	269	ASP	2.2
1	C	214	MET	2.2
1	C	235	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	355	ILE	2.2
1	A	215	MET	2.1
1	C	355	ILE	2.1
1	A	213	TRP	2.1
1	C	384	ILE	2.0
1	C	382	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	429	6/6	0.73	0.47	89,91,95,97	0
4	GOL	C	425	6/6	0.74	0.40	71,76,78,80	0
3	PO4	C	7	5/5	0.79	0.27	57,60,66,68	5
3	PO4	C	8	5/5	0.83	0.31	38,40,51,52	5
4	GOL	A	427	6/6	0.84	0.29	73,78,79,80	0
3	PO4	B	3	5/5	0.85	0.25	59,60,66,68	5
4	GOL	A	425	6/6	0.87	0.25	71,77,79,83	0
4	GOL	C	426	6/6	0.87	0.20	68,76,83,87	0
4	GOL	A	431	6/6	0.88	0.19	61,66,70,76	0
3	PO4	B	12	5/5	0.88	0.36	60,64,67,68	5
3	PO4	A	2	5/5	0.88	0.28	52,61,63,65	5
3	PO4	A	10	5/5	0.89	0.14	53,56,61,62	5
3	PO4	C	4	5/5	0.89	0.15	59,60,64,65	5
4	GOL	A	426	6/6	0.91	0.24	71,75,77,83	0
4	GOL	B	425	6/6	0.91	0.40	67,75,77,77	0
3	PO4	A	11	5/5	0.92	0.32	53,58,63,67	5
3	PO4	C	5	5/5	0.92	0.15	94,95,96,99	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	A	428	1/1	0.93	0.37	69,69,69,69	0
2	CL	B	426	1/1	0.94	0.19	70,70,70,70	0
3	PO4	C	6	5/5	0.94	0.36	38,44,52,54	5
2	CL	B	427	1/1	0.94	0.59	74,74,74,74	0
2	CL	A	430	1/1	0.95	0.13	66,66,66,66	0
2	CL	B	428	1/1	0.95	0.20	68,68,68,68	0
3	PO4	A	9	5/5	0.97	0.15	87,87,87,88	0
2	CL	C	427	1/1	0.97	0.41	69,69,69,69	0
2	CL	B	1	1/1	0.97	0.19	56,56,56,56	0
2	CL	A	1	1/1	0.98	0.22	54,54,54,54	0
2	CL	C	1	1/1	0.98	0.20	53,53,53,53	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.