

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 17, 2024 – 11:57 AM EDT

PDB ID : 3N07

Title: Structure of putative 3-deoxy-D-manno-octulosonate 8-phosphate phosphatese

from Vibrio cholerae

Authors: Liu, W.; Ramagopal, U.A.; Toro, R.; Burley, S.K.; Almo, S.C.; New York SGX

Research Center for Structural Genomics (NYSGXRC)

Deposited on : 2010-05-13

Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 2.37.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	195	% 	7% • 8%
1	В	195	88%	•• 10%
1	С	195	6% 79%	10% • 10%
1	D	195	81%	9% • 9%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 3-deoxy-D-manno-octulosonate 8-phosphate phosphatase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Λ	179	Total	С	N	О	S	Se	0	5	0
1	Λ	119	1421	896	243	271	5	6	0	3	
1	В	176	Total	С	N	О	S	Se	0	4	0
1	Ъ	170	1389	875	239	266	4	5	0	4	0
1	С	176	Total	С	N	О	S	Se	0	4	0
1		170	1390	879	238	263	4	6	0	4	0
1	D	177	Total	С	N	О	S	Se	0	6	0
1	ש	111	1415	892	243	270	5	5		0	U

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP Q9KP52
A	0	SER	-	expression tag	UNP Q9KP52
A	1	LEU	-	expression tag	UNP Q9KP52
A	186	GLU	-	expression tag	UNP Q9KP52
A	187	GLY	-	expression tag	UNP Q9KP52
A	188	HIS	-	expression tag	UNP Q9KP52
A	189	HIS	-	expression tag	UNP Q9KP52
A	190	HIS	-	expression tag	UNP Q9KP52
A	191	HIS	-	expression tag	UNP Q9KP52
A	192	HIS	-	expression tag	UNP Q9KP52
A	193	HIS	-	expression tag	UNP Q9KP52
В	-1	MSE	-	expression tag	UNP Q9KP52
В	0	SER	-	expression tag	UNP Q9KP52
В	1	LEU	-	expression tag	UNP Q9KP52
В	186	GLU	-	expression tag	UNP Q9KP52
В	187	GLY	-	expression tag	UNP Q9KP52
В	188	HIS	-	expression tag	UNP Q9KP52
В	189	HIS	-	expression tag	UNP Q9KP52
В	190	HIS	-	expression tag	UNP Q9KP52
В	191	HIS	-	expression tag	UNP Q9KP52
В	192	HIS	-	expression tag	UNP Q9KP52



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
В	193	HIS	-	expression tag	UNP Q9KP52
С	-1	MSE	-	expression tag	UNP Q9KP52
С	0	SER	-	expression tag	UNP Q9KP52
С	1	LEU	-	expression tag	UNP Q9KP52
С	186	GLU	-	expression tag	UNP Q9KP52
С	187	GLY	-	expression tag	UNP Q9KP52
С	188	HIS	-	expression tag	UNP Q9KP52
С	189	HIS	-	expression tag	UNP Q9KP52
С	190	HIS	-	expression tag	UNP Q9KP52
С	191	HIS	-	expression tag	UNP Q9KP52
С	192	HIS	-	expression tag	UNP Q9KP52
С	193	HIS	-	expression tag	UNP Q9KP52
D	-1	MSE	-	expression tag	UNP Q9KP52
D	0	SER	-	expression tag	UNP Q9KP52
D	1	LEU	-	expression tag	UNP Q9KP52
D	186	GLU	-	expression tag	UNP Q9KP52
D	187	GLY	-	expression tag	UNP Q9KP52
D	188	HIS	-	expression tag	UNP Q9KP52
D	189	HIS	-	expression tag	UNP Q9KP52
D	190	HIS		expression tag	UNP Q9KP52
D	191	HIS	-	expression tag	UNP Q9KP52
D	192	HIS	-	expression tag	UNP Q9KP52
D	193	HIS	-	expression tag	UNP Q9KP52

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	В	1	Total Mg 1 1	0	0
2	С	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

• Molecule 3 is water.

-	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	3	A	98	Total O 98 98	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	97	Total O 97 97	0	0
3	С	52	Total O 52 52	0	0
3	D	68	Total O 68 68	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-deoxy-D-manno-octulosonate 8-phosphate phosphatase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	63.95Å 77.11Å 78.50Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 106.65° 90.00°	Depositor
Resolution (Å)	50.00 - 1.76	Depositor
Resolution (A)	33.46 - 1.76	EDS
% Data completeness	99.2 (50.00-1.76)	Depositor
(in resolution range)	99.2 (33.46-1.76)	EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.57 (at 1.76Å)	Xtriage
Refinement program	REFMAC	Depositor
$R, R_{free}$	0.170 , 0.201	Depositor
It, It free	0.185 , $0.214$	DCC
$R_{free}$ test set	3634  reflections  (5.04%)	wwPDB-VP
Wilson B-factor $(\mathring{A}^2)$	20.3	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 45.8	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5934	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
Mol	Chain	RMSZ  #  Z  > 5		RMSZ	# Z  > 5	
1	A	0.63	0/1437	0.69	1/1935 (0.1%)	
1	В	0.69	0/1404	0.71	0/1890	
1	С	0.57	0/1405	0.64	0/1890	
1	D	0.64	0/1430	0.70	0/1924	
All	All	0.63	0/5676	0.68	1/7639 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

N	Лol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
	1	A	141	ASP	CB-CG-OD1	5.51	123.26	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1421	0	1427	6	0
1	В	1389	0	1398	3	0
1	С	1390	0	1410	14	0
1	D	1415	0	1426	12	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	98	0	0	0	0
3	В	97	0	0	0	0
3	С	52	0	0	1	0
3	D	68	0	0	1	0
All	All	5934	0	5661	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
1 A 00 ACD OD0	1 A 101 CLNI II CO	distance (Å)	overlap (Å)
1:A:98:ASP:OD2	1:A:101:GLN:HG2	1.87	0.73
1:C:17:LEU:O	1:C:21:LYS:HG3	1.95	0.65
1:A:175:ASN:HD21	1:B:136:ARG:NH1	1.98	0.61
1:C:109[B]:LYS:O	1:C:109[B]:LYS:HG3	2.03	0.58
1:D:62:LEU:HD23	1:D:69:ILE:HD11	1.85	0.58
1:D:97:ASP:N	1:D:97:ASP:OD1	2.36	0.57
1:A:177[A]:LEU:C	1:A:177[A]:LEU:HD23	2.26	0.55
1:C:91:LEU:HD21	1:C:109[B]:LYS:HD3	1.86	0.55
1:C:59:VAL:HG12	1:C:63:MSE:HE2	1.89	0.55
1:C:22:GLN:O	1:C:22:GLN:HG2	2.07	0.54
1:D:124:LEU:HD21	1:D:145:LEU:HB3	1.89	0.53
1:D:125:ILE:HG13	3:D:315:HOH:O	2.11	0.51
1:D:130[A]:MSE:SE	1:D:146:LEU:HD11	2.60	0.51
1:D:29[B]:ASP:HB3	1:D:33:VAL:HG21	1.94	0.50
1:C:39:ILE:HG22	1:C:41[B]:MSE:SE	2.62	0.49
1:D:59:VAL:HG12	1:D:63:MSE:HE2	1.95	0.48
1:D:59:VAL:HG11	1:D:89:ILE:HD11	1.95	0.47
1:D:27:ILE:O	1:D:119:TYR:HA	2.15	0.47
1:C:113:ALA:HB3	1:C:116:GLN:OE1	2.14	0.46
1:A:39:ILE:HG22	1:A:41[B]:MSE:SE	2.66	0.46
1:B:59:VAL:HG11	1:B:89:ILE:HD11	1.97	0.46
1:C:63:MSE:HE1	1:C:87:LEU:O	2.15	0.45
1:C:130[B]:MSE:CG	1:C:136:ARG:HB3	2.46	0.45
1:D:81:GLU:HG2	1:D:94:GLN:NE2	2.32	0.45
1:B:98:ASP:CG	1:B:98:ASP:O	2.55	0.44
1:C:130[A]:MSE:SE	1:C:146:LEU:HD11	2.68	0.44
1:C:27:ILE:O	1:C:119:TYR:HA	2.18	0.43



$\alpha \cdots \alpha$		
Continued fr	om $previous$	paae

Atom-1	Atom-2	Interatomic	Clash
7100111 1	1100111 2	distance (Å)	overlap (Å)
1:D:130[B]:MSE:CG	1:D:136:ARG:HB3	2.48	0.43
1:A:40:TYR:HB2	1:A:48:LEU:HB2	2.01	0.43
1:C:21:LYS:HG2	1:C:173:ALA:O	2.18	0.43
1:C:41[B]:MSE:SE	3:C:210:HOH:O	2.87	0.42
1:A:27:ILE:O	1:A:119:TYR:HA	2.20	0.42
1:D:40:TYR:O	1:D:47:GLU:HA	2.21	0.41
1:C:24:LYS:HD3	1:C:24:LYS:HA	1.82	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	182/195 (93%)	178 (98%)	4 (2%)	0	100	100
1	В	178/195 (91%)	176 (99%)	2 (1%)	0	100	100
1	С	178/195 (91%)	175 (98%)	3 (2%)	0	100	100
1	D	181/195 (93%)	177 (98%)	4 (2%)	0	100	100
All	All	719/780 (92%)	706 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	153/157 (98%)	148 (97%)	5 (3%)	38 15
1	В	149/157 (95%)	147 (99%)	2 (1%)	69 54
1	С	149/157~(95%)	146 (98%)	3 (2%)	55 34
1	D	152/157 (97%)	150 (99%)	2 (1%)	69 54
All	All	$603/628 \ (96\%)$	591 (98%)	12 (2%)	53 34

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	LYS
1	A	62	LEU
1	A	83	ARG
1	A	98	ASP
1	A	108	GLN
1	В	98	ASP
1	В	144	PRO
1	С	3	SER
1	С	22	GLN
1	С	85	LYS
1	D	125	ILE
1	D	145	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	44	GLN
1	D	148	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



### 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	175/195~(89%)	-0.08	1 (0%) 89 92	17, 27, 43, 54	0
1	В	172/195~(88%)	-0.18	4 (2%) 60 67	17, 24, 36, 48	0
1	С	172/195 (88%)	0.33	11 (6%) 19 25	20, 33, 53, 74	0
1	D	173/195 (88%)	-0.11	4 (2%) 60 67	18, 28, 46, 55	0
All	All	692/780 (88%)	-0.01	20 (2%) 51 57	17, 28, 47, 74	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	2	SER	10.2
1	С	3	SER	4.9
1	В	177	LEU	4.4
1	С	97	ASP	4.2
1	В	98	ASP	3.6
1	D	108	GLN	3.5
1	С	66	GLY	3.0
1	В	97	ASP	2.8
1	С	64	ASN	2.8
1	D	110	LEU	2.8
1	С	98	ASP	2.7
1	D	3	SER	2.7
1	С	18	GLU	2.5
1	С	5	VAL	2.5
1	С	65	ALA	2.4
1	С	14	PRO	2.4
1	В	101	GLN	2.4
1	A	3	SER	2.3
1	D	97	ASP	2.2
1	С	177	LEU	2.1



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	MG	D	200	1/1	0.79	0.20	14,14,14,14	1
2	MG	A	200	1/1	0.98	0.04	20,20,20,20	0
2	MG	С	200	1/1	0.99	0.08	21,21,21,21	0
2	MG	В	200	1/1	1.00	0.03	16,16,16,16	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

