



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2024 – 11:20 AM EDT

PDB ID : 3PRK  
Title : INHIBITION OF PROTEINASE K BY METHOXYSUCCINYL-ALA-AL  
A-PRO-ALA-CHLOROMETHYL KETONE. AN X-RAY STUDY AT 2.2-  
ANGSTROMS RESOLUTION  
Authors : Wolf, W.M.; Bajorath, J.; Mueller, A.; Raghunathan, S.; Singh, T.P.; Hinrichs,  
W.; Saenger, W.  
Deposited on : 1991-08-07  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	279	 71% 23% 6% •
2	I	6	 67% 33%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEINASE K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	279	2022	1243	355	414	10	0	0	0

- Molecule 2 is a protein called METHOXYSUCCINYL-ALA-ALA-PRO-ALA-CHLOROMETHYL KETONE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	I	6	31	20	4	7	0	0	1

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	E	1	1	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	E	164	164	164	0	0
4	I	6	6	6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEINASE K



- Molecule 2: METHOXYSUCCINYL-ALA-ALA-PRO-ALA-CHLOROMETHYL KETONE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.30Å 68.30Å 108.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.20 57.79 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.20) 88.2 (57.79-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.09 (at 2.19Å)	Xtrriage
Refinement program	TNT	Depositor
R, $R_{free}$	0.198 , (Not available) 0.192 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.2	Xtrriage
Anisotropy	0.269	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 71.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2224	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0QE, MSU, ALV, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	E	0.74	5/2061 (0.2%)	1.16	20/2800 (0.7%)
2	I	0.42	0/17	1.35	0/23
All	All	0.73	5/2078 (0.2%)	1.16	20/2823 (0.7%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	27	GLU	CD-OE2	6.57	1.32	1.25
1	E	174	GLU	CD-OE1	6.36	1.32	1.25
1	E	50	GLU	CD-OE2	5.62	1.31	1.25
1	E	48	GLU	CD-OE2	5.36	1.31	1.25
1	E	43	GLU	CD-OE1	5.15	1.31	1.25

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	147	ARG	NE-CZ-NH2	-10.05	115.27	120.30
1	E	200	ASP	CB-CG-OD2	8.42	125.88	118.30
1	E	200	ASP	CB-CG-OD1	-8.38	110.75	118.30
1	E	184	ASP	CB-CG-OD1	7.31	124.88	118.30
1	E	184	ASP	CB-CG-OD2	-7.17	111.85	118.30
1	E	39	ASP	CB-CG-OD2	6.77	124.40	118.30
1	E	39	ASP	CB-CG-OD1	-6.53	112.42	118.30
1	E	26	ASP	CB-CG-OD1	-6.13	112.78	118.30
1	E	187	ASP	CB-CG-OD1	-6.03	112.87	118.30
1	E	26	ASP	CB-CG-OD2	5.92	123.63	118.30
1	E	112	ASP	CB-CG-OD1	-5.91	112.98	118.30
1	E	112	ASP	CB-CG-OD2	5.86	123.57	118.30
1	E	147	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	E	187	ASP	CB-CG-OD2	5.60	123.34	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	117	ASP	CB-CG-OD2	-5.48	113.37	118.30
1	E	185	ARG	NE-CZ-NH1	5.39	122.99	120.30
1	E	106	THR	N-CA-CB	5.30	120.36	110.30
1	E	98	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	E	189	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	E	58	THR	N-CA-CB	5.08	119.94	110.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2022	0	1922	66	0
2	I	31	0	27	4	0
3	E	1	0	0	0	0
4	E	164	0	0	4	0
4	I	6	0	0	0	0
All	All	2224	0	1949	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:185:ARG:HH11	1:E:185:ARG:HG2	1.29	0.97
1:E:261:LEU:H	1:E:270:ASN:HD21	1.08	0.94
1:E:58:THR:HG21	1:E:62:SER:O	1.74	0.87
1:E:46:HIS:HD2	1:E:48:GLU:H	1.19	0.86
1:E:28:SER:O	1:E:31:GLN:HG2	1.85	0.77
1:E:185:ARG:HH11	1:E:185:ARG:CG	2.00	0.74
1:E:164:ALA:H	1:E:194:ASN:HD22	1.37	0.72
1:E:164:ALA:H	1:E:194:ASN:ND2	1.89	0.70
1:E:162:ASN:HB2	1:E:194:ASN:HD21	1.58	0.69

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:18:PRO:HG3	1:E:187:ASP:OD2	1.94	0.67
1:E:250:ARG:NH2	4:E:333:HOH:O	2.28	0.65
1:E:72:HIS:CD2	1:E:210:SER:HB3	2.32	0.64
1:E:107:ILE:CD1	1:E:111:MET:HE1	2.29	0.62
1:E:30:GLY:C	1:E:239:THR:HG21	2.21	0.61
1:E:30:GLY:HA2	1:E:239:THR:CG2	2.31	0.61
1:E:104:TYR:CD1	2:I:280:MSU:H21	2.35	0.60
1:E:35:VAL:HG11	1:E:77:VAL:HG11	1.83	0.60
1:E:46:HIS:CD2	1:E:48:GLU:H	2.10	0.60
1:E:100:GLY:O	2:I:283:PRO:HD3	2.02	0.59
1:E:75:GLY:HA2	1:E:79:SER:HB3	1.84	0.59
1:E:107:ILE:HD12	1:E:111:MET:CE	2.33	0.58
1:E:30:GLY:HA2	1:E:239:THR:HG23	1.85	0.57
1:E:107:ILE:HD12	1:E:111:MET:HE1	1.87	0.57
1:E:224:SER:O	1:E:228:PRO:HD3	2.05	0.56
1:E:261:LEU:H	1:E:270:ASN:ND2	1.91	0.56
1:E:8:TRP:CH2	1:E:204:PRO:HB3	2.42	0.54
1:E:13:ILE:HD11	1:E:229:HIS:HB3	1.89	0.54
1:E:261:LEU:N	1:E:270:ASN:HD21	1.92	0.54
1:E:132:SER:HB3	1:E:224:SER:CB	2.37	0.54
1:E:186:TYR:O	1:E:187:ASP:HB2	2.08	0.54
1:E:69:HIS:CD2	2:I:283:PRO:HB2	2.43	0.54
1:E:72:HIS:NE2	1:E:210:SER:HB3	2.23	0.54
1:E:80:ARG:O	1:E:86:LYS:NZ	2.43	0.51
1:E:192:PHE:CE1	1:E:222:GLY:HA2	2.46	0.50
1:E:126:GLY:HA3	1:E:238:MET:CE	2.42	0.50
1:E:128:VAL:HG22	1:E:154:MET:HB3	1.95	0.49
1:E:162:ASN:CB	1:E:194:ASN:HD21	2.25	0.49
1:E:218:ARG:HG2	1:E:220:ILE:HG23	1.95	0.49
1:E:132:SER:HB3	1:E:224:SER:OG	2.12	0.49
1:E:105:SER:HB2	4:E:413:HOH:O	2.12	0.48
1:E:38:ILE:HD11	1:E:114:VAL:HG21	1.96	0.48
1:E:133:LEU:CD2	1:E:133:LEU:C	2.82	0.47
1:E:185:ARG:CG	1:E:185:ARG:NH1	2.65	0.47
1:E:201:ILE:HG13	1:E:202:PHE:N	2.30	0.47
1:E:227:THR:HB	1:E:228:PRO:HD3	1.97	0.47
1:E:225:MET:C	1:E:228:PRO:HD2	2.36	0.46
1:E:125:LYS:HG3	1:E:239:THR:HB	1.97	0.46
1:E:50:GLU:HG2	4:E:414:HOH:O	2.15	0.46
1:E:93:VAL:HG12	1:E:95:VAL:HG13	1.96	0.46
1:E:270:ASN:ND2	4:E:308:HOH:O	2.47	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:126:GLY:HA3	1:E:238:MET:HE2	1.98	0.45
1:E:224:SER:O	1:E:228:PRO:CD	2.64	0.45
1:E:143:SER:OG	1:E:147:ARG:NH2	2.49	0.45
1:E:30:GLY:HA2	1:E:239:THR:HG21	1.99	0.44
1:E:107:ILE:HD11	1:E:111:MET:HE1	1.99	0.44
1:E:270:ASN:HD22	1:E:270:ASN:C	2.20	0.44
1:E:72:HIS:CE1	1:E:210:SER:HB3	2.53	0.43
1:E:69:HIS:CG	2:I:283:PRO:HB2	2.53	0.43
1:E:250:ARG:HE	1:E:250:ARG:HB3	1.61	0.43
1:E:227:THR:N	1:E:228:PRO:CD	2.82	0.42
1:E:58:THR:CG2	1:E:94:LYS:HD3	2.49	0.42
1:E:165:ASP:OD1	1:E:167:ARG:HB3	2.20	0.41
1:E:162:ASN:H	1:E:194:ASN:ND2	2.18	0.41
1:E:30:GLY:CA	1:E:239:THR:HG21	2.50	0.41
1:E:107:ILE:HD12	1:E:111:MET:HE2	2.03	0.41
1:E:240:LEU:HD12	1:E:240:LEU:HA	1.78	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	277/279 (99%)	267 (96%)	9 (3%)	1 (0%)	30	34
2	I	2/6 (33%)	2 (100%)	0	0	100	100
All	All	279/285 (98%)	269 (96%)	9 (3%)	1 (0%)	30	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	39	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	211/213 (99%)	190 (90%)	21 (10%)	6	6
2	I	1/1 (100%)	1 (100%)	0	100	100
All	All	212/214 (99%)	191 (90%)	21 (10%)	6	6

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	13	ILE
1	E	17	SER
1	E	39	ASP
1	E	58	THR
1	E	63	SER
1	E	106	THR
1	E	107	ILE
1	E	118	LYS
1	E	133	LEU
1	E	143	SER
1	E	154	MET
1	E	167	ARG
1	E	185	ARG
1	E	198	VAL
1	E	225	MET
1	E	230	VAL
1	E	239	THR
1	E	240	LEU
1	E	250	ARG
1	E	258	LYS
1	E	270	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	3	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	5	ASN
1	E	46	HIS
1	E	119	ASN
1	E	162	ASN
1	E	168	ASN
1	E	194	ASN
1	E	229	HIS
1	E	270	ASN
1	E	276	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ALV	I	284	2,1	4,4,5	0.98	0	1,4,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALV	I	284	2,1	-	1/2/2/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	284	ALV	O-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	279/279 (100%)	-0.07	0 100 100	8, 14, 25, 38	0
2	I	3/6 (50%)	0.88	0 100 100	26, 26, 27, 32	0
All	All	282/285 (98%)	-0.06	0 100 100	8, 14, 26, 38	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ALV	I	284	5/6	0.87	0.10	20,21,22,24	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	E	500	1/1	0.97	0.04	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.