



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 12, 2024 – 05:15 PM EST

PDB ID : 3R6L
Title : Caspase-2 T380A bound with Ac-VDVAD-CHO
Authors : Tang, Y.; Wells, J.; Arkin, M.
Deposited on : 2011-03-21
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 2% 91% 6% 2%
1	C	160	 2% 89% 9% 2%
2	B	112	 4% 79% 6% 11%
2	D	112	 3% 78% 9% 10%
3	E	6	 83% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	6	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ASA	F	406	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-2 subunit p18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1212	772	210	221	9	1	2	0
1	C	160	1250	792	216	233	9	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	174	MET	-	expression tag	UNP P42575
C	174	MET	-	expression tag	UNP P42575

- Molecule 2 is a protein called Caspase-2 subunit p12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	97	762	480	137	131	14	0	2	0
2	D	98	782	494	140	133	15	2	3	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	380	ALA	THR	engineered mutation	UNP P42575
B	453	LEU	-	expression tag	UNP P42575
B	454	GLU	-	expression tag	UNP P42575
B	455	HIS	-	expression tag	UNP P42575
B	456	HIS	-	expression tag	UNP P42575
B	457	HIS	-	expression tag	UNP P42575
B	458	HIS	-	expression tag	UNP P42575
B	459	HIS	-	expression tag	UNP P42575
B	460	HIS	-	expression tag	UNP P42575
D	380	ALA	THR	engineered mutation	UNP P42575

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	453	LEU	-	expression tag	UNP P42575
D	454	GLU	-	expression tag	UNP P42575
D	455	HIS	-	expression tag	UNP P42575
D	456	HIS	-	expression tag	UNP P42575
D	457	HIS	-	expression tag	UNP P42575
D	458	HIS	-	expression tag	UNP P42575
D	459	HIS	-	expression tag	UNP P42575
D	460	HIS	-	expression tag	UNP P42575

- Molecule 3 is a protein called Peptide Inhibitor (ACE)VDVAD-CHO.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	6	Total	C	N	O	0	0	0
			38	23	5	10			
3	E	6	Total	C	N	O	0	0	0
			38	23	5	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	129	Total	O	0	0
			129	129		
4	B	59	Total	O	0	0
			59	59		
4	C	128	Total	O	0	0
			128	128		
4	D	86	Total	O	0	0
			86	86		
4	F	4	Total	O	0	0
			4	4		
4	E	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Caspase-2 subunit p18




- Molecule 1: Caspase-2 subunit p18




- Molecule 2: Caspase-2 subunit p12




- Molecule 2: Caspase-2 subunit p12



- Molecule 3: Peptide Inhibitor (ACE)VDVAD-CHO



- Molecule 3: Peptide Inhibitor (ACE)VDVAD-CHO



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.91Å 97.06Å 98.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 50.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-1.90) 99.4 (50.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.85 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.168 , 0.194 0.169 , 0.194	Depositor DCC
R_{free} test set	2408 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.009 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4489	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASA, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.62	0/1242	0.65	0/1680
1	C	0.66	0/1278	0.72	3/1728 (0.2%)
2	B	0.70	0/786	0.75	2/1058 (0.2%)
2	D	0.79	2/810 (0.2%)	0.77	2/1088 (0.2%)
3	E	0.47	0/27	0.66	0/37
3	F	0.59	0/27	0.75	0/37
All	All	0.68	2/4170 (0.0%)	0.71	7/5628 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	440[A]	CYS	CB-SG	-7.63	1.69	1.82
2	D	440[B]	CYS	CB-SG	-7.63	1.69	1.82

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	417	ARG	NE-CZ-NH1	6.51	123.55	120.30
2	B	417	ARG	NE-CZ-NH2	-6.17	117.22	120.30
1	C	219	ARG	NE-CZ-NH2	-5.54	117.53	120.30
2	D	401[A]	MET	CG-SD-CE	5.30	108.69	100.20
2	D	401[B]	MET	CG-SD-CE	5.30	108.69	100.20
1	C	190	LEU	CA-CB-CG	5.14	127.13	115.30
1	C	301	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1212	0	1184	8	0
1	C	1250	0	1215	16	0
2	B	762	0	747	7	0
2	D	782	0	784	7	0
3	E	38	0	34	3	0
3	F	38	0	34	6	0
4	A	129	0	0	1	0
4	B	59	0	0	2	0
4	C	128	0	0	0	0
4	D	86	0	0	5	0
4	E	1	0	0	0	0
4	F	4	0	0	0	0
All	All	4489	0	3998	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:320:CYS:SG	3:F:406:ASA:C	2.21	1.29
2:D:366[A]:CYS:SG	4:D:580:HOH:O	1.98	1.21
1:A:320:CYS:SG	3:E:406:ASA:C	2.36	1.14
1:A:294:GLN:NE2	4:A:620:HOH:O	1.83	1.08
2:B:366[A]:CYS:SG	4:B:555:HOH:O	2.13	1.06
1:C:320:CYS:SG	3:F:406:ASA:HXT	2.03	0.98
1:C:187:HIS:O	1:C:190:LEU:HD22	1.82	0.79
1:A:194:LEU:HD22	1:A:199:ARG:HD2	1.70	0.72
1:C:325:THR:HG21	2:D:425:GLU:HG3	1.77	0.65
1:C:177:LYS:HD2	4:D:531:HOH:O	1.98	0.64
1:A:320:CYS:SG	3:E:406:ASA:HXT	2.34	0.63
2:D:354:PRO:O	2:D:355:LYS:HB3	2.00	0.62
4:D:584:HOH:O	3:F:401:ACE:H1	1.99	0.61
2:B:417:ARG:HD3	4:B:516:HOH:O	2.00	0.61
1:C:249:GLN:O	1:C:253:GLU:HG3	2.01	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:424:THR:HG23	2:B:426:PHE:H	1.66	0.60
1:C:248:ALA:HA	1:C:251:MET:HE2	1.87	0.57
1:A:320:CYS:SG	3:E:406:ASA:O	2.65	0.55
1:C:320:CYS:HG	3:F:406:ASA:HXT	1.75	0.52
1:C:306:PRO:HA	1:C:309:GLN:HG3	1.92	0.51
1:A:248:ALA:HA	1:A:251:MET:HE2	1.95	0.48
2:D:381:LYS:HG3	4:D:584:HOH:O	2.13	0.48
1:C:325:THR:HG21	2:D:425:GLU:CG	2.45	0.47
1:A:179[B]:CYS:SG	2:B:449:HIS:HB2	2.57	0.45
1:C:320:CYS:SG	3:F:406:ASA:O	2.67	0.45
2:B:417:ARG:HG2	2:B:418:GLU:N	2.31	0.45
1:C:205:LEU:HD23	1:C:243:LEU:HB2	1.98	0.45
1:C:200:GLY:HA2	1:C:267:THR:HB	2.00	0.44
2:B:417:ARG:HG2	2:B:418:GLU:H	1.83	0.43
1:C:320:CYS:SG	3:F:406:ASA:CA	3.00	0.43
1:C:205:LEU:HD23	1:C:243:LEU:CB	2.49	0.43
1:A:248:ALA:HA	1:A:251:MET:CE	2.47	0.43
2:D:396:GLU:CG	4:D:581:HOH:O	2.67	0.42
2:B:378:ARG:HA	2:B:384:SER:HA	2.02	0.41
1:C:248:ALA:HA	1:C:251:MET:CE	2.51	0.40
2:D:396:GLU:CG	2:D:397:ARG:HG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/160 (96%)	148 (97%)	5 (3%)	0	100	100
1	C	159/160 (99%)	155 (98%)	4 (2%)	0	100	100
2	B	97/112 (87%)	95 (98%)	2 (2%)	0	100	100
2	D	99/112 (88%)	97 (98%)	1 (1%)	1 (1%)	13	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	4/6 (67%)	4 (100%)	0	0	100	100
3	F	4/6 (67%)	4 (100%)	0	0	100	100
All	All	516/556 (93%)	503 (98%)	12 (2%)	1 (0%)	44	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	355	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/140 (94%)	126 (96%)	5 (4%)	28	21
1	C	135/140 (96%)	133 (98%)	2 (2%)	60	59
2	B	80/95 (84%)	80 (100%)	0	100	100
2	D	84/95 (88%)	82 (98%)	2 (2%)	44	39
3	E	3/3 (100%)	3 (100%)	0	100	100
3	F	3/3 (100%)	3 (100%)	0	100	100
All	All	436/476 (92%)	427 (98%)	9 (2%)	52	45

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	179[A]	CYS
1	A	179[B]	CYS
1	A	194	LEU
1	A	265	ARG
1	A	332	GLN
1	C	190	LEU
1	C	325	THR
2	D	379	ASN
2	D	451	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	257	ASN
1	A	332	GLN
1	C	257	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ASA	F	406	3	6,7,7	0.97	0	4,8,8	1.09	0
3	ASA	E	406	3	6,7,7	0.98	0	4,8,8	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASA	F	406	3	-	3/5/6/6	-
3	ASA	E	406	3	-	3/5/6/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	406	ASA	O-C-CA-CB
3	F	406	ASA	C-CA-CB-CG
3	E	406	ASA	C-CA-CB-CG
3	E	406	ASA	N-CA-CB-CG
3	F	406	ASA	N-CA-CB-CG
3	E	406	ASA	CA-CB-CG-OD2

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	406	ASA	5	0
3	E	406	ASA	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	155/160 (96%)	-0.23	3 (1%) 66 68	8, 17, 28, 33	6 (3%)
1	C	160/160 (100%)	-0.21	3 (1%) 66 68	7, 17, 30, 38	3 (1%)
2	B	97/112 (86%)	-0.17	4 (4%) 42 43	6, 14, 36, 41	2 (2%)
2	D	98/112 (87%)	-0.38	3 (3%) 51 53	4, 14, 26, 43	4 (4%)
3	E	4/6 (66%)	0.76	0 100 100	28, 30, 33, 34	0
3	F	4/6 (66%)	0.11	0 100 100	20, 21, 23, 27	0
All	All	518/556 (93%)	-0.23	13 (2%) 58 60	4, 16, 32, 43	15 (2%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	THR	5.4
2	D	354	PRO	4.3
2	B	380	ALA	3.6
2	D	451	PRO	3.2
2	B	451	PRO	2.7
1	C	333	ASP	2.5
2	B	450	PRO	2.4
1	C	324	GLU	2.4
1	A	325	THR	2.4
1	C	190	LEU	2.1
2	D	355	LYS	2.1
2	B	424	THR	2.1
1	A	216	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ASA	E	406	8/8	0.85	0.14	33,33,33,34	0
3	ASA	F	406	8/8	0.93	0.09	22,23,24,25	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.