



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2024 – 02:29 PM EST

PDB ID : 3UJI  
Title : Crystal structure of anti-HIV-1 V3 Fab 2558 in complex with MN peptide  
Authors : Kong, X.P.  
Deposited on : 2011-11-07  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	209	 83% 15%
2	H	223	 % 88% 9%
3	P	23	 4% 70% 22%
4	A	3	 33% 67%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4113 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Light chain of anti-HIV-1 V3 monoclonal antibody 2558.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	208	1554	977	252	320	5	0	0	0


- Molecule 2 is a protein called Fab region of the heavy chain of anti-HIV-1 V3 monoclonal antibody 2558.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	218	1650	1055	264	324	7	0	0	0

- Molecule 3 is a protein called Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	P	18	154	99	32	23	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	A	3	38	22	2	14	0	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	1	Total Ca 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	L	372	Total O 372 372	0	0
6	H	324	Total O 324 324	0	0
6	P	20	Total O 20 20	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Light chain of anti-HIV-1 V3 monoclonal antibody 2558



- Molecule 2: Fab region of the heavy chain of anti-HIV-1 V3 monoclonal antibody 2558



- Molecule 3: Envelope glycoprotein gp160



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.35Å 73.10Å 88.66Å 90.00° 121.98° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 50.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-1.60) 99.1 (50.00-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.10 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.161 , 0.194 0.161 , 0.193	Depositor DCC
$R_{free}$ test set	3660 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.1	Xtrriage
Anisotropy	0.021	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4113	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	1.47	7/1592 (0.4%)	1.47	13/2175 (0.6%)
2	H	1.35	5/1696 (0.3%)	1.28	6/2310 (0.3%)
3	P	1.30	0/158	1.45	2/210 (1.0%)
All	All	1.41	12/3446 (0.3%)	1.38	21/4695 (0.4%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	61	ARG	CZ-NH1	7.31	1.42	1.33
2	H	82	TRP	CD2-CE2	6.50	1.49	1.41
2	H	164	HIS	CA-CB	-6.34	1.40	1.53
1	L	50	GLU	CD-OE1	6.07	1.32	1.25
1	L	34	SER	CA-CB	5.95	1.61	1.52
2	H	33	TRP	CD2-CE2	5.83	1.48	1.41
1	L	175	SER	CB-OG	-5.25	1.35	1.42
2	H	148	GLU	CD-OE1	5.22	1.31	1.25
1	L	191	TYR	CG-CD1	5.20	1.46	1.39
1	L	9	SER	CB-OG	-5.08	1.35	1.42
2	H	177	SER	CB-OG	5.05	1.48	1.42
1	L	91	TRP	CD2-CE2	5.03	1.47	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	82	ASP	CB-CG-OD2	-8.99	110.21	118.30
1	L	61	ARG	NE-CZ-NH2	8.95	124.78	120.30
1	L	51	ASP	CB-CG-OD1	7.55	125.09	118.30
1	L	61	ARG	NE-CZ-NH1	-7.06	116.77	120.30
2	H	101	ASP	CB-CG-OD2	-6.85	112.13	118.30
1	L	180	LEU	CB-CG-CD1	-6.30	100.28	111.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	94	ARG	NE-CZ-NH1	-6.17	117.21	120.30
1	L	124	GLU	OE1-CD-OE2	6.14	130.67	123.30
1	L	60	ASP	CB-CG-OD1	-5.98	112.92	118.30
1	L	198	GLU	OE1-CD-OE2	5.97	130.46	123.30
2	H	38	ARG	NE-CZ-NH2	5.79	123.19	120.30
2	H	101	ASP	CB-CG-OD1	5.66	123.39	118.30
1	L	82	ASP	OD1-CG-OD2	5.61	133.95	123.30
1	L	110	LYS	CD-CE-NZ	-5.59	98.84	111.70
1	L	60	ASP	N-CA-C	5.48	125.81	111.00
3	P	304	ARG	NE-CZ-NH2	5.42	123.01	120.30
2	H	100	ASP	CB-CG-OD2	-5.36	113.48	118.30
3	P	315	ARG	NE-CZ-NH1	5.31	122.96	120.30
1	L	172	TYR	CB-CG-CD2	-5.30	117.82	121.00
2	H	12	LYS	CD-CE-NZ	-5.15	99.85	111.70
1	L	30	ASP	CB-CG-OD1	5.14	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1554	0	1509	16	0
2	H	1650	0	1585	13	0
3	P	154	0	158	4	0
4	A	38	0	34	2	0
5	L	1	0	0	0	0
6	H	324	0	0	3	1
6	L	372	0	0	2	3
6	P	20	0	0	0	1
All	All	4113	0	3286	31	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:315:ARG:HH11	3:P:315:ARG:HG3	1.08	1.16
2:H:6:GLN:H	2:H:105:GLN:HE22	1.10	0.98
1:L:181:THR:H	1:L:184:GLN:HE21	1.06	0.97
2:H:105:GLN:HE21	2:H:105:GLN:H	1.15	0.91
3:P:315:ARG:HH11	3:P:315:ARG:CG	1.91	0.81
3:P:315:ARG:HG3	3:P:315:ARG:NH1	1.86	0.81
2:H:70:SER:OG	2:H:81:HIS:HE1	1.79	0.66
2:H:6:GLN:H	2:H:105:GLN:NE2	1.91	0.64
2:H:6:GLN:HE21	2:H:104:GLY:HA3	1.63	0.64
1:L:167:GLN:HE21	1:L:169:ASN:HD21	1.46	0.63
1:L:203:GLU:OE1	6:L:308:HOH:O	2.16	0.61
1:L:2:TYR:N	4:A:1:NAG:HO3	1.98	0.60
1:L:2:TYR:N	4:A:2:NAG:HO6	2.00	0.60
1:L:181:THR:H	1:L:184:GLN:NE2	1.90	0.58
2:H:164:HIS:ND1	6:H:570:HOH:O	2.33	0.55
1:L:181:THR:N	1:L:184:GLN:HE21	1.90	0.52
2:H:164:HIS:CD2	6:H:488:HOH:O	2.63	0.50
1:L:167:GLN:HE21	1:L:169:ASN:ND2	2.10	0.49
1:L:181:THR:OG1	1:L:184:GLN:HG3	2.13	0.49
2:H:105:GLN:HE21	2:H:105:GLN:N	1.97	0.48
1:L:120:PRO:HD3	1:L:132:LEU:CD2	2.44	0.47
2:H:105:GLN:H	2:H:105:GLN:NE2	1.98	0.47
1:L:169:ASN:C	1:L:169:ASN:HD22	2.17	0.47
2:H:18:LEU:HD12	6:H:409:HOH:O	2.16	0.45
3:P:315:ARG:NH1	3:P:315:ARG:CG	2.60	0.45
1:L:185:TRP:CZ2	1:L:208:PRO:HA	2.52	0.45
2:H:6:GLN:HE22	2:H:91:TYR:HA	1.82	0.43
1:L:169:ASN:ND2	1:L:171:LYS:H	2.17	0.42
1:L:96:VAL:HB	2:H:47:TRP:CG	2.56	0.41
1:L:184:GLN:NE2	6:L:345:HOH:O	2.39	0.41
1:L:160:GLU:HB3	2:H:169:VAL:HG21	2.02	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:L:296:HOH:O	6:L:405:HOH:O[3_545]	1.75	0.45
6:L:350:HOH:O	6:L:373:HOH:O[3_545]	1.94	0.26
6:H:409:HOH:O	6:P:255:HOH:O[4_445]	1.95	0.25
6:L:313:HOH:O	6:L:336:HOH:O[4_556]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	206/209 (99%)	202 (98%)	4 (2%)	0	100	100
2	H	214/223 (96%)	210 (98%)	4 (2%)	0	100	100
3	P	16/23 (70%)	16 (100%)	0	0	100	100
All	All	436/455 (96%)	428 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	176/177 (99%)	169 (96%)	7 (4%)	27	8
2	H	183/188 (97%)	180 (98%)	3 (2%)	58	37
3	P	15/19 (79%)	14 (93%)	1 (7%)	13	3
All	All	374/384 (97%)	363 (97%)	11 (3%)	37	15

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	2	TYR
1	L	56	SER
1	L	89	GLN
1	L	155	VAL
1	L	169	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	L	183	GLU
1	L	189	LYS
2	H	105	GLN
2	H	161	SER
2	H	167	PRO
3	P	315	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	42	GLN
1	L	79	GLN
1	L	89	GLN
1	L	108	GLN
1	L	169	ASN
1	L	170	ASN
1	L	184	GLN
2	H	1	GLN
2	H	6	GLN
2	H	66	GLN
2	H	81	HIS
2	H	105	GLN
2	H	164	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1	2,4	14,14,15	1.06	1 (7%)	17,19,21	1.97	4 (23%)
4	NAG	A	2	4	14,14,15	1.65	2 (14%)	17,19,21	2.45	7 (41%)
4	FUC	A	3	4	10,10,11	0.98	0	14,14,16	1.74	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	A	2	4	-	0/6/23/26	0/1/1/1
4	FUC	A	3	4	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2	NAG	C8-C7	4.38	1.59	1.50
4	A	2	NAG	C2-N2	3.63	1.52	1.46
4	A	1	NAG	O7-C7	2.30	1.28	1.23

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2	NAG	O5-C1-C2	-5.35	103.02	111.29
4	A	1	NAG	O5-C1-C2	-4.56	104.24	111.29
4	A	2	NAG	O7-C7-N2	4.38	129.73	121.98
4	A	2	NAG	O5-C5-C6	-3.67	100.51	107.66
4	A	3	FUC	C1-C2-C3	-3.57	104.45	109.64
4	A	1	NAG	C4-C3-C2	3.30	115.86	111.02
4	A	2	NAG	C2-N2-C7	-3.28	118.51	122.90
4	A	1	NAG	C3-C4-C5	-3.26	104.32	110.23
4	A	2	NAG	C3-C4-C5	-3.17	104.48	110.23
4	A	1	NAG	C1-C2-N2	2.75	114.77	110.43
4	A	2	NAG	O7-C7-C8	-2.47	117.65	122.05
4	A	3	FUC	C2-C3-C4	2.43	115.14	110.86
4	A	3	FUC	C3-C4-C5	-2.35	106.25	109.81

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2	NAG	C8-C7-N2	-2.19	112.48	116.12
4	A	3	FUC	O4-C4-C3	-2.14	105.32	110.38
4	A	3	FUC	O2-C2-C3	2.08	114.45	110.15
4	A	3	FUC	C6-C5-C4	2.07	116.86	113.08

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1	NAG	C1-C2-N2-C7
4	A	1	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	NAG	1	0
4	A	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	208/209 (99%)	-0.62	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	9, 15, 31, 53	0
2	H	218/223 (97%)	-0.49	3 (1%) <span style="border: 1px solid blue; padding: 2px;">73</span> <span style="border: 1px solid blue; padding: 2px;">76</span>	11, 17, 38, 71	0
3	P	18/23 (78%)	-0.18	1 (5%) <span style="border: 1px solid red; padding: 2px;">31</span> <span style="border: 1px solid red; padding: 2px;">30</span>	12, 18, 42, 57	0
All	All	444/455 (97%)	-0.54	4 (0%) <span style="border: 1px solid blue; padding: 2px;">81</span> <span style="border: 1px solid blue; padding: 2px;">83</span>	9, 17, 34, 71	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	133	GLY	3.0
2	H	127	SER	2.5
3	P	301	TYR	2.4
2	H	134	GLY	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CA	L	209	1/1	1.00	0.05	17,17,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.