



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 19, 2024 – 04:37 AM EDT

PDB ID : 4CN9  
Title : structure of proximal thread matrix protein 1 (PTMP1) from the mussel byssus with zinc occupied MIDAS motif  
Authors : Gertz, M.; Suhre, M.H.; Scheibel, T.; Steegborn, C.  
Deposited on : 2014-01-21  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	 2% 80% 6% 13%
1	B	454	 3% 82% 5% 13%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12621 atoms, of which 6110 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROXIMAL THREAD MATRIX PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	397	6070	1905	3037	514	598	16	0	1	0
1	B	396	6069	1905	3039	513	595	17	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q8T5C2
A	10	LYS	GLU	conflict	UNP Q8T5C2
A	38	ARG	GLN	conflict	UNP Q8T5C2
A	328	THR	SER	conflict	UNP Q8T5C2
B	0	SER	-	expression tag	UNP Q8T5C2
B	10	LYS	GLU	conflict	UNP Q8T5C2
B	38	ARG	GLN	conflict	UNP Q8T5C2
B	328	THR	SER	conflict	UNP Q8T5C2

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	10	2	6	2	0	0
3	B	1	10	2	6	2	0	0
3	B	1	10	2	6	2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	14	3	8	3	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C H O 14 3 8 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	228	Total O 228 228	0	0
6	B	170	Total O 170 170	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROXIMAL THREAD MATRIX PROTEIN 1



- Molecule 1: PROXIMAL THREAD MATRIX PROTEIN 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.54Å 63.07Å 122.35Å 90.00° 101.83° 90.00°	Depositor
Resolution (Å)	43.56 – 1.90 43.56 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.56-1.90) 99.9 (43.56-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 1.89Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.173 , 0.208 0.177 , 0.212	Depositor DCC
$R_{free}$ test set	3626 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtrriage
Anisotropy	0.277	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 50.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12621	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3079	0.58	0/4145
1	B	0.39	0/3079	0.55	0/4144
All	All	0.40	0/6158	0.56	0/8289

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3033	3037	3032	20	2
1	B	3030	3039	3037	17	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	4	6	6	1	0
3	B	8	12	12	3	0
4	A	12	8	16	4	0
4	B	12	8	16	2	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	228	0	0	9	2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	170	0	0	5	2
All	All	6511	6110	6119	43	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:ASP:OD2	6:A:2027:HOH:O	1.99	0.79
6:A:2146:HOH:O	1:B:287:ASN:OD1	2.07	0.72
4:A:1453:GOL:O3	6:A:2227:HOH:O	2.00	0.71
1:A:184:HIS:NE2	1:A:188:LYS:HE3	2.06	0.70
3:A:1450:EDO:O1	6:A:2183:HOH:O	2.11	0.62
3:B:1450:EDO:H22	6:B:2119:HOH:O	2.01	0.60
1:B:184:HIS:CD2	1:B:188:LYS:HE3	2.36	0.60
4:A:1451:GOL:O2	6:A:2226:HOH:O	2.11	0.60
1:B:147:HIS:ND1	4:B:1454:GOL:H2	2.17	0.59
1:A:317:LYS:HB2	4:A:1451:GOL:H11	1.84	0.59
1:B:117:LYS:NZ	6:B:2041:HOH:O	2.34	0.59
1:B:147:HIS:HB2	4:B:1454:GOL:H31	1.85	0.57
1:A:183:GLN:OE1	1:A:212:GLN:HB3	2.05	0.56
1:A:183:GLN:OE1	1:A:212:GLN:CB	2.56	0.53
1:A:317:LYS:HG3	4:A:1451:GOL:H32	1.90	0.52
1:A:164:LYS:HE2	6:A:2081:HOH:O	2.10	0.51
1:A:325:ASP:OD2	6:A:2182:HOH:O	2.19	0.51
1:B:358:GLU:O	1:B:359:GLU:HB3	2.11	0.51
1:A:117:LYS:NZ	6:A:2053:HOH:O	2.32	0.49
1:A:339:ASP:OD2	1:A:375:HIS:HD2	1.95	0.49
1:A:272:TYR:CE2	1:A:276:LYS:HE3	2.48	0.48
3:B:1450:EDO:H11	6:B:2134:HOH:O	2.14	0.47
1:B:184:HIS:NE2	1:B:188:LYS:CE	2.78	0.47
1:A:93:ASN:OD1	1:A:93:ASN:N	2.48	0.46
1:A:208:THR:O	1:A:212:GLN:HG3	2.16	0.46
1:A:358:GLU:O	1:A:359:GLU:CB	2.64	0.46
1:A:47:GLU:HB3	1:A:445:HIS:O	2.15	0.46
1:B:183:GLN:OE1	1:B:212:GLN:HB3	2.17	0.45
1:B:235:LEU:HG	6:B:2092:HOH:O	2.15	0.45
1:B:259:ASP:OD2	1:B:369:ASP:OD1	2.34	0.45
1:B:262:SER:OG	1:B:331:ILE:C	2.55	0.45
1:A:252:ALA:HA	1:A:360:VAL:CG1	2.47	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:97:GLY:O	1:B:164:LYS:HE3	2.17	0.45
3:B:1450:EDO:C2	6:B:2119:HOH:O	2.61	0.45
1:A:206:VAL:HG22	1:A:209:GLU:HB3	1.99	0.44
1:B:267:ASN:O	1:B:268:ASN:HB2	2.17	0.44
1:A:259:ASP:OD2	1:A:369:ASP:OD1	2.36	0.44
1:B:245:VAL:HG12	1:B:246:VAL:N	2.34	0.43
1:B:272:TYR:CE2	1:B:276:LYS:HE3	2.55	0.42
1:B:267:ASN:O	1:B:268:ASN:CB	2.69	0.41
1:A:424:LYS:CE	6:A:2218:HOH:O	2.68	0.41
1:A:183:GLN:OE1	1:A:183:GLN:HA	2.21	0.40
1:B:184:HIS:NE2	1:B:188:LYS:HE2	2.36	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303:ASP:OD2	6:B:2123:HOH:O[1_565]	1.95	0.25
1:A:423:ASP:OD2	6:A:2027:HOH:O[1_655]	2.02	0.18
6:A:2169:HOH:O	6:B:2046:HOH:O[2_655]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/454 (87%)	383 (97%)	9 (2%)	2 (0%)	29	18
1	B	394/454 (87%)	384 (98%)	8 (2%)	2 (0%)	29	18
All	All	788/908 (87%)	767 (97%)	17 (2%)	4 (0%)	29	18

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	B	268	ASN
1	A	106	SER
1	B	106	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/378 (89%)	329 (98%)	8 (2%)	49	43
1	B	337/378 (89%)	334 (99%)	3 (1%)	78	79
All	All	674/756 (89%)	663 (98%)	11 (2%)	62	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ASP
1	A	47	GLU
1	A	206	VAL
1	A	241	CYS
1	A	357	ARG
1	A	359	GLU
1	A	367	LEU
1	A	369	ASP
1	B	48	GLU
1	B	241	CYS
1	B	369	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	375	HIS
1	B	287	ASN
1	B	380	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	1451	-	5,5,5	0.58	0	5,5,5	0.32	0
5	SO4	A	1452	-	4,4,4	0.30	0	6,6,6	0.28	0
3	EDO	A	1450	-	3,3,3	0.35	0	2,2,2	0.26	0
4	GOL	A	1453	-	5,5,5	0.32	0	5,5,5	0.32	0
4	GOL	B	1451	-	5,5,5	0.38	0	5,5,5	0.30	0
3	EDO	B	1450	-	3,3,3	0.29	0	2,2,2	1.05	0
5	SO4	B	1452	-	4,4,4	0.24	0	6,6,6	0.32	0
3	EDO	B	1453	-	3,3,3	0.35	0	2,2,2	0.36	0
4	GOL	B	1454	-	5,5,5	0.26	0	5,5,5	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1451	-	-	4/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1450	-	-	0/1/1/1	-
4	GOL	A	1453	-	-	4/4/4/4	-
4	GOL	B	1451	-	-	4/4/4/4	-
3	EDO	B	1450	-	-	1/1/1/1	-
3	EDO	B	1453	-	-	1/1/1/1	-
4	GOL	B	1454	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1451	GOL	O1-C1-C2-C3
4	A	1451	GOL	C1-C2-C3-O3
4	B	1451	GOL	O1-C1-C2-C3
4	B	1454	GOL	C1-C2-C3-O3
4	B	1451	GOL	O1-C1-C2-O2
4	B	1451	GOL	O2-C2-C3-O3
4	A	1453	GOL	O1-C1-C2-C3
4	A	1453	GOL	C1-C2-C3-O3
4	B	1451	GOL	C1-C2-C3-O3
4	A	1451	GOL	O1-C1-C2-O2
4	A	1451	GOL	O2-C2-C3-O3
4	B	1454	GOL	O2-C2-C3-O3
4	A	1453	GOL	O1-C1-C2-O2
4	A	1453	GOL	O2-C2-C3-O3
3	B	1453	EDO	O1-C1-C2-O2
3	B	1450	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1451	GOL	3	0
3	A	1450	EDO	1	0
4	A	1453	GOL	1	0
3	B	1450	EDO	3	0
4	B	1454	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	397/454 (87%)	0.06	10 (2%) 57 60	15, 25, 43, 89	0
1	B	396/454 (87%)	0.08	15 (3%) 40 43	16, 30, 52, 83	0
All	All	793/908 (87%)	0.07	25 (3%) 47 50	15, 27, 49, 89	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	ARG	8.7
1	B	95	ARG	8.5
1	A	96	ASN	7.1
1	B	267	ASN	6.4
1	A	93	ASN	5.8
1	B	358	GLU	5.5
1	B	96	ASN	4.9
1	B	359	GLU	4.6
1	A	45	ASP	4.5
1	B	92	VAL	4.4
1	B	93	ASN	4.4
1	A	358	GLU	4.2
1	A	359	GLU	3.8
1	A	92	VAL	3.8
1	B	447	LYS	3.4
1	B	91	GLY	3.4
1	B	266	ALA	2.9
1	A	443	ARG	2.9
1	B	441	LYS	2.6
1	B	94	GLY	2.4
1	A	267	ASN	2.3
1	B	97	GLY	2.2
1	B	443	ARG	2.2
1	A	442	PRO	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	357	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	1453	6/6	0.72	0.21	35,41,43,45	0
4	GOL	B	1451	6/6	0.75	0.14	52,62,71,73	0
4	GOL	B	1454	6/6	0.78	0.12	24,30,41,41	0
4	GOL	A	1451	6/6	0.81	0.18	30,45,50,60	0
2	ZN	B	1448	1/1	0.88	0.08	46,46,46,46	1
3	EDO	B	1453	4/4	0.90	0.13	37,45,48,52	0
3	EDO	B	1450	4/4	0.92	0.11	30,37,44,44	0
2	ZN	A	1449	1/1	0.93	0.06	33,33,33,33	1
3	EDO	A	1450	4/4	0.95	0.11	24,29,43,48	0
5	SO4	B	1452	5/5	0.98	0.08	28,32,39,44	0
5	SO4	A	1452	5/5	0.99	0.08	26,26,38,40	0
2	ZN	A	1448	1/1	0.99	0.09	25,25,25,25	1
2	ZN	B	1449	1/1	1.00	0.09	30,30,30,30	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.