



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 08:38 am GMT

PDB ID : 4CVY  
Title : Crystal structure of the M. tuberculosis sulfate ester dioxygenase Rv3406 in complex with iron.  
Authors : Neres, J.; Hartkoorn, R.C.; Chiarelli, L.R.; Gadupudi, R.; Pasca, M.; Mori, G.; Farina, D.; Salina, S.; Makarov, V.; Kolly, G.S.; Molteni, E.; Binda, C.; Dhar, N.; Ferrari, S.; Brodin, P.; Delorme, V.; Landry, V.; de Jesus Lopes Ribeiro, A.L.; Saxena, P.; Pojer, F.; Venturelli, A.; Carta, A.; Luciani, R.; Porta, A.; Zanoni, G.; De Rossi, E.; Costi, M.P.; Riccardi, G.; Cole, S.T.  
Deposited on : 2014-03-31  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	 2% 73% 8% 19%
1	B	295	 2% 71% 8% 19%
1	C	295	 3% 66% 13% 20%

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.36

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	D	295	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '3%', a large green segment labeled '68%', a small yellow segment labeled '10%', and a grey segment at the end labeled '19%'. The total length of the bar represents 100%.</p>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIOXYGENASE RV3406/MT3514.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	Total 1882	C 1188	N 344	O 345	S 5	0	0	0
1	B	240	Total 1880	C 1186	N 347	O 343	S 4	0	0	0
1	C	236	Total 1855	C 1173	N 340	O 338	S 4	0	0	0
1	D	235	Total 1849	C 1168	N 339	O 338	S 4	0	0	0

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	N O		
2	A	1	Total 4	N O 1 3	0	0

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe 1 1	0	0
3	B	1	Total Fe 1 1	0	0
3	C	1	Total Fe 1 1	0	0
3	D	1	Total Fe 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	50	Total O 50 50	0	0
4	B	54	Total O 54 54	0	0
4	C	42	Total O 42 42	0	0
4	D	42	Total O 42 42	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DIOXYGENASE RV3406/MT3514




- Molecule 1: DIOXYGENASE RV3406/MT3514



- Molecule 1: DIOXYGENASE RV3406/MT3514



- Molecule 1: DIOXYGENASE RV3406/MT3514



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.97Å 128.15Å 138.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.04 – 2.00 43.41 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.8 (94.04-2.00) 97.8 (43.41-2.00)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, $R_{free}$	0.208 , 0.254 0.217 , 0.259	Depositor DCC
$R_{free}$ test set	3773 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.8	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7662	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/1926	0.92	3/2624 (0.1%)
1	B	0.81	1/1925 (0.1%)	0.95	6/2623 (0.2%)
1	C	0.71	0/1899	0.94	5/2588 (0.2%)
1	D	0.78	0/1892	0.99	8/2578 (0.3%)
All	All	0.77	1/7642 (0.0%)	0.95	22/10413 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	138	GLU	CG-CD	5.18	1.59	1.51

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	47	ARG	NE-CZ-NH2	-11.45	114.57	120.30
1	B	47	ARG	NE-CZ-NH2	-10.16	115.22	120.30
1	A	47	ARG	NE-CZ-NH2	-9.94	115.33	120.30
1	B	47	ARG	NE-CZ-NH1	9.62	125.11	120.30
1	D	47	ARG	NE-CZ-NH1	7.71	124.15	120.30
1	C	240	ASP	CB-CG-OD1	7.45	125.01	118.30
1	C	224	ARG	NE-CZ-NH2	-6.82	116.89	120.30
1	A	47	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	D	233	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	D	210	ASP	CB-CG-OD1	6.13	123.82	118.30
1	C	47	ARG	NE-CZ-NH1	5.98	123.29	120.30
1	D	21	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	272	ASP	CB-CG-OD1	5.88	123.59	118.30
1	D	113	ARG	NE-CZ-NH2	-5.55	117.52	120.30
1	D	113	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	B	266	ARG	NE-CZ-NH1	5.44	123.02	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	233	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	27	ASP	CB-CG-OD1	5.34	123.10	118.30
1	C	35	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	C	224	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	266	ARG	NE-CZ-NH1	5.20	122.90	120.30
1	B	113	ARG	CG-CD-NE	-5.06	101.18	111.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1882	0	1840	17	0
1	B	1880	0	1840	23	0
1	C	1855	0	1818	27	0
1	D	1849	0	1810	22	0
2	A	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	50	0	0	0	0
4	B	54	0	0	0	0
4	C	42	0	0	1	0
4	D	42	0	0	3	0
All	All	7662	0	7308	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:ARG:HD3	1:A:240:ASP:OD1	1.77	0.84

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:222:GLN:HE22	1:D:250:GLN:HE22	1.27	0.83
1:D:47:ARG:HD3	1:D:240:ASP:OD1	1.84	0.77
1:B:54:ASP:OD1	1:B:113:ARG:NH2	2.18	0.75
1:B:178:ASP:C	1:B:178:ASP:OD1	2.28	0.72
1:B:216:VAL:HG13	1:D:216:VAL:HG13	1.71	0.71
1:A:116:SER:OG	1:A:261:HIS:NE2	2.24	0.71
1:B:47:ARG:CD	1:B:240:ASP:OD1	2.41	0.69
1:D:54:ASP:OD1	1:D:113:ARG:NH2	2.22	0.68
1:B:47:ARG:HD3	1:B:240:ASP:OD1	1.94	0.67
1:A:216:VAL:HG13	1:C:216:VAL:HG13	1.78	0.65
1:D:47:ARG:CD	1:D:240:ASP:OD1	2.45	0.65
1:C:3:LEU:HD13	1:C:4:ILE:HG23	1.81	0.63
1:A:216:VAL:HG13	1:C:216:VAL:CG1	2.29	0.62
1:C:101:THR:OG1	1:C:151:HIS:CD2	2.53	0.61
1:B:216:VAL:CG1	1:D:216:VAL:HG13	2.33	0.59
1:A:6:VAL:O	1:A:6:VAL:HG23	2.03	0.58
1:C:101:THR:OG1	1:C:151:HIS:HD2	1.85	0.58
1:A:47:ARG:CD	1:A:240:ASP:OD1	2.52	0.57
1:C:138:GLU:N	1:C:139:PRO:CD	2.68	0.57
1:C:47:ARG:HD3	1:C:240:ASP:OD1	2.05	0.56
1:A:216:VAL:CG1	1:C:216:VAL:HG13	2.35	0.56
1:C:47:ARG:CD	1:C:240:ASP:OD1	2.54	0.56
1:D:57:GLN:NE2	1:D:113:ARG:HE	2.03	0.55
1:B:47:ARG:HD2	1:B:240:ASP:OD1	2.06	0.54
1:B:101:THR:OG1	1:B:151:HIS:HD2	1.91	0.54
1:C:94:ASN:HD22	1:C:255:ASP:HB2	1.73	0.53
1:A:216:VAL:CG1	1:C:216:VAL:CG1	2.88	0.51
1:A:54:ASP:OD1	1:A:113:ARG:NH2	2.41	0.50
1:C:120:TYR:OH	1:C:258:ASP:OD2	2.14	0.50
1:D:276:ASP:HB2	4:D:2027:HOH:O	2.12	0.50
1:D:57:GLN:HE21	1:D:113:ARG:HE	1.60	0.49
1:B:4:ILE:HG12	1:B:33:GLU:OE2	2.13	0.49
1:B:216:VAL:HG13	1:D:216:VAL:CG1	2.41	0.49
1:C:21:ARG:HE	1:C:51:GLN:HG3	1.78	0.49
1:A:57:GLN:HE21	1:A:113:ARG:HE	1.60	0.48
1:B:99:ASP:HB3	1:B:100:VAL:HG23	1.95	0.48
1:D:101:THR:O	1:D:183:HIS:NE2	2.47	0.48
1:D:277:VAL:HG22	4:D:2027:HOH:O	2.14	0.48
1:C:107:PRO:HB3	1:C:268:THR:CG2	2.44	0.47
1:B:153:ASN:O	1:B:154:ARG:C	2.53	0.47
1:A:224:ARG:HD2	1:C:213:GLU:OE2	2.14	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96:TRP:CH2	1:B:125:LEU:HB3	2.50	0.46
1:B:266:ARG:HH11	1:B:266:ARG:CG	2.28	0.46
1:D:138:GLU:N	1:D:139:PRO:CD	2.79	0.45
1:B:266:ARG:HH11	1:B:266:ARG:HB3	1.81	0.45
1:B:178:ASP:OD1	1:B:178:ASP:O	2.33	0.45
1:C:54:ASP:OD1	1:C:113:ARG:NH2	2.43	0.45
1:D:215:ARG:HB3	4:D:2031:HOH:O	2.15	0.45
1:A:276:ASP:OD2	1:A:280:GLN:NE2	2.48	0.45
1:A:6:VAL:O	1:A:6:VAL:CG2	2.65	0.45
1:C:17:ILE:HD12	1:C:46:PHE:CZ	2.52	0.44
1:C:99:ASP:HB3	1:C:100:VAL:HG23	2.00	0.44
1:D:4:ILE:HG22	1:D:20:VAL:HG22	2.00	0.44
1:B:215:ARG:NH2	1:B:219:GLU:OE2	2.49	0.43
1:C:47:ARG:HD2	1:C:240:ASP:OD1	2.17	0.43
1:C:126:TRP:CD1	1:C:251:HIS:HB3	2.54	0.43
1:A:213:GLU:OE2	1:C:224:ARG:HD2	2.18	0.43
1:A:4:ILE:HG21	1:A:20:VAL:CG2	2.48	0.43
1:A:144:THR:HB	1:A:186:VAL:HG23	2.01	0.42
1:B:138:GLU:N	1:B:139:PRO:HD2	2.33	0.42
1:C:61:ALA:CB	1:C:267:VAL:HG11	2.50	0.42
1:B:213:GLU:OE2	1:D:224:ARG:HD2	2.19	0.42
1:D:127:ALA:O	1:D:249:THR:HA	2.19	0.42
1:B:266:ARG:HH11	1:B:266:ARG:HG2	1.84	0.42
1:C:215:ARG:CZ	4:C:2026:HOH:O	2.67	0.42
1:D:276:ASP:OD1	1:D:278:TYR:N	2.49	0.41
1:D:7:LYS:O	1:D:15:ALA:HA	2.19	0.41
1:C:22:LEU:HD22	1:C:60:PHE:CE1	2.55	0.41
1:A:138:GLU:N	1:A:139:PRO:CD	2.84	0.41
1:B:210:ASP:OD1	1:B:210:ASP:C	2.59	0.41
1:D:149:ALA:O	1:D:182:GLU:HA	2.20	0.41
1:B:139:PRO:HB2	1:D:143:LEU:HD12	2.03	0.41
1:C:4:ILE:HB	1:C:20:VAL:HG23	2.03	0.41
1:D:137:PRO:HD2	1:D:140:LEU:HD12	2.04	0.40
1:B:209:LEU:HD23	1:B:217:LEU:HD12	2.03	0.40
1:C:17:ILE:HB	1:C:46:PHE:CE2	2.57	0.40
1:C:22:LEU:HD22	1:C:60:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/295 (79%)	225 (96%)	9 (4%)	0	100	100
1	B	234/295 (79%)	226 (97%)	8 (3%)	0	100	100
1	C	230/295 (78%)	218 (95%)	11 (5%)	1 (0%)	34	30
1	D	229/295 (78%)	222 (97%)	7 (3%)	0	100	100
All	All	927/1180 (79%)	891 (96%)	35 (4%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	199	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/238 (82%)	190 (97%)	6 (3%)	40	40
1	B	195/238 (82%)	184 (94%)	11 (6%)	21	17
1	C	193/238 (81%)	184 (95%)	9 (5%)	26	22
1	D	193/238 (81%)	187 (97%)	6 (3%)	40	40
All	All	777/952 (82%)	745 (96%)	32 (4%)	30	28

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LEU
1	A	100	VAL
1	A	107	PRO
1	A	113	ARG
1	A	178	ASP
1	A	204	ARG
1	B	4	ILE
1	B	6	VAL
1	B	25	ASP
1	B	58	LEU
1	B	100	VAL
1	B	113	ARG
1	B	142	CYS
1	B	154	ARG
1	B	178	ASP
1	B	228	PRO
1	B	266	ARG
1	C	3	LEU
1	C	4	ILE
1	C	6	VAL
1	C	8	LYS
1	C	25	ASP
1	C	52	LEU
1	C	58	LEU
1	C	142	CYS
1	C	187	ARG
1	D	4	ILE
1	D	35	ARG
1	D	100	VAL
1	D	113	ARG
1	D	178	ASP
1	D	216	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	265	HIS
1	B	41	HIS
1	B	151	HIS
1	C	57	GLN
1	C	151	HIS
1	C	265	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	57	GLN
1	D	222	GLN
1	D	265	HIS
1	D	280	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NO3	A	301	-	1,3,3	3.38	1 (100%)	0,3,3	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	NO3	O1-N	3.38	1.39	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	240/295 (81%)	0.08	6 (2%) 57 56	26, 39, 61, 78	0
1	B	240/295 (81%)	0.03	6 (2%) 57 56	24, 34, 60, 89	0
1	C	236/295 (80%)	0.04	8 (3%) 45 44	23, 38, 64, 85	0
1	D	235/295 (79%)	-0.02	10 (4%) 35 34	24, 37, 64, 84	0
All	All	951/1180 (80%)	0.04	30 (3%) 47 46	23, 37, 62, 89	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	ASP	5.6
1	D	286	SER	5.3
1	C	68	ILE	4.2
1	C	179	PHE	3.9
1	B	177	PRO	3.8
1	A	4	ILE	3.6
1	B	261	HIS	3.5
1	B	288	ALA	3.4
1	D	179	PHE	3.3
1	D	3	LEU	3.2
1	C	284	VAL	3.0
1	A	21	ARG	2.9
1	C	259	ASP	2.9
1	C	21	ARG	2.9
1	D	180	ARG	2.9
1	A	259	ASP	2.8
1	D	285	ILE	2.6
1	C	25	ASP	2.5
1	C	27	ASP	2.5
1	A	5	THR	2.5
1	D	270	MET	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	154	ARG	2.4
1	D	28	PRO	2.4
1	D	68	ILE	2.4
1	C	178	ASP	2.3
1	D	65	GLY	2.3
1	A	94	ASN	2.3
1	A	179	PHE	2.2
1	D	4	ILE	2.2
1	B	287	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NO3	A	301	4/4	0.85	0.11	40,44,46,50	0
3	FE	C	1286	1/1	0.96	0.06	57,57,57,57	0
3	FE	D	1287	1/1	0.96	0.09	49,49,49,49	0
3	FE	B	1290	1/1	0.97	0.08	51,51,51,51	0
3	FE	A	1293	1/1	0.98	0.16	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.