



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:35 PM UTC

PDB ID : 4F09 / pdb\_00004f09  
Title : Discovery and Optimization of C-2 Methyl Imidazo-pyrrolopyridines as Potent and Orally Bioavailable JAK1 Inhibitors with Selectivity over JAK2  
Authors : Murray, J.M.  
Deposited on : 2012-05-03  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2508 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	290	2398	1520	415	447	2	14	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	831	GLY	-	expression tag	UNP O60674
A	832	SER	-	expression tag	UNP O60674

- Molecule 2 is 2-methyl-1-(piperidin-4-yl)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine (CCD ID: JAK) (formula: C<sub>14</sub>H<sub>17</sub>N<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
2	A	1	19	14	5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	91	Total O 91 91	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.44Å 69.49Å 50.23Å 90.00° 101.17° 90.00°	Depositor
Resolution (Å)	49.28 – 2.40 49.28 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.28-2.40) 99.7 (49.28-2.40)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.50 (at 2.39Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.11.2	Depositor
R, $R_{free}$	0.195 , 0.254 0.206 , 0.262	Depositor DCC
$R_{free}$ test set	575 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtriage
Anisotropy	0.616	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2508	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, JAK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.00	16/2415 (0.7%)	1.38	13/3247 (0.4%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	869	ASP	C-N	8.32	1.40	1.33
1	A	853	GLN	CA-C	-7.84	1.43	1.52
1	A	1019	PHE	C-O	-7.56	1.14	1.24
1	A	869	ASP	N-CA	-6.21	1.39	1.46
1	A	853	GLN	N-CA	-6.10	1.39	1.46
1	A	974	HIS	CD2-NE2	-6.01	1.31	1.37
1	A	846	GLU	C-O	-5.69	1.17	1.24
1	A	852	LEU	CA-C	5.61	1.57	1.52
1	A	854	GLN	C-O	-5.52	1.16	1.23
1	A	870	PRO	N-CD	5.44	1.55	1.47
1	A	870	PRO	CA-C	-5.44	1.47	1.52
1	A	856	GLY	C-O	-5.38	1.16	1.23
1	A	967	LEU	C-O	-5.30	1.18	1.24
1	A	1048	PHE	C-O	-5.28	1.17	1.24
1	A	855	LEU	N-CA	-5.25	1.40	1.46
1	A	853	GLN	C-O	-5.11	1.17	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	852	LEU	N-CA-C	11.69	128.15	112.13
1	A	1076	PHE	CA-CB-CG	7.76	121.56	113.80
1	A	855	LEU	N-CA-C	7.73	122.62	112.72
1	A	869	ASP	O-C-N	7.53	128.04	121.34
1	A	869	ASP	N-CA-C	5.84	119.22	108.94

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1085	ASN	N-CA-C	5.71	119.54	112.58
1	A	968	GLY	CA-C-N	5.64	128.40	120.28
1	A	968	GLY	C-N-CA	5.64	128.40	120.28
1	A	989	ARG	N-CA-C	5.63	118.66	108.69
1	A	869	ASP	CA-C-N	-5.39	114.80	120.94
1	A	869	ASP	C-N-CA	-5.39	114.80	120.94
1	A	856	GLY	N-CA-C	-5.31	100.59	113.18
1	A	952	LYS	N-CA-C	5.07	116.80	111.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2398	0	2368	46	2
2	A	19	0	17	2	0
3	A	91	0	0	0	0
All	All	2508	0	2385	47	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:858:GLY:HA2	1:A:859:ASN:CB	1.68	1.17
1:A:858:GLY:CA	1:A:859:ASN:HB3	1.84	1.07
1:A:858:GLY:HA2	1:A:859:ASN:HB3	1.03	1.01
1:A:850:LYS:HE3	1:A:869:ASP:HB3	1.64	0.80
1:A:869:ASP:OD2	1:A:874:ASN:HA	1.82	0.78
1:A:1011:LYS:HE2	1:A:1012:GLU:H	1.50	0.76
1:A:1011:LYS:HE2	1:A:1012:GLU:N	2.05	0.72
1:A:857:LYS:HG3	1:A:858:GLY:N	2.07	0.70
1:A:1007:PTR:O3P	1:A:1030:LYS:NZ	2.20	0.69

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:859:ASN:O	1:A:860:PHE:HB2	1.92	0.68
1:A:867:ARG:HG2	1:A:869:ASP:HB2	1.75	0.68
1:A:1008:PTR:HE1	1:A:1010:VAL:HB	1.75	0.68
1:A:858:GLY:HA3	1:A:861:GLY:O	1.95	0.66
1:A:1011:LYS:HA	1:A:1011:LYS:CE	2.31	0.61
1:A:858:GLY:N	1:A:861:GLY:O	2.36	0.59
1:A:918:TYR:HB3	1:A:922:ARG:HA	1.84	0.58
1:A:1011:LYS:HD2	1:A:1029:SER:OG	2.04	0.58
1:A:858:GLY:CA	1:A:861:GLY:O	2.54	0.56
1:A:948:ILE:HG23	1:A:952:LYS:HB3	1.88	0.56
1:A:858:GLY:CA	1:A:859:ASN:CB	2.47	0.55
1:A:897:ARG:O	1:A:901:ILE:HG12	2.08	0.53
1:A:902:LEU:HG	1:A:913:TYR:HB2	1.91	0.52
1:A:857:LYS:HG3	1:A:858:GLY:H	1.73	0.51
1:A:852:LEU:C	1:A:853:GLN:HG3	2.35	0.50
1:A:966:TYR:O	1:A:969:THR:HB	2.13	0.48
1:A:971:ARG:NH2	1:A:1005:LYS:O	2.47	0.47
1:A:858:GLY:HA2	1:A:859:ASN:HB2	1.79	0.47
1:A:858:GLY:HA3	1:A:861:GLY:H	1.80	0.47
1:A:973:ILE:HD11	1:A:1001:LEU:HD11	1.98	0.46
1:A:865:MET:HE3	1:A:878:VAL:HG11	1.96	0.46
1:A:1008:PTR:HD1	1:A:1010:VAL:HG12	1.98	0.45
1:A:859:ASN:O	1:A:860:PHE:CB	2.63	0.45
1:A:852:LEU:C	1:A:853:GLN:CG	2.88	0.45
1:A:1055:LYS:NZ	1:A:1092:ASP:H	2.15	0.45
1:A:1069:LYS:HD3	1:A:1073:MET:HG3	1.99	0.44
1:A:953:LEU:HD22	1:A:1047:LEU:HB3	1.99	0.44
1:A:1002:PRO:O	1:A:1003:GLN:C	2.61	0.44
1:A:942:GLN:O	1:A:945:LYS:HE3	2.20	0.42
1:A:850:LYS:CE	1:A:869:ASP:HB3	2.43	0.42
1:A:952:LYS:HE3	1:A:952:LYS:HB2	1.33	0.42
1:A:846:GLU:HG3	1:A:919:SER:HB2	2.01	0.41
1:A:1010:VAL:CG2	1:A:1012:GLU:HG3	2.50	0.41
1:A:1038:TRP:CE3	1:A:1106:TRP:HA	2.56	0.41
1:A:1083:LYS:HA	1:A:1108:ASN:HD21	1.86	0.41
2:A:1201:JAK:H1	2:A:1201:JAK:H11	1.78	0.41
1:A:856:GLY:HA3	2:A:1201:JAK:H13	2.02	0.40
1:A:1085:ASN:O	1:A:1085:ASN:CG	2.64	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:922:ARG:NH1	1:A:1072:GLN:NE2[2_555]	1.83	0.37
1:A:875:THR:CG2	1:A:1096:ASP:OD2[2_656]	1.90	0.30

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	286/302 (95%)	273 (96%)	13 (4%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	261/271 (96%)	246 (94%)	15 (6%)	<b>18</b> <b>33</b>

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	864	GLU
1	A	870	PRO
1	A	925	LEU
1	A	947	ARG
1	A	952	LYS
1	A	967	LEU
1	A	969	THR

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	977	LEU
1	A	1003	GLN
1	A	1006	GLU
1	A	1010	VAL
1	A	1011	LYS
1	A	1072	GLN
1	A	1096	ASP
1	A	1121	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	886	HIS
1	A	909	ASN
1	A	944	HIS
1	A	1112	GLN
1	A	1125	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PTR	A	1007	1	15,16,17	1.40	2 (13%)	17,22,24	1.76	5 (29%)
1	PTR	A	1008	1	15,16,17	1.50	4 (26%)	17,22,24	2.00	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1007	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1008	1	-	0/10/11/13	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1007	PTR	OH-CZ	-2.72	1.34	1.40
1	A	1008	PTR	OH-CZ	-2.68	1.34	1.40
1	A	1007	PTR	CE1-CD1	-2.51	1.34	1.38
1	A	1008	PTR	P-OH	-2.51	1.55	1.59
1	A	1008	PTR	P-O3P	-2.21	1.46	1.54
1	A	1008	PTR	P-O2P	-2.20	1.46	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1007	PTR	OH-P-O1P	-4.82	93.39	109.48
1	A	1008	PTR	OH-P-O1P	-3.88	96.53	109.48
1	A	1008	PTR	P-OH-CZ	3.82	137.48	123.88
1	A	1008	PTR	OH-CZ-CE2	2.91	127.95	119.22
1	A	1008	PTR	OH-CZ-CE1	-2.89	110.56	119.22
1	A	1008	PTR	O3P-P-O2P	2.58	117.48	107.80
1	A	1008	PTR	CB-CG-CD1	-2.44	116.37	120.90
1	A	1007	PTR	CE2-CD2-CG	-2.38	117.87	121.00
1	A	1007	PTR	P-OH-CZ	2.29	132.05	123.88
1	A	1007	PTR	O3P-P-O2P	2.14	115.84	107.80
1	A	1007	PTR	O2P-P-OH	2.01	111.27	105.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1007	PTR	1	0
1	A	1008	PTR	2	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	JAK	A	1201	-	21,22,22	1.09	1 (4%)	24,32,32	1.62	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JAK	A	1201	-	-	0/4/12/12	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	JAK	C9-C4	-3.04	1.37	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	JAK	C16-C15-N14	3.90	118.16	110.81
2	A	1201	JAK	C12-C13-N14	3.40	117.22	110.81
2	A	1201	JAK	C9-C8-C17	3.40	137.88	132.05
2	A	1201	JAK	C13-C12-C11	-2.28	106.63	110.77
2	A	1201	JAK	C9-C4-N3	-2.22	108.36	110.29
2	A	1201	JAK	C8-C9-C4	-2.21	118.72	121.19

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1201	JAK	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/302 (95%)	0.38	28 (9%) <b>13</b> <b>10</b>	8, 21, 57, 78	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	873	ASP	7.0
1	A	859	ASN	5.6
1	A	1132	GLY	5.4
1	A	858	GLY	5.2
1	A	860	PHE	4.8
1	A	875	THR	4.6
1	A	871	LEU	4.4
1	A	1131	ALA	4.2
1	A	869	ASP	3.8
1	A	887	SER	3.8
1	A	870	PRO	3.5
1	A	886	HIS	3.5
1	A	876	GLY	3.5
1	A	857	LYS	3.1
1	A	877	GLU	3.0
1	A	1013	PRO	3.0
1	A	1111	ASN	3.0
1	A	1012	GLU	2.9
1	A	947	ARG	2.7
1	A	1011	LYS	2.5
1	A	1014	GLY	2.4
1	A	861	GLY	2.3
1	A	1085	ASN	2.3
1	A	1130	MET	2.2
1	A	946	GLU	2.2
1	A	1100	MET	2.2
1	A	1003	GLN	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	987	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PTR	A	1007	16/17	0.82	0.14	20,20,20,20	0
1	PTR	A	1008	16/17	0.86	0.17	20,20,20,20	0

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	JAK	A	1201	19/19	0.93	0.08	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.