



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 02:06 PM EDT

PDB ID : 4GOA
Title : Crystal structure of jack bean urease inhibited with fluoride
Authors : Balasubramania, A.; Ponnuraj, K.
Deposited on : 2012-08-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	840	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	KCX	A	490	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	837	6338	3979	1104	1214	41	0	1	0

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ni	0	0
			2	2		

- Molecule 3 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	F	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	686	Total	O	0	0
			686	686		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Urease



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	139.49Å 139.49Å 198.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.92 – 2.20 29.92 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.92-2.20) 99.1 (29.92-2.20)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.161 , 0.210 0.164 , 0.211	Depositor DCC
R_{free} test set	2919 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtrriage
Anisotropy	0.040	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7028	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, KCX, CME, F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.09	7/6400 (0.1%)	0.96	15/8672 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	451	ARG	CZ-NH1	7.31	1.42	1.33
1	A	408	VAL	CB-CG1	5.47	1.64	1.52
1	A	786	ALA	CA-CB	5.42	1.63	1.52
1	A	142	GLU	CB-CG	-5.14	1.42	1.52
1	A	616	ALA	CA-CB	5.07	1.63	1.52
1	A	281	ASN	CB-CG	5.06	1.62	1.51
1	A	391	VAL	CB-CG1	5.04	1.63	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	451	ARG	NE-CZ-NH2	-9.22	115.69	120.30
1	A	451	ARG	NE-CZ-NH1	8.70	124.65	120.30
1	A	590	MET	CA-CB-CG	-7.67	100.27	113.30
1	A	609	ARG	NE-CZ-NH1	-7.12	116.74	120.30
1	A	783	ARG	NE-CZ-NH2	-6.97	116.82	120.30
1	A	6	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	A	276	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	A	139	CYS	CA-CB-SG	6.20	125.15	114.00
1	A	733	ASP	CB-CG-OD2	-6.04	112.87	118.30
1	A	68	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	A	314	VAL	CG1-CB-CG2	-5.31	102.40	110.90
1	A	277	LYS	CD-CE-NZ	5.25	123.78	111.70
1	A	68	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	A	185	ARG	NE-CZ-NH1	5.07	122.83	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	ASP	CB-CA-C	-5.03	100.34	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6338	0	6322	98	2
2	A	2	0	0	0	0
3	A	2	0	0	0	0
4	A	686	0	0	17	6
All	All	7028	0	6322	98	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:LYS:HE2	4:A:1459:HOH:O	1.59	1.00
1:A:63:HIS:HD2	4:A:1231:HOH:O	1.49	0.93
1:A:442:THR:OG1	1:A:490:KCX:HG2	1.69	0.92
1:A:275:HIS:HD2	1:A:277:LYS:H	1.08	0.92
1:A:666:SER:HA	4:A:1516:HOH:O	1.70	0.90
1:A:247:LYS:CE	4:A:1459:HOH:O	2.21	0.86
1:A:657:GLN:OE1	1:A:825:VAL:HG22	1.76	0.86
1:A:222:ARG:CG	1:A:222:ARG:HH11	1.90	0.84
1:A:490:KCX:HD3	1:A:490:KCX:C	2.10	0.81
1:A:66:GLY:H	1:A:69:GLN:HE21	1.29	0.78
1:A:479:HIS:ND1	1:A:512:HIS:HE1	1.84	0.75
1:A:222:ARG:HH11	1:A:222:ARG:HG2	1.50	0.75
1:A:515:GLN:HE21	1:A:516:ILE:H	1.32	0.75
1:A:275:HIS:CD2	1:A:277:LYS:H	1.99	0.75
1:A:184:ARG:H	1:A:251:HIS:HD2	1.35	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:692:GLN:H	1:A:692:GLN:HE21	1.39	0.70
1:A:490:KCX:C	1:A:490:KCX:CD	2.69	0.69
1:A:668:ASN:HD22	1:A:670:ASN:H	1.41	0.69
1:A:725:MET:HE3	1:A:760:LEU:HD11	1.74	0.69
1:A:683:ASN:HD21	1:A:799:LYS:H	1.40	0.67
1:A:753:LEU:HB3	4:A:1636:HOH:O	1.95	0.66
1:A:653:LYS:HD3	1:A:654:MET:CE	2.25	0.65
1:A:597:ARG:HD3	4:A:1039:HOH:O	1.97	0.65
1:A:69:GLN:NE2	1:A:105:GLY:H	1.95	0.64
1:A:141:ASP:OD1	1:A:275:HIS:HE1	1.81	0.63
1:A:135:GLY:HA2	1:A:292:ARG:O	1.99	0.62
1:A:66:GLY:H	1:A:69:GLN:NE2	1.97	0.60
1:A:63:HIS:CD2	4:A:1231:HOH:O	2.35	0.60
1:A:740:THR:N	1:A:741:PRO:HD3	2.17	0.59
1:A:222:ARG:HH11	1:A:222:ARG:HG3	1.68	0.58
1:A:619:ASP:HB3	1:A:654:MET:CE	2.34	0.57
1:A:471:SER:HA	1:A:491:LEU:HD23	1.88	0.55
1:A:715:THR:HG23	1:A:716:LYS:N	2.21	0.55
1:A:277:LYS:HE3	4:A:1194:HOH:O	2.07	0.55
1:A:442:THR:HG1	1:A:490:KCX:HG2	1.70	0.54
1:A:69:GLN:HE22	1:A:105:GLY:H	1.56	0.54
1:A:816:VAL:HG12	1:A:823:LEU:HD12	1.90	0.54
1:A:63:HIS:HE1	4:A:1355:HOH:O	1.90	0.53
1:A:130:ASP:N	4:A:1577:HOH:O	2.41	0.52
1:A:222:ARG:CG	1:A:222:ARG:NH1	2.58	0.52
1:A:683:ASN:HD21	1:A:799:LYS:N	2.05	0.52
1:A:715:THR:CG2	1:A:716:LYS:N	2.72	0.52
1:A:775:ASN:HB3	1:A:784:VAL:HG21	1.91	0.52
1:A:131:ASN:CB	4:A:1415:HOH:O	2.58	0.51
1:A:597:ARG:CD	4:A:1039:HOH:O	2.54	0.51
1:A:221:ILE:O	1:A:222:ARG:HG3	2.11	0.51
1:A:60:LEU:C	1:A:60:LEU:HD23	2.32	0.50
1:A:39:ILE:O	1:A:43:ILE:HG12	2.11	0.50
1:A:772:GLN:HG2	4:A:1119:HOH:O	2.12	0.49
1:A:519:HIS:CE1	1:A:551:GLY:HA3	2.48	0.49
1:A:111:LEU:HD23	1:A:118:VAL:HA	1.95	0.49
1:A:172:HIS:HE1	1:A:324:GLY:O	1.95	0.48
1:A:407:HIS:CE1	1:A:544:TYR:CD2	3.02	0.48
1:A:491:LEU:HD11	1:A:505:CYS:HB3	1.95	0.48
1:A:612:LYS:HE2	4:A:1455:HOH:O	2.13	0.47
1:A:515:GLN:NE2	1:A:516:ILE:H	2.07	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:653:LYS:HD3	1:A:654:MET:HE2	1.95	0.46
1:A:772:GLN:NE2	4:A:1119:HOH:O	2.41	0.46
1:A:683:ASN:ND2	1:A:799:LYS:H	2.11	0.46
1:A:827:GLU:H	1:A:827:GLU:HG2	1.57	0.45
1:A:490:KCX:HD3	1:A:490:KCX:O	2.15	0.45
1:A:619:ASP:HB3	1:A:654:MET:HE1	1.97	0.45
1:A:132:ARG:HE	1:A:132:ARG:HB3	1.39	0.45
1:A:632:SER:O	1:A:638:GLY:HA3	2.17	0.45
1:A:490:KCX:HG2	1:A:490:KCX:HZ	1.57	0.45
1:A:360:LYS:HB3	1:A:365:ALA:HB2	1.99	0.45
1:A:17:GLY:O	1:A:21:GLN:HG3	2.17	0.45
1:A:529:VAL:HA	1:A:553:HIS:CD2	2.52	0.44
1:A:558:ILE:HD11	1:A:621:LEU:HD12	1.99	0.44
1:A:573:PRO:HG3	1:A:637:MET:O	2.17	0.44
1:A:490:KCX:HB3	4:A:1099:HOH:O	2.16	0.44
1:A:576:PRO:HB2	1:A:653:LYS:HD2	1.99	0.44
1:A:730:ASP:O	1:A:731:ILE:HG12	2.17	0.44
1:A:82:GLN:NE2	1:A:740:THR:OG1	2.46	0.44
1:A:147:ASN:HD21	1:A:191:ARG:HH22	1.66	0.43
1:A:725:MET:CE	1:A:760:LEU:HD11	2.46	0.43
1:A:707:MET:HG3	1:A:719:MET:HG2	2.00	0.43
1:A:408:VAL:O	1:A:429:GLY:HA3	2.19	0.43
1:A:740:THR:H	1:A:741:PRO:HD3	1.82	0.43
1:A:569:SER:HB3	1:A:629:ILE:HB	2.01	0.42
1:A:268:CYS:HB3	1:A:271:ASN:ND2	2.34	0.42
1:A:593:HIS:HE1	4:A:1325:HOH:O	2.02	0.42
1:A:479:HIS:ND1	1:A:512:HIS:CE1	2.75	0.42
1:A:707:MET:CG	1:A:719:MET:HG2	2.49	0.42
1:A:715:THR:CG2	1:A:716:LYS:H	2.31	0.42
1:A:222:ARG:HG3	1:A:222:ARG:NH1	2.30	0.42
1:A:509:ALA:HB2	1:A:516:ILE:HD11	2.01	0.42
1:A:469:LYS:HE2	1:A:495:TRP:CE2	2.54	0.42
1:A:775:ASN:HD22	1:A:775:ASN:H	1.67	0.41
1:A:461:PRO:HG3	1:A:752:THR:HG22	2.01	0.41
1:A:442:THR:HG1	1:A:490:KCX:CG	2.33	0.41
1:A:668:ASN:ND2	1:A:670:ASN:H	2.13	0.41
1:A:42:GLN:NE2	1:A:42:GLN:HA	2.35	0.41
1:A:431:GLY:HA3	1:A:438:THR:HG23	2.02	0.41
1:A:565:ASN:ND2	1:A:565:ASN:H	2.19	0.40
1:A:67:ARG:NH1	1:A:103:GLU:HG2	2.36	0.40
1:A:268:CYS:HA	1:A:269:PRO:HD2	1.91	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:TYR:O	1:A:545:HIS:C	2.60	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1635:HOH:O	4:A:1638:HOH:O[2_665]	1.06	1.14
4:A:1659:HOH:O	4:A:1661:HOH:O[10_664]	1.50	0.70
4:A:1636:HOH:O	4:A:1643:HOH:O[2_665]	1.93	0.27
1:A:459:ASP:OD2	4:A:1635:HOH:O[3_565]	1.97	0.23
4:A:1661:HOH:O	4:A:1661:HOH:O[10_664]	2.07	0.13
1:A:1:MET:N	4:A:1211:HOH:O[2_665]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	829/840 (99%)	795 (96%)	33 (4%)	1 (0%)	51 60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	637	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	672/683 (98%)	646 (96%)	26 (4%)	32 41

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ARG
1	A	49	ASP
1	A	111	LEU
1	A	132	ARG
1	A	139	CYS
1	A	148	ILE
1	A	222	ARG
1	A	245	ARG
1	A	329	CYS
1	A	448	THR
1	A	474	LYS
1	A	476	ASP
1	A	508	ILE
1	A	555	PRO
1	A	597	ARG
1	A	598	GLU
1	A	652	ASP
1	A	668	ASN
1	A	692	GLN
1	A	746	MET
1	A	775	ASN
1	A	780	LEU
1	A	822	LEU
1	A	825	VAL
1	A	835	ARG
1	A	839	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	63	HIS
1	A	69	GLN
1	A	79	ASN
1	A	82	GLN
1	A	147	ASN
1	A	172	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	251	HIS
1	A	275	HIS
1	A	512	HIS
1	A	515	GLN
1	A	565	ASN
1	A	593	HIS
1	A	607	HIS
1	A	668	ASN
1	A	683	ASN
1	A	692	GLN
1	A	775	ASN
1	A	781	ASN
1	A	836	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	A	59[A]	1	8,9,10	0.80	0	6,9,11	1.79	1 (16%)
1	CME	A	592	1	8,9,10	0.72	0	6,9,11	1.59	1 (16%)
1	CME	A	207	1	8,9,10	0.41	0	6,9,11	0.78	0
1	KCX	A	490	2,1	10,11,12	0.89	0	6,12,14	1.43	1 (16%)
1	CME	A	59[B]	1	8,9,10	0.85	0	6,9,11	1.59	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	59[A]	1	-	3/5/8/10	-
1	CME	A	592	1	-	1/5/8/10	-
1	CME	A	207	1	-	1/5/8/10	-
1	KCX	A	490	2,1	-	4/9/10/12	-
1	CME	A	59[B]	1	-	1/5/8/10	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59[A]	CME	CB-SG-SD	3.83	113.77	103.86
1	A	592	CME	CE-SD-SG	-3.24	89.23	103.46
1	A	490	KCX	OQ1-CX-NZ	-3.20	120.06	124.92
1	A	59[B]	CME	CB-SG-SD	2.75	110.97	103.86

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	59[A]	CME	CE-SD-SG-CB
1	A	59[B]	CME	SD-CE-CZ-OH
1	A	490	KCX	O-C-CA-CB
1	A	490	KCX	CG-CD-CE-NZ
1	A	490	KCX	CA-CB-CG-CD
1	A	59[A]	CME	SD-CE-CZ-OH
1	A	59[A]	CME	CZ-CE-SD-SG
1	A	207	CME	CZ-CE-SD-SG
1	A	592	CME	CZ-CE-SD-SG
1	A	490	KCX	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	490	KCX	8	0

5.5 Carbohydrates

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	833/840 (99%)	-0.89	11 (1%) 77 75	3, 8, 24, 50	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	ASP	5.5
1	A	666	SER	3.0
1	A	380	PHE	2.6
1	A	598	GLU	2.5
1	A	267	ASN	2.4
1	A	266	PRO	2.3
1	A	245	ARG	2.3
1	A	131	ASN	2.3
1	A	597	ARG	2.2
1	A	824	CYS	2.2
1	A	665	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	490	12/13	0.71	0.26	6,10,12,15	0
1	CME	A	59[B]	10/11	0.96	0.12	9,10,17,21	10
1	CME	A	59[A]	10/11	0.96	0.12	12,15,38,38	10
1	CME	A	592	10/11	0.96	0.09	15,16,20,23	0
1	CME	A	207	10/11	0.98	0.07	13,13,19,19	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	F	A	903	1/1	0.98	0.15	15,15,15,15	0
3	F	A	904	1/1	0.99	0.07	11,11,11,11	0
2	NI	A	901	1/1	1.00	0.05	5,5,5,5	0
2	NI	A	902	1/1	1.00	0.07	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.