



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 05:54 PM EDT

PDB ID : 4HK0  
Title : UCA Fab (unbound) from CH65-CH67 Lineage  
Authors : Schmidt, A.G.; Harrison, S.C.  
Deposited on : 2012-10-14  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	 8% 76% 14% • 9%
1	C	237	 3% 79% 14% • 6%
2	B	214	 % 83% 14% •
2	D	214	 82% 15% ••

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6548 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UCA heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	Total 1624	C 1029	N 273	O 313	S 9	0	0	0
1	C	222	Total 1684	C 1066	N 283	O 326	S 9	0	0	0

- Molecule 2 is a protein called UCA light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	210	Total 1567	C 976	N 267	O 320	S 4	0	0	0
2	D	210	Total 1567	C 976	N 267	O 320	S 4	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total 10	O 10	0	0
3	B	27	Total 27	O 27	0	0
3	C	39	Total 39	O 39	0	0
3	D	30	Total 30	O 30	0	0



L182	E185	Q186	W187	V208	A209	P210	THR	GLU	CYS	SER
------	------	------	------	------	------	------	-----	-----	-----	-----

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.00Å 102.00Å 163.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.20 – 2.50 48.68 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.20-2.50) 97.4 (48.68-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.12 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, $R_{free}$	0.206 , 0.258 0.203 , 0.252	Depositor DCC
$R_{free}$ test set	1681 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtrriage
Anisotropy	0.494	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 33.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6548	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	1/1666 (0.1%)	0.72	1/2269 (0.0%)
1	C	0.68	0/1729	0.77	2/2357 (0.1%)
2	B	0.66	0/1606	0.77	0/2197
2	D	0.70	1/1606 (0.1%)	0.81	1/2197 (0.0%)
All	All	0.65	2/6607 (0.0%)	0.77	4/9020 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	185	GLU	CG-CD	5.68	1.60	1.51
1	A	96	CYS	CB-SG	-5.23	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	19	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	C	191	LEU	CA-CB-CG	5.71	128.44	115.30
1	A	191	LEU	CA-CB-CG	5.58	128.12	115.30
1	C	151	LEU	CA-CB-CG	5.50	127.96	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1624	0	1577	19	1
1	C	1684	0	1633	21	0
2	B	1567	0	1511	23	0
2	D	1567	0	1511	21	0
3	A	10	0	0	0	0
3	B	27	0	0	0	0
3	C	39	0	0	2	0
3	D	30	0	0	0	1
All	All	6548	0	6232	81	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:169:GLN:HE21	2:D:175:ALA:HB2	1.34	0.93
1:A:35:HIS:HD2	1:A:47:TRP:HE1	1.26	0.82
1:C:153:CYS:SG	1:C:209:CYS:CB	2.68	0.80
2:B:121:PRO:HB3	2:B:208:VAL:HG11	1.64	0.79
2:D:121:PRO:HB3	2:D:208:VAL:HG11	1.65	0.78
1:C:184:GLN:NE2	3:C:319:HOH:O	2.17	0.77
1:C:223:ARG:NH1	1:C:225:GLU:OE2	2.20	0.75
1:C:176:VAL:HG22	1:C:195:VAL:HG12	1.67	0.74
1:C:201:SER:HG	1:C:207:TYR:HH	1.28	0.74
2:B:53:ARG:CZ	2:B:59:GLU:HG2	2.24	0.68
1:A:35:HIS:CD2	1:A:47:TRP:HE1	2.12	0.67
2:D:187:TRP:HH2	2:D:208:VAL:HG12	1.60	0.67
2:D:16:GLN:O	2:D:77:VAL:HG23	1.96	0.65
2:D:140:ASP:H	2:D:169:GLN:HE22	1.42	0.65
2:D:168:LYS:NZ	2:D:172:ASN:OD1	2.17	0.65
2:D:110:GLN:HB2	2:D:142:TYR:CE1	2.32	0.64
2:B:121:PRO:HB3	2:B:208:VAL:CG1	2.29	0.62
1:A:7:SER:HB3	1:A:21:SER:OG	2.00	0.61
1:C:153:CYS:CB	1:C:209:CYS:SG	2.82	0.61
1:A:35:HIS:HE1	1:A:99:GLY:O	1.86	0.59
2:B:182:LEU:HD13	2:B:187:TRP:HB2	1.85	0.58
2:D:132:ALA:HB3	2:D:182:LEU:O	2.03	0.58
2:D:121:PRO:HB3	2:D:208:VAL:CG1	2.34	0.56
1:C:60:TYR:CE1	1:C:70:MET:HE2	2.40	0.56
1:A:11:VAL:HG11	1:A:160:PRO:HG3	1.88	0.55
1:A:136:PRO:HB3	1:A:224:VAL:HG22	1.90	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:26:ASN:H	2:D:68:ASN:ND2	2.06	0.54
2:D:110:GLN:HB2	2:D:142:TYR:CZ	2.42	0.54
2:B:123:SER:OG	2:B:126:GLU:HG3	2.06	0.54
1:C:156:LYS:NZ	3:C:319:HOH:O	2.40	0.54
1:C:47:TRP:CD1	2:D:97:VAL:HB	2.44	0.53
2:B:16:GLN:O	2:B:77:VAL:HG23	2.09	0.53
2:D:30:LYS:O	2:D:31:SER:HB3	2.09	0.53
1:A:212:ASN:HD21	1:A:214:LYS:HG3	1.74	0.53
2:B:53:ARG:NH2	2:B:59:GLU:HG2	2.24	0.52
1:C:176:VAL:HG22	1:C:195:VAL:CG1	2.39	0.52
2:D:140:ASP:H	2:D:169:GLN:NE2	2.06	0.52
1:A:167:TRP:CH2	1:A:209:CYS:HB3	2.47	0.50
1:A:40:ALA:HB3	1:A:43:GLN:HG3	1.92	0.50
2:B:182:LEU:CD1	2:B:187:TRP:HB2	2.42	0.50
1:C:37:VAL:O	1:C:37:VAL:HG23	2.11	0.49
2:D:14:PRO:HD3	2:D:108:LEU:O	2.13	0.49
2:B:38:LYS:HB3	2:B:39:PRO:HD2	1.95	0.49
2:D:34:TRP:CZ3	2:D:87:CYS:HB3	2.47	0.49
1:C:37:VAL:HG21	1:C:116:TRP:CZ3	2.48	0.48
1:A:60:TYR:CE1	1:A:70:MET:HE2	2.49	0.48
2:D:38:LYS:HB3	2:D:39:PRO:HD2	1.95	0.47
2:B:207:THR:HG22	2:B:208:VAL:N	2.30	0.46
1:C:198:PRO:O	1:C:201:SER:HB3	2.16	0.46
1:A:175:GLY:O	1:A:195:VAL:HA	2.16	0.46
2:D:31:SER:HA	2:D:50:ASP:OD1	2.15	0.46
2:B:127:LEU:HD23	2:B:132:ALA:HB2	1.99	0.45
1:A:22:CYS:O	1:A:78:THR:HA	2.17	0.45
1:A:47:TRP:CG	2:B:97:VAL:HB	2.52	0.44
2:B:151:LYS:HB2	2:B:194:SER:HB2	1.99	0.44
2:D:126:GLU:HG2	2:D:131:LYS:HB2	1.99	0.44
1:C:11:VAL:CG2	1:C:215:PRO:HB3	2.46	0.44
2:B:38:LYS:O	2:B:41:GLN:HG2	2.18	0.43
1:A:151:LEU:C	1:A:151:LEU:HD12	2.39	0.43
2:B:142:TYR:CG	2:B:143:PRO:HA	2.54	0.43
1:A:13:LYS:NZ	1:A:126:SER:O	2.51	0.43
1:A:47:TRP:CD1	2:B:97:VAL:HB	2.53	0.43
1:C:197:VAL:HG11	1:C:207:TYR:CE1	2.54	0.43
1:C:199:SER:O	1:C:202:LEU:HG	2.19	0.43
1:C:213:HIS:HB3	1:C:218:THR:OG1	2.19	0.42
1:C:167:TRP:CH2	1:C:209:CYS:HB3	2.53	0.42
2:B:122:PRO:HB3	2:B:132:ALA:HA	2.02	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:LEU:HD21	1:A:207:TYR:CG	2.55	0.42
2:D:28:GLY:HA2	2:D:65:ASN:OD1	2.20	0.41
2:D:36:GLN:HG3	2:D:85:TYR:CE2	2.54	0.41
2:B:60:ARG:HB3	2:B:75:SER:O	2.20	0.41
2:B:198:THR:HG22	2:B:203:THR:HG23	2.01	0.41
2:B:35:TYR:O	2:B:85:TYR:HA	2.19	0.41
2:B:60:ARG:NH1	2:B:81:ASP:OD2	2.42	0.41
1:C:11:VAL:HG23	1:C:215:PRO:HB3	2.02	0.41
2:B:121:PRO:CB	2:B:208:VAL:HG11	2.42	0.41
1:A:212:ASN:HD21	1:A:214:LYS:CG	2.34	0.40
1:A:2:VAL:HG13	1:A:27:TYR:CD1	2.57	0.40
1:C:87:ARG:HH11	1:C:87:ARG:HD3	1.75	0.40
2:B:82:GLU:HG3	2:B:107:VAL:HG23	2.04	0.40
1:C:157:ASP:HB3	1:C:188:LEU:HD13	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLY:O	3:D:302:HOH:O[3_455]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/237 (88%)	203 (97%)	6 (3%)	0	100	100
1	C	218/237 (92%)	211 (97%)	7 (3%)	0	100	100
2	B	208/214 (97%)	198 (95%)	8 (4%)	2 (1%)	15	28
2	D	208/214 (97%)	199 (96%)	7 (3%)	2 (1%)	15	28
All	All	843/902 (94%)	811 (96%)	28 (3%)	4 (0%)	29	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	132	ALA
2	D	31	SER
2	B	94	SER
2	D	94	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/200 (89%)	172 (97%)	6 (3%)	37	63
1	C	185/200 (92%)	178 (96%)	7 (4%)	33	58
2	B	177/181 (98%)	177 (100%)	0	100	100
2	D	177/181 (98%)	176 (99%)	1 (1%)	86	95
All	All	717/762 (94%)	703 (98%)	14 (2%)	55	79

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	SER
1	A	72	ARG
1	A	151	LEU
1	A	153	CYS
1	A	191	LEU
1	A	210	ASN
1	C	72	ARG
1	C	98	ARG
1	C	101	LEU
1	C	125	SER
1	C	140	SER
1	C	153	CYS
1	C	191	LEU
2	D	110	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	HIS
1	A	177	HIS
1	A	212	ASN
2	B	16	GLN
2	B	169	GLN
1	C	177	HIS
1	C	184	GLN
2	D	68	ASN
2	D	169	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/237 (90%)	0.52	18 (8%) 11 11	22, 52, 81, 123	0
1	C	222/237 (93%)	0.04	7 (3%) 47 51	16, 32, 64, 125	0
2	B	210/214 (98%)	-0.00	3 (1%) 75 77	21, 40, 70, 134	0
2	D	210/214 (98%)	-0.10	1 (0%) 91 91	17, 32, 57, 91	0
All	All	857/902 (95%)	0.12	29 (3%) 45 48	16, 39, 73, 134	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	185	SER	4.0
1	A	227	LYS	3.7
1	A	73	ASP	3.7
1	A	24	ALA	3.6
1	A	76	ILE	3.5
1	A	5	VAL	3.3
1	C	218	THR	3.3
1	A	220	VAL	3.1
1	A	174	SER	3.1
2	D	1	GLN	3.0
1	A	1	GLN	2.9
1	A	217	ASN	2.8
1	C	220	VAL	2.7
1	A	3	GLN	2.7
1	C	108	TYR	2.7
1	A	56	GLY	2.5
1	A	26	GLY	2.5
1	A	77	SER	2.5
1	A	54	ASN	2.5
2	B	1	GLN	2.3
1	A	9	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	222	LYS	2.2
1	A	71	THR	2.2
1	A	223	ARG	2.2
1	C	202	LEU	2.2
1	A	208	ILE	2.1
2	B	59	GLU	2.1
1	C	186	SER	2.1
2	B	131	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.