



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2024 – 08:11 AM EST

PDB ID : 4K96
Title : Structure of Binary Complex of cGAS with Bound dsDNA
Authors : Gao, P.; Wu, Y.; Patel, D.J.
Deposited on : 2013-04-19
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div>
1	B	362	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: orange;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div>
2	E	17	<div style="display: flex; align-items: center;"> <div style="width: 18%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 71%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 29%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div>
2	G	17	<div style="display: flex; align-items: center;"> <div style="width: 24%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 71%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 29%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div>
3	F	17	<div style="display: flex; align-items: center;"> <div style="width: 18%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 65%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 18%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 18%; height: 10px; background-color: orange; margin-right: 5px;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	17	 <p>6% 71% 24% 6%</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7981 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic GMP-AMP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	355	Total	C	N	O	S	0	0	0
			2927	1883	495	536	13			
1	B	358	Total	C	N	O	S	0	0	0
			2948	1895	498	542	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	SER	-	expression tag	UNP Q8C6L5
B	146	SER	-	expression tag	UNP Q8C6L5

- Molecule 2 is a DNA chain called DNA-F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	E	17	Total	C	N	O	P	0	0	0
			350	167	73	94	16			
2	G	17	Total	C	N	O	P	0	0	0
			350	167	73	94	16			

- Molecule 3 is a DNA chain called DNA-R.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	F	17	Total	C	N	O	P	0	0	0
			341	166	53	106	16			
3	H	17	Total	C	N	O	P	0	0	0
			341	166	53	106	16			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	264	Total	O	0	0
			264	264		
5	B	329	Total	O	0	0
			329	329		
5	E	37	Total	O	0	0
			37	37		
5	F	27	Total	O	0	0
			27	27		
5	G	41	Total	O	0	0
			41	41		
5	H	24	Total	O	0	0
			24	24		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclic GMP-AMP synthase




- Molecule 1: Cyclic GMP-AMP synthase




- Molecule 2: DNA-F




- Molecule 2: DNA-F



- Molecule 3: DNA-R



- Molecule 3: DNA-R



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.91Å 93.80Å 75.51Å 90.00° 97.72° 90.00°	Depositor
Resolution (Å)	45.07 – 2.08 45.07 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.07-2.08) 94.6 (45.07-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.08Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.202 , 0.226 0.203 , 0.226	Depositor DCC
R_{free} test set	2000 reflections (2.69%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtrriage
Anisotropy	0.401	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7981	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	1/2990 (0.0%)	0.44	1/4013 (0.0%)
1	B	0.41	2/3012 (0.1%)	0.44	1/4044 (0.0%)
2	E	0.49	0/395	1.14	0/608
2	G	0.43	0/395	1.02	0/608
3	F	0.54	0/379	1.71	16/583 (2.7%)
3	H	0.53	0/379	1.20	1/583 (0.2%)
All	All	0.43	3/7550 (0.0%)	0.73	19/10439 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	193	GLU	CD-OE2	-5.93	1.19	1.25
1	A	361	GLU	CD-OE1	-5.29	1.19	1.25
1	B	193	GLU	CD-OE1	-5.24	1.19	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	DT	OP2-P-O3'	-10.72	81.62	105.20
3	F	1	DT	OP1-P-O3'	-10.67	81.73	105.20
3	F	2	DT	C1'-O4'-C4'	-8.83	101.27	110.10
3	F	2	DT	OP1-P-OP2	7.71	131.16	119.60
3	F	3	DT	C3'-C2'-C1'	-6.97	94.14	102.50
3	F	3	DT	C5-C4-O4	-6.68	120.23	124.90
3	H	1	DT	O4'-C1'-N1	6.66	112.66	108.00
3	F	3	DT	N3-C4-O4	6.51	123.81	119.90
3	F	3	DT	C4'-C3'-C2'	-6.36	97.38	103.10
3	F	2	DT	C5-C4-O4	-5.70	120.91	124.90
1	B	146	SER	C-N-CD	5.59	140.14	128.40
3	F	3	DT	O4'-C1'-C2'	-5.58	101.43	105.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	DT	O4'-C4'-C3'	-5.49	102.31	104.50
3	F	1	DT	C1'-O4'-C4'	-5.46	104.64	110.10
1	A	343	GLU	C-N-CD	5.29	139.51	128.40
3	F	1	DT	C4-C5-C7	5.25	122.15	119.00
3	F	3	DT	O4'-C1'-N1	5.09	111.56	108.00
3	F	2	DT	N3-C4-O4	5.09	122.95	119.90
3	F	9	DT	N3-C4-O4	5.05	122.93	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2927	0	2963	24	0
1	B	2948	0	2982	19	0
2	E	350	0	191	5	0
2	G	350	0	191	7	0
3	F	341	0	197	5	0
3	H	341	0	197	6	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	264	0	0	6	0
5	B	329	0	0	5	0
5	E	37	0	0	0	0
5	F	27	0	0	0	0
5	G	41	0	0	2	0
5	H	24	0	0	0	0
All	All	7981	0	6721	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:15:DG:N2	3:F:4:DC:N3	2.22	0.84
2:E:15:DG:H1	3:F:4:DC:H42	1.27	0.80
1:B:180:ARG:NH1	5:B:890:HOH:O	2.20	0.74
1:A:180:ARG:NH2	5:A:870:HOH:O	2.20	0.73
1:B:160:LYS:NZ	5:B:987:HOH:O	2.23	0.72
1:B:183:GLN:NE2	1:B:190:LYS:O	2.25	0.70
2:G:15:DG:H1	3:H:4:DC:H42	1.42	0.66
1:B:481:ILE:O	1:B:486:LYS:NZ	2.32	0.63
1:A:341:ARG:NH1	5:A:756:HOH:O	2.34	0.59
1:B:341:ARG:NH1	5:B:734:HOH:O	2.24	0.59
2:G:8:DC:H5	5:G:140:HOH:O	1.85	0.58
1:A:183:GLN:NE2	1:A:190:LYS:O	2.38	0.57
1:B:182:MET:HE2	1:B:189:PHE:HB2	1.88	0.55
1:A:422:HIS:HD2	1:A:475:LEU:HD23	1.72	0.54
1:A:448:CYS:HA	1:A:451:LYS:HE2	1.89	0.54
2:E:16:DA:C6	2:E:17:DA:N6	2.75	0.54
1:A:293:ALA:HB2	1:A:309:ILE:HG12	1.90	0.53
2:G:17:DA:N6	3:H:2:DT:H3	2.06	0.53
1:A:151:LYS:NZ	5:A:949:HOH:O	2.37	0.52
1:A:161:ARG:HH12	3:F:8:DT:H1'	1.75	0.52
2:G:14:DC:H2''	2:G:15:DG:C8	2.45	0.51
3:F:2:DT:H2'	3:F:3:DT:C6	2.45	0.51
1:B:182:MET:HG2	1:B:192:VAL:HG21	1.92	0.51
1:B:448:CYS:HA	1:B:451:LYS:HE2	1.93	0.50
1:A:342:ARG:HH12	3:H:13:DC:H1'	1.76	0.50
1:A:272:GLU:HG3	5:A:929:HOH:O	2.13	0.48
1:B:163:ASP:OD1	5:B:816:HOH:O	2.20	0.48
1:A:220:VAL:HG21	1:A:312:LEU:HB3	1.96	0.47
3:F:1:DT:H2'	3:F:2:DT:H4'	1.96	0.47
2:E:3:DA:H2''	2:E:4:DT:H72	1.97	0.46
1:A:338:THR:O	1:A:342:ARG:HG3	2.16	0.46
1:B:460:LEU:HD13	1:B:476:PHE:CE2	2.51	0.46
1:B:293:ALA:HB2	1:B:309:ILE:HG12	1.98	0.46
1:A:436:GLN:HB2	1:A:439:GLN:HG3	1.97	0.45
1:B:182:MET:HE3	1:B:269:ILE:HG21	1.98	0.45
3:H:1:DT:H2''	3:H:2:DT:O5'	2.16	0.45
1:A:492:LYS:HD3	1:A:492:LYS:HA	1.82	0.45
2:G:15:DG:H1	3:H:4:DC:N4	2.10	0.45
2:G:2:DA:OP2	5:G:137:HOH:O	2.21	0.45
1:B:148:ASP:O	1:B:152:LYS:HD3	2.17	0.44
1:A:253:GLU:O	5:A:850:HOH:O	2.21	0.43
1:B:239:PHE:HE2	1:B:257:LEU:HD13	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:271:LYS:HG2	1:B:284:VAL:HG21	1.99	0.43
1:A:292:PRO:HB3	1:A:349:PRO:O	2.17	0.43
1:A:315:LYS:HB3	3:H:15:DA:H5''	2.00	0.43
1:B:479:GLU:OE1	1:B:479:GLU:N	2.43	0.43
2:E:16:DA:C5	2:E:17:DA:N6	2.88	0.42
1:A:182:MET:HG2	1:A:192:VAL:HG21	2.02	0.42
1:A:409:LYS:HE2	1:A:409:LYS:HB3	1.84	0.42
1:A:464:LYS:NZ	5:A:913:HOH:O	2.49	0.42
1:B:223:ILE:HD11	1:B:237:VAL:HG13	2.02	0.41
1:B:229:TYR:HB3	1:B:230:GLU:H	1.68	0.41
1:A:463:GLU:HG2	1:A:486:LYS:HD2	2.01	0.41
1:B:433:GLN:OE1	5:B:810:HOH:O	2.22	0.41
1:A:465:LEU:HB2	1:A:476:PHE:CD2	2.56	0.41
2:G:15:DG:H8	2:G:15:DG:H5''	1.86	0.40
1:A:170:THR:HG23	1:A:280:ILE:HD13	2.02	0.40
1:A:244:ARG:O	1:A:246:ASN:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/362 (97%)	340 (97%)	10 (3%)	1 (0%)	41	39
1	B	354/362 (98%)	346 (98%)	7 (2%)	1 (0%)	41	39
All	All	705/724 (97%)	686 (97%)	17 (2%)	2 (0%)	41	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	ASN
1	A	300	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/335 (98%)	314 (96%)	13 (4%)	31	31
1	B	330/335 (98%)	315 (96%)	15 (4%)	27	26
All	All	657/670 (98%)	629 (96%)	28 (4%)	29	28

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	LYS
1	A	185	ARG
1	A	218	LEU
1	A	219	GLU
1	A	275	LYS
1	A	297	LEU
1	A	302	GLU
1	A	350	LYS
1	A	430	MET
1	A	433	GLN
1	A	452	LEU
1	A	460	LEU
1	A	465	LEU
1	B	147	PRO
1	B	149	LYS
1	B	159	LEU
1	B	178	LEU
1	B	183	GLN
1	B	195	LEU
1	B	218	LEU
1	B	297	LEU
1	B	406	GLU
1	B	414	GLU
1	B	430	MET
1	B	452	LEU
1	B	460	LEU
1	B	465	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	478	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	183	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/362 (98%)	0.23	18 (5%) 28 32	20, 41, 77, 117	0
1	B	358/362 (98%)	0.12	14 (3%) 39 44	20, 33, 65, 110	0
2	E	17/17 (100%)	0.27	3 (17%) 1 1	24, 41, 119, 120	0
2	G	17/17 (100%)	0.45	4 (23%) 0 0	25, 53, 131, 159	0
3	F	17/17 (100%)	0.69	3 (17%) 1 1	29, 51, 157, 179	0
3	H	17/17 (100%)	0.30	1 (5%) 22 26	31, 59, 127, 131	0
All	All	781/792 (98%)	0.20	43 (5%) 25 30	20, 38, 81, 179	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	245	GLY	9.4
1	A	229	TYR	7.4
1	A	244	ARG	6.6
1	B	185	ARG	6.6
3	F	1	DT	5.6
1	B	243	PRO	5.3
1	A	185	ARG	5.1
1	B	229	TYR	4.7
1	A	222	ARG	4.6
2	G	17	DA	4.0
1	A	246	ASN	4.0
1	A	243	PRO	4.0
1	A	240	LYS	3.9
1	A	239	PHE	3.7
1	B	239	PHE	3.6
3	F	2	DT	3.6
1	A	417	ALA	3.6
1	B	244	ARG	3.6
1	A	184	LYS	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	255	GLU	3.5
2	G	16	DA	3.3
2	E	17	DA	3.2
1	A	228	TYR	3.2
1	B	246	ASN	3.2
1	B	245	GLY	3.1
1	B	223	ILE	3.0
1	A	250	HIS	2.9
1	B	222	ARG	2.9
1	B	186	GLU	2.7
2	G	1	DA	2.7
2	E	16	DA	2.7
1	B	247	PRO	2.5
1	A	186	GLU	2.4
2	G	15	DG	2.3
3	F	3	DT	2.3
1	A	197	THR	2.3
1	A	248	LEU	2.2
1	B	413	GLN	2.2
1	A	414	GLU	2.2
1	A	255	GLU	2.1
3	H	3	DT	2.1
2	E	15	DG	2.1
1	B	392	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ZN	B	601	1/1	0.96	0.30	88,88,88,88	0
4	ZN	A	601	1/1	1.00	0.10	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.