



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 08:07 PM EDT

PDB ID : 4LB9  
Title : X-ray study of human serum albumin complexed with etoposide  
Authors : Wang, Z.; Ho, J.X.; Ruble, J.; Rose, J.P.; Carter, D.C.  
Deposited on : 2013-06-20  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERUM ALBUMIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	583	4635	2926	784	884	41	0	0	0


- Molecule 2 is MYRISTIC ACID (three-letter code: MYR) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	16	14	2	0	0
2	A	1	16	14	2	0	0
2	A	1	16	14	2	0	0
2	A	1	16	14	2	0	0

- Molecule 3 is (5S,5aR,8aR,9R)-9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxo-5,5a,6,8,8a,9-hexahydrofuro[3',4':6,7]naphtho[2,3-d][1,3]dioxol-5-yl 4,6-O-[(1R)-ethylidene]-beta-D-glucopyra

noside (three-letter code: EVP) (formula:  $C_{29}H_{32}O_{13}$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	C	O	0	0
			42	29	13		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total	O	0	0
			68	68		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SERUM ALBUMIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.54Å 38.68Å 99.22Å 90.00° 104.56° 90.00°	Depositor
Resolution (Å)	25.68 – 2.70 25.68 – 2.70	Depositor EDS
% Data completeness (in resolution range)	86.6 (25.68-2.70) 81.0 (25.68-2.70)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.72Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.195 , 0.249 0.197 , 0.253	Depositor DCC
$R_{free}$ test set	765 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtrriage
Anisotropy	0.533	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 31.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4809	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EVP, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/4725	0.47	0/6373

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4635	0	4557	74	0
2	A	64	0	108	10	0
3	A	42	0	32	2	0
4	A	68	0	0	16	0
All	All	4809	0	4697	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:519:LYS:NZ	4:A:762:HOH:O	2.18	0.76
1:A:212:LYS:NZ	4:A:732:HOH:O	2.20	0.75
1:A:254:ALA:HB1	2:A:601:MYR:H42	1.75	0.69
1:A:369:CYS:O	4:A:716:HOH:O	2.09	0.68
1:A:541:LYS:NZ	4:A:755:HOH:O	2.27	0.68
1:A:348:ARG:NH2	1:A:482:VAL:O	2.27	0.67
1:A:294:GLU:OE1	4:A:734:HOH:O	2.11	0.67
3:A:605:EVP:H8A	3:A:605:EVP:O13	1.95	0.67
1:A:182:LEU:O	1:A:186:ARG:HG2	1.97	0.64
1:A:160:ARG:NH1	1:A:188:GLU:OE1	2.31	0.64
1:A:167:GLU:OE1	4:A:740:HOH:O	2.15	0.63
1:A:160:ARG:NH2	4:A:715:HOH:O	2.28	0.63
1:A:66:LEU:HD21	2:A:601:MYR:H122	1.80	0.62
1:A:314:ASP:HB2	1:A:317:LYS:HB3	1.81	0.62
1:A:22:LEU:HD13	2:A:601:MYR:H82	1.81	0.61
1:A:561:ALA:O	1:A:564:LYS:NZ	2.34	0.61
1:A:206:PHE:O	4:A:741:HOH:O	2.16	0.61
1:A:276:LYS:NZ	1:A:280:GLU:OE2	2.34	0.61
1:A:233:LYS:NZ	1:A:237:ASP:OD1	2.35	0.60
1:A:460:LEU:HD21	2:A:603:MYR:H132	1.85	0.59
1:A:299:PRO:HB2	1:A:302:LEU:HD21	1.86	0.58
1:A:34:CYS:N	1:A:84:TYR:OH	2.38	0.57
1:A:479:GLU:CG	1:A:483:ASN:HB3	2.34	0.57
1:A:198:LEU:HB2	1:A:458:ASN:HD22	1.69	0.56
1:A:333:GLU:OE1	1:A:337:ARG:NH1	2.38	0.56
1:A:383:GLU:OE1	1:A:485:ARG:NH1	2.39	0.56
1:A:136:LYS:NZ	4:A:728:HOH:O	2.21	0.55
1:A:411:TYR:HB3	2:A:603:MYR:H111	1.89	0.55
1:A:66:LEU:HA	1:A:69:LEU:HD12	1.88	0.54
1:A:516:LEU:O	1:A:521:ARG:NH1	2.40	0.53
1:A:61:ASN:HB2	1:A:64:LYS:HD2	1.91	0.53
1:A:60:GLU:HG2	1:A:61:ASN:H	1.74	0.52
1:A:479:GLU:HG2	1:A:483:ASN:HB3	1.92	0.52
1:A:414:LYS:NZ	4:A:759:HOH:O	2.34	0.51
1:A:170:GLN:OE1	4:A:721:HOH:O	2.19	0.51
1:A:82:GLU:N	1:A:82:GLU:OE1	2.43	0.51
1:A:415:VAL:HG21	2:A:603:MYR:H143	1.93	0.50
1:A:197:ARG:HD3	3:A:605:EVP:H24	1.94	0.50
1:A:149:PHE:CE2	1:A:193:SER:HB2	2.47	0.49
1:A:479:GLU:HG3	1:A:483:ASN:HB3	1.95	0.49
1:A:551:PHE:CD2	2:A:604:MYR:H52	2.48	0.49
1:A:429:ASN:ND2	4:A:747:HOH:O	2.46	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:GLU:H	1:A:37:GLU:CD	2.16	0.48
1:A:408:LEU:HD23	1:A:529:LEU:HD23	1.95	0.48
1:A:479:GLU:HG3	1:A:480:SER:H	1.79	0.48
1:A:351:LYS:NZ	4:A:767:HOH:O	2.38	0.48
1:A:389:LYS:NZ	1:A:442:GLU:OE2	2.47	0.47
1:A:61:ASN:OD1	1:A:61:ASN:N	2.47	0.47
1:A:430:LEU:HB3	2:A:602:MYR:H141	1.96	0.47
1:A:472:ARG:H	1:A:472:ARG:HG2	1.50	0.47
1:A:60:GLU:HG2	1:A:61:ASN:OD1	2.15	0.46
1:A:471:ASP:OD1	1:A:471:ASP:N	2.47	0.46
1:A:379:PRO:HA	1:A:382:GLU:HB2	1.97	0.46
1:A:388:ILE:HG22	2:A:602:MYR:H72	1.98	0.46
1:A:262:LYS:HE2	1:A:266:GLU:OE2	2.16	0.45
1:A:383:GLU:HB3	1:A:384:PRO:HD3	1.98	0.45
1:A:237:ASP:O	1:A:241:VAL:HG13	2.16	0.45
1:A:305:LEU:HD21	1:A:337:ARG:CZ	2.46	0.45
1:A:479:GLU:HG3	1:A:480:SER:N	2.32	0.45
1:A:519:LYS:O	1:A:523:ILE:HG12	2.17	0.44
1:A:416:PRO:O	1:A:534:LYS:HE2	2.18	0.44
1:A:428:ARG:NH1	4:A:761:HOH:O	2.50	0.44
1:A:20:LYS:NZ	4:A:756:HOH:O	2.50	0.43
1:A:507:PHE:CD2	2:A:604:MYR:H121	2.53	0.43
1:A:59:ALA:HB1	1:A:60:GLU:OE1	2.19	0.43
1:A:218:ARG:NH1	4:A:707:HOH:O	2.37	0.42
1:A:151:ALA:HB3	1:A:152:PRO:HD3	2.02	0.42
1:A:240:LYS:HE2	1:A:244:GLU:OE1	2.19	0.42
1:A:103:LEU:HD21	1:A:249:ASP:HB2	2.01	0.42
1:A:219:LEU:HD23	1:A:219:LEU:HA	1.75	0.42
1:A:218:ARG:HA	1:A:218:ARG:HD3	1.88	0.42
1:A:30:TYR:HE1	1:A:102:PHE:HB3	1.85	0.41
1:A:398:LEU:O	1:A:402:LYS:HB2	2.19	0.41
1:A:30:TYR:CE1	1:A:102:PHE:HB3	2.55	0.41
1:A:38:ASP:O	1:A:42:LEU:HG	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	581/585 (99%)	547 (94%)	26 (4%)	8 (1%)	9 24

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	PRO
1	A	321	GLU
1	A	442	GLU
1	A	60	GLU
1	A	130	ASN
1	A	59	ALA
1	A	78	ALA
1	A	301	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	509/511 (100%)	469 (92%)	40 (8%)	10 25

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	9	HIS
1	A	58	SER

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	80	LEU
1	A	95	GLU
1	A	97	GLU
1	A	103	LEU
1	A	107	ASP
1	A	114	ARG
1	A	178	LEU
1	A	179	LEU
1	A	199	LYS
1	A	205	LYS
1	A	247	HIS
1	A	270	SER
1	A	297	GLU
1	A	301	ASP
1	A	334	TYR
1	A	337	ARG
1	A	344	VAL
1	A	345	LEU
1	A	370	TYR
1	A	378	LYS
1	A	388	ILE
1	A	389	LYS
1	A	396	GLU
1	A	407	LEU
1	A	417	GLN
1	A	433	VAL
1	A	436	LYS
1	A	451	ASP
1	A	459	GLN
1	A	472	ARG
1	A	480	SER
1	A	483	ASN
1	A	484	ARG
1	A	485	ARG
1	A	495	GLU
1	A	514	CYS
1	A	544	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	458	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EVP	A	605	-	48,48,48	3.83	18 (37%)	71,73,73	2.93	33 (46%)
2	MYR	A	603	-	15,15,15	0.56	0	15,15,15	1.09	1 (6%)
2	MYR	A	601	-	15,15,15	0.52	0	15,15,15	1.04	1 (6%)
2	MYR	A	604	-	15,15,15	0.53	0	15,15,15	1.09	1 (6%)
2	MYR	A	602	-	15,15,15	0.59	0	15,15,15	0.94	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EVP	A	605	-	-	5/12/76/76	0/7/7/7
2	MYR	A	603	-	-	5/13/13/13	-
2	MYR	A	601	-	-	6/13/13/13	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MYR	A	604	-	-	2/13/13/13	-
2	MYR	A	602	-	-	2/13/13/13	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	605	EVP	C19-C14	11.17	1.55	1.39
3	A	605	EVP	O3-C7	10.41	1.57	1.35
3	A	605	EVP	C16-C17	-9.91	1.26	1.40
3	A	605	EVP	C18-C17	-8.35	1.29	1.40
3	A	605	EVP	O3-C8	7.44	1.60	1.45
3	A	605	EVP	C15-C14	6.29	1.48	1.39
3	A	605	EVP	C9-C10	-5.48	1.44	1.53
3	A	605	EVP	C11-C4	4.90	1.49	1.40
3	A	605	EVP	O7-C18	4.61	1.44	1.37
3	A	605	EVP	C11-C10	4.03	1.58	1.50
3	A	605	EVP	C19-C18	4.02	1.45	1.38
3	A	605	EVP	C14-C5	3.84	1.57	1.52
3	A	605	EVP	C4-C5	3.74	1.57	1.51
3	A	605	EVP	C6-C7	-3.14	1.47	1.51
3	A	605	EVP	C15-C16	-2.81	1.34	1.38
3	A	605	EVP	O5-C16	2.63	1.41	1.37
3	A	605	EVP	O6-C17	2.39	1.42	1.36
3	A	605	EVP	C12-C11	-2.37	1.35	1.39

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	605	EVP	C11-C10-C9	-9.92	96.74	111.35
3	A	605	EVP	O5-C16-C17	7.27	122.14	114.53
3	A	605	EVP	C19-C14-C15	-6.35	109.44	118.09
3	A	605	EVP	C19-C18-C17	5.62	125.93	120.59
3	A	605	EVP	C24-O10-C25	5.22	116.78	111.67
3	A	605	EVP	O8-C22-C28	5.20	120.89	108.09
3	A	605	EVP	C8-O3-C7	-4.99	105.35	110.31
3	A	605	EVP	C8-C9-C6	4.92	109.37	101.73
3	A	605	EVP	O5-C16-C15	-4.68	116.02	124.08
3	A	605	EVP	C22-C28-C27	-4.14	101.30	110.01
3	A	605	EVP	C16-C15-C14	4.06	126.12	119.86
3	A	605	EVP	O3-C7-C6	-4.05	103.47	109.50
3	A	605	EVP	C19-C14-C5	3.92	127.89	120.41
3	A	605	EVP	O7-C18-C19	-3.54	117.99	124.08

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	605	EVP	C18-C19-C14	-3.39	114.63	119.86
3	A	605	EVP	C20-O5-C16	3.35	122.43	117.51
3	A	605	EVP	C28-C27-C26	-3.29	102.22	109.68
3	A	605	EVP	O2-C2-C3	3.10	131.98	127.86
3	A	605	EVP	C14-C5-C4	2.96	117.51	112.95
3	A	605	EVP	C21-O7-C18	-2.78	113.44	117.51
3	A	605	EVP	O4-C7-C6	2.70	132.82	129.38
3	A	605	EVP	C11-C4-C5	2.70	121.41	114.39
3	A	605	EVP	O13-C28-C22	2.66	116.42	110.08
3	A	605	EVP	O13-C28-C27	-2.65	104.14	110.38
3	A	605	EVP	C24-C23-C26	-2.61	105.50	109.40
3	A	605	EVP	O11-C26-C27	2.53	114.15	109.72
3	A	605	EVP	C22-O9-C23	2.51	118.62	113.72
3	A	605	EVP	O1-C13-C12	2.49	131.16	127.86
3	A	605	EVP	C18-C17-C16	2.47	122.06	118.79
2	A	604	MYR	C3-C2-C1	-2.31	108.48	114.51
3	A	605	EVP	O2-C1-O1	-2.29	104.50	108.09
3	A	605	EVP	O11-C25-O10	-2.28	108.56	110.86
2	A	603	MYR	O2-C1-C2	2.18	120.88	114.00
3	A	605	EVP	O3-C7-O4	2.14	123.71	121.43
3	A	605	EVP	O12-C27-C26	2.14	115.41	109.94
2	A	602	MYR	O2-C1-C2	2.13	120.72	114.00
2	A	601	MYR	C3-C2-C1	-2.13	108.96	114.51

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	605	EVP	C28-C22-O8-C10
3	A	605	EVP	C19-C18-O7-C21
3	A	605	EVP	C15-C16-O5-C20
2	A	603	MYR	C1-C2-C3-C4
3	A	605	EVP	C17-C18-O7-C21
2	A	603	MYR	C9-C10-C11-C12
3	A	605	EVP	C17-C16-O5-C20
2	A	604	MYR	C1-C2-C3-C4
2	A	601	MYR	C4-C5-C6-C7
2	A	604	MYR	C9-C10-C11-C12
2	A	601	MYR	C5-C6-C7-C8
2	A	603	MYR	C11-C12-C13-C14
2	A	603	MYR	C5-C6-C7-C8
2	A	601	MYR	C3-C4-C5-C6

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	601	MYR	C6-C7-C8-C9
2	A	603	MYR	C10-C11-C12-C13
2	A	601	MYR	O2-C1-C2-C3
2	A	601	MYR	O1-C1-C2-C3
2	A	602	MYR	C5-C6-C7-C8
2	A	602	MYR	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	605	EVP	2	0
2	A	603	MYR	3	0
2	A	601	MYR	3	0
2	A	604	MYR	2	0
2	A	602	MYR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	583/585 (99%)	-0.67	0 <b>100</b> <b>100</b>	16, 31, 51, 80	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EVP	A	605	42/42	0.84	0.11	25,39,49,63	0
2	MYR	A	603	16/16	0.89	0.11	18,27,38,40	0
2	MYR	A	601	16/16	0.89	0.09	21,25,31,32	0
2	MYR	A	604	16/16	0.95	0.08	21,28,34,41	0
2	MYR	A	602	16/16	0.95	0.07	17,23,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.