



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 11:48 AM EST

PDB ID : 4NL4
Title : PriA Helicase Bound to ADP
Authors : Bhattacharyya, B.; George, N.P.; Keck, J.L.
Deposited on : 2013-11-13
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	747	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5547 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Primosome assembly protein PriA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	698	5490	3486	1006	981	17	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-15	HIS	-	expression tag	UNP A6TGC5
H	-14	HIS	-	expression tag	UNP A6TGC5
H	-13	HIS	-	expression tag	UNP A6TGC5
H	-12	HIS	-	expression tag	UNP A6TGC5
H	-11	HIS	-	expression tag	UNP A6TGC5
H	-10	HIS	-	expression tag	UNP A6TGC5
H	-9	SER	-	expression tag	UNP A6TGC5
H	-8	SER	-	expression tag	UNP A6TGC5
H	-7	GLY	-	expression tag	UNP A6TGC5
H	-6	LEU	-	expression tag	UNP A6TGC5
H	-5	VAL	-	expression tag	UNP A6TGC5
H	-4	PRO	-	expression tag	UNP A6TGC5
H	-3	ARG	-	expression tag	UNP A6TGC5
H	-2	GLY	-	expression tag	UNP A6TGC5
H	-1	SER	-	expression tag	UNP A6TGC5
H	0	HIS	-	expression tag	UNP A6TGC5

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	H	1	27	10	5	10	2	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	H	2	2	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	H	28	28	28	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Primosome assembly protein PriA



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	85.43Å 85.43Å 111.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.14 – 2.65 35.12 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.14-2.65) 100.0 (35.12-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 2.65Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.218 , 0.254 0.217 , 0.251	Depositor DCC
R_{free} test set	1342 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	64.4	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.005 for -h,-k,l 0.035 for h,-h-k,-l 0.018 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5547	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.29	0/5624	0.50	0/7646

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	5490	0	5484	45	0
2	H	27	0	12	0	0
3	H	2	0	0	0	0
4	H	28	0	0	2	0
All	All	5547	0	5496	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:303:ARG:HH11	1:H:328:GLN:HE21	1.09	0.97
1:H:303:ARG:NH1	1:H:328:GLN:HE21	1.74	0.85
1:H:367:ARG:HH11	1:H:367:ARG:HB2	1.42	0.82
1:H:663:PRO:O	1:H:664:LEU:HB2	1.80	0.82
1:H:644:ASN:O	1:H:646:GLN:N	2.13	0.80
1:H:667:GLU:O	1:H:667:GLU:HG3	1.83	0.76
1:H:644:ASN:OD1	1:H:647:GLN:HG2	1.86	0.76
1:H:367:ARG:HH11	1:H:367:ARG:CB	2.05	0.70
1:H:644:ASN:C	1:H:646:GLN:H	1.95	0.69
1:H:642:ASP:CB	1:H:648:ALA:HB2	2.23	0.68
1:H:305:SER:O	1:H:308:THR:CG2	2.43	0.67
1:H:642:ASP:HB2	1:H:648:ALA:HB2	1.77	0.65
1:H:122:GLN:O	1:H:126:VAL:HG23	2.02	0.59
1:H:366:VAL:HG11	1:H:373:GLN:HE21	1.68	0.59
1:H:231:GLU:OE2	1:H:267:ARG:NH2	2.36	0.58
1:H:14:ARG:HG3	4:H:926:HOH:O	2.03	0.57
1:H:305:SER:O	1:H:308:THR:HG22	2.05	0.57
1:H:321:GLU:HG3	1:H:353:LEU:HB3	1.87	0.57
1:H:667:GLU:O	1:H:667:GLU:CG	2.53	0.56
1:H:411:ARG:NH2	1:H:500:LEU:O	2.37	0.56
1:H:190:SER:C	1:H:192:ALA:H	2.11	0.53
1:H:292:LYS:HE2	4:H:924:HOH:O	2.07	0.53
1:H:17:ASP:HB2	1:H:54:LEU:HD12	1.91	0.53
1:H:641:GLU:HG2	1:H:687:ARG:HG2	1.90	0.53
1:H:420:VAL:HG22	1:H:552:LEU:HB3	1.89	0.52
1:H:305:SER:O	1:H:308:THR:HG23	2.09	0.52
1:H:646:GLN:HB3	1:H:647:GLN:HE21	1.76	0.51
1:H:646:GLN:O	1:H:648:ALA:N	2.37	0.51
1:H:31:ARG:NH2	1:H:107:LYS:O	2.43	0.50
1:H:644:ASN:C	1:H:646:GLN:N	2.62	0.50
1:H:642:ASP:HB3	1:H:648:ALA:HB2	1.97	0.47
1:H:666:ASP:O	1:H:667:GLU:HG2	2.15	0.46
1:H:494:GLU:HA	1:H:497:LEU:HB2	1.97	0.46
1:H:131:LEU:C	1:H:133:ARG:H	2.20	0.43
1:H:168:ARG:HA	1:H:173:ALA:O	2.19	0.43
1:H:292:LYS:HB2	1:H:309:PRO:HG2	2.01	0.43
1:H:672:LEU:HB2	1:H:691:LEU:HB3	2.00	0.43
1:H:638:ILE:HD12	1:H:690:ILE:HD11	2.01	0.42
1:H:620:ALA:O	1:H:624:ARG:HG2	2.20	0.42
1:H:541:LEU:HD22	1:H:547:PHE:HE2	1.83	0.42
1:H:638:ILE:HG12	1:H:724:LEU:HD22	2.00	0.42
1:H:194:GLU:H	1:H:194:GLU:HG2	1.69	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:704:VAL:HA	1:H:707:THR:HG22	2.02	0.41
1:H:199:ASN:ND2	1:H:202:GLN:H	2.18	0.41
1:H:193:GLY:O	1:H:194:GLU:C	2.58	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	687/747 (92%)	641 (93%)	39 (6%)	7 (1%)	15 23

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	664	LEU
1	H	645	ASN
1	H	666	ASP
1	H	132	LYS
1	H	647	GLN
1	H	135	ARG
1	H	191	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	H	577/614 (94%)	559 (97%)	18 (3%)	40 57

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	56	LEU
1	H	132	LYS
1	H	190	SER
1	H	194	GLU
1	H	195	ARG
1	H	197	ARG
1	H	199	ASN
1	H	308	THR
1	H	329	GLN
1	H	367	ARG
1	H	415	GLN
1	H	494	GLU
1	H	646	GLN
1	H	660	GLN
1	H	666	ASP
1	H	672	LEU
1	H	688	TRP
1	H	710	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	199	ASN
1	H	328	GLN
1	H	329	GLN
1	H	373	GLN
1	H	418	ASN
1	H	419	GLN
1	H	463	HIS
1	H	482	HIS
1	H	495	GLN
1	H	528	HIS
1	H	546	HIS
1	H	590	GLN
1	H	647	GLN
1	H	660	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	H	801	-	24,29,29	1.03	2 (8%)	29,45,45	1.46	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	H	801	-	-	4/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	801	ADP	C5-C4	2.67	1.48	1.40
2	H	801	ADP	C2-N3	2.00	1.35	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	801	ADP	C3'-C2'-C1'	3.53	106.29	100.98
2	H	801	ADP	N3-C2-N1	-3.45	123.28	128.68
2	H	801	ADP	PA-O3A-PB	-3.00	122.52	132.83
2	H	801	ADP	C4-C5-N7	-2.60	106.69	109.40

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	801	ADP	C5'-O5'-PA-O2A
2	H	801	ADP	C5'-O5'-PA-O3A
2	H	801	ADP	C5'-O5'-PA-O1A
2	H	801	ADP	C4'-C5'-O5'-PA

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	698/747 (93%)	0.22	46 (6%) 18 15	27, 52, 138, 169	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	490	THR	6.9
1	H	52	SER	6.0
1	H	53	GLU	5.8
1	H	51	ARG	5.2
1	H	714	LEU	4.2
1	H	54	LEU	4.1
1	H	132	LYS	4.0
1	H	658	LEU	4.0
1	H	491	GLU	3.9
1	H	169	GLY	3.8
1	H	650	LEU	3.8
1	H	586	ARG	3.7
1	H	712	ASN	3.6
1	H	523	HIS	3.3
1	H	710	LEU	3.2
1	H	713	THR	3.2
1	H	587	ALA	2.9
1	H	429	PHE	2.9
1	H	642	ASP	2.8
1	H	188	ALA	2.7
1	H	427	ARG	2.7
1	H	654	GLN	2.7
1	H	670	TRP	2.7
1	H	408	SER	2.6
1	H	722	TRP	2.6
1	H	458	HIS	2.6
1	H	38	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	716	GLU	2.5
1	H	511	ARG	2.5
1	H	544	GLY	2.5
1	H	133	ARG	2.5
1	H	426	ARG	2.4
1	H	699	ARG	2.4
1	H	128	LEU	2.4
1	H	709	ALA	2.4
1	H	666	ASP	2.3
1	H	57	ASP	2.3
1	H	55	PRO	2.3
1	H	196	LEU	2.2
1	H	448	ASP	2.2
1	H	124	GLN	2.1
1	H	119	ALA	2.1
1	H	193	GLY	2.1
1	H	428	GLY	2.1
1	H	708	LEU	2.0
1	H	385	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ADP	H	801	27/27	0.90	0.20	79,126,134,138	0
3	ZN	H	802	1/1	0.91	0.10	93,93,93,93	0
3	ZN	H	803	1/1	0.98	0.17	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.