



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 04:58 PM EDT

PDB ID : 4PRP
Title : Crystal structure of TK3 TCR-HLA-B*35:01-HPVG-Q5 complex
Authors : Yu Chih, L.; Rossjohn, J.; Gras, S.
Deposited on : 2014-03-06
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 7% 87% 12%
2	B	99	 7% 84% 12%
3	C	11	 91% 9%
4	D	200	 88% 11%
5	E	241	 89% 10%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	2262	1410	414	431	7	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	156	LEU	ARG	SEE REMARK 999	UNP C5MK56

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	829	528	140	158	3	0	0	0

- Molecule 3 is a protein called Epstein-Barr nuclear antigen 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	11	95	62	14	19	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	5	GLN	GLU	SEE REMARK 999	UNP P03211

- Molecule 4 is a protein called TK3 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	200	1557	971	258	321	7	0	0	0

- Molecule 5 is a protein called TK3 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	241	1919	1207	334	373	5	0	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
6	A	1	5	4	1	0	0
6	E	1	5	4	1	0	0
6	E	1	5	4	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	26	26	26	0	0
7	B	7	7	7	0	0
7	C	3	3	3	0	0
7	D	26	26	26	0	0
7	E	15	15	15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Epstein-Barr nuclear antigen 1



- Molecule 4: TK3 TCR alpha chain



- Molecule 5: TK3 TCR beta chain





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	45.14Å 62.60Å 97.81Å 92.04° 102.53° 109.70°	Depositor
Resolution (Å)	47.36 – 2.50 47.36 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.2 (47.36-2.50) 97.2 (47.36-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.51Å)	Xtrriage
Refinement program	PHENIX, BUSTER 2.10.0	Depositor
R, R_{free}	0.180 , 0.229 0.185 , 0.236	Depositor DCC
R_{free} test set	1669 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtrriage
Anisotropy	0.713	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.038 for h,-h-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6754	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2328	0.66	0/3164
2	B	0.47	0/852	0.67	0/1152
3	C	0.62	0/99	0.64	0/133
4	D	0.49	0/1589	0.71	0/2152
5	E	0.51	0/1968	0.67	0/2675
All	All	0.49	0/6836	0.67	0/9276

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2262	0	2134	16	0
2	B	829	0	796	9	0
3	C	95	0	78	1	0
4	D	1557	0	1488	9	0
5	E	1919	0	1825	12	0
6	A	5	0	0	0	0
6	E	10	0	0	0	0
7	A	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	7	0	0	0	0
7	C	3	0	0	0	0
7	D	26	0	0	0	0
7	E	15	0	0	0	0
All	All	6754	0	6321	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:THR:HA	2:B:86:THR:HG21	1.60	0.83
1:A:79[B]:ARG:HE	1:A:82:ARG:HH22	1.37	0.72
2:B:36:GLU:HB3	2:B:83:ASN:HB2	1.73	0.69
2:B:25:CYS:HG	2:B:80:CYS:HG	0.76	0.68
1:A:206:LEU:HD23	1:A:242:GLN:HB3	1.79	0.65
2:B:33:SER:HB2	2:B:54:LEU:HD21	1.84	0.58
4:D:40:PHE:HE1	5:E:112:GLU:HG3	1.71	0.56
4:D:96:PRO:HA	4:D:126:VAL:HB	1.88	0.56
1:A:214:THR:HG23	1:A:262:GLN:HB2	1.89	0.55
2:B:9:VAL:HG23	2:B:93:VAL:HG22	1.89	0.55
1:A:229:GLU:HB2	1:A:246:ALA:HB3	1.89	0.54
4:D:171:TYR:O	4:D:192:ALA:HA	2.08	0.52
4:D:185:PHE:CE2	4:D:187:SER:HB2	2.44	0.52
1:A:109:LEU:HB2	1:A:165:VAL:HG11	1.92	0.52
5:E:221:GLN:HG3	5:E:244:ILE:HD13	1.91	0.51
4:D:96:PRO:HG2	4:D:186:LYS:HE2	1.92	0.50
5:E:10:HIS:HD1	5:E:164:HIS:HD1	1.60	0.50
1:A:234:ARG:HE	1:A:242:GLN:HE21	1.60	0.49
5:E:155:CYS:CB	5:E:220:CYS:SG	3.00	0.49
5:E:107:SER:HB3	5:E:113:LEU:HD23	1.94	0.48
4:D:107:GLN:HG3	4:D:116:LEU:HD23	1.96	0.48
5:E:107:SER:HB3	5:E:113:LEU:CD2	2.44	0.48
1:A:15:PRO:HA	1:A:16:GLY:HA2	1.73	0.47
4:D:140:LEU:HB2	4:D:150:VAL:HG12	1.96	0.47
1:A:187:THR:HG21	1:A:261:VAL:HG21	1.96	0.47
1:A:203:CYS:HG	1:A:259:CYS:HG	1.63	0.46
1:A:203:CYS:SG	1:A:259:CYS:SG	3.13	0.45
1:A:234:ARG:HG3	1:A:242:GLN:HG3	1.99	0.45
1:A:143:THR:HG23	3:C:11:TYR:HA	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:TRP:HZ2	2:B:99:MET:HB3	1.82	0.44
2:B:7:ILE:HG12	2:B:93:VAL:HG13	2.00	0.44
1:A:7:TYR:O	1:A:98:MET:HA	2.19	0.43
4:D:21:LEU:HD12	4:D:21:LEU:N	2.33	0.43
5:E:237:ARG:HH12	5:E:240:PRO:HG3	1.84	0.43
4:D:13:LEU:HD13	4:D:94:PRO:HG3	2.00	0.42
2:B:27:VAL:HG21	2:B:37:VAL:HG21	2.01	0.42
5:E:3:GLY:HA3	5:E:26:ARG:HG2	2.01	0.42
1:A:50:PRO:HD3	1:A:239:ARG:HH22	1.85	0.41
5:E:180:VAL:HA	5:E:203:ARG:O	2.19	0.41
1:A:201:LEU:O	1:A:246:ALA:HA	2.21	0.41
5:E:154:VAL:HG22	5:E:203:ARG:HG3	2.02	0.41
5:E:1:ASP:HA	5:E:2:SER:HA	1.93	0.41
5:E:132:PRO:HD3	5:E:240:PRO:HB3	2.03	0.41
2:B:7:ILE:HG12	2:B:93:VAL:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/276 (100%)	259 (94%)	14 (5%)	2 (1%)	22	39
2	B	97/99 (98%)	91 (94%)	6 (6%)	0	100	100
3	C	9/11 (82%)	8 (89%)	1 (11%)	0	100	100
4	D	198/200 (99%)	192 (97%)	5 (2%)	1 (0%)	29	48
5	E	239/241 (99%)	229 (96%)	9 (4%)	1 (0%)	34	54
All	All	818/827 (99%)	779 (95%)	35 (4%)	4 (0%)	29	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	ASP
1	A	252	GLY
4	D	145	SER
5	E	229	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/234 (100%)	225 (96%)	10 (4%)	29	53
2	B	94/94 (100%)	88 (94%)	6 (6%)	17	33
3	C	9/9 (100%)	9 (100%)	0	100	100
4	D	178/178 (100%)	170 (96%)	8 (4%)	27	51
5	E	209/209 (100%)	202 (97%)	7 (3%)	38	64
All	All	725/724 (100%)	694 (96%)	31 (4%)	29	53

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	GLU
1	A	106	ASP
1	A	110	LEU
1	A	111	ARG
1	A	177	GLU
1	A	214	THR
1	A	228	THR
1	A	231	VAL
1	A	247	VAL
1	A	253	GLU
2	B	7	ILE
2	B	44	GLU
2	B	70	PHE
2	B	80	CYS
2	B	86	THR
2	B	93	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	26	THR
4	D	58	SER
4	D	117	THR
4	D	147	ASP
4	D	150	VAL
4	D	160	THR
4	D	166	LYS
4	D	176	CYS
5	E	11	LEU
5	E	120	ARG
5	E	141	SER
5	E	156	LEU
5	E	185	GLN
5	E	223	GLN
5	E	244	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	GLN
1	A	242	GLN
5	E	86	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	E	302	-	4,4,4	0.12	0	6,6,6	0.17	0
6	SO4	E	301	-	4,4,4	0.16	0	6,6,6	0.13	0
6	SO4	A	301	-	4,4,4	0.19	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	0.21	20 (7%) 15 16	23, 48, 105, 114	65 (23%)
2	B	99/99 (100%)	0.27	7 (7%) 16 16	29, 57, 86, 103	23 (23%)
3	C	11/11 (100%)	-0.57	0 100 100	26, 30, 41, 43	0
4	D	200/200 (100%)	-0.28	2 (1%) 82 84	19, 39, 80, 93	27 (13%)
5	E	241/241 (100%)	-0.25	2 (0%) 86 87	20, 43, 70, 92	35 (14%)
All	All	827/827 (100%)	-0.05	31 (3%) 41 45	19, 44, 87, 114	150 (18%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	GLU	7.0
1	A	194	VAL	6.9
1	A	254	GLU	6.0
1	A	195	SER	5.7
1	A	274	TRP	5.6
1	A	255	GLN	5.0
1	A	252	GLY	4.7
1	A	193	PRO	4.1
1	A	221	GLY	3.5
1	A	197	HIS	3.3
1	A	220	ASP	3.1
1	A	248	VAL	3.0
1	A	257	TYR	2.8
1	A	196	ASP	2.8
2	B	72	PRO	2.6
1	A	198	GLU	2.6
2	B	20	SER	2.6
1	A	192	HIS	2.5
4	D	147	ASP	2.5
2	B	73	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	74	GLU	2.5
1	A	249	VAL	2.4
2	B	15	ALA	2.4
5	E	231	ASP	2.4
1	A	261	VAL	2.4
2	B	23	LEU	2.3
1	A	250	PRO	2.2
2	B	98	ASP	2.2
5	E	230	ASN	2.1
4	D	146	SER	2.1
1	A	223	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	A	301	5/5	0.97	0.15	47,47,48,49	5
6	SO4	E	301	5/5	0.97	0.20	89,90,91,91	0
6	SO4	E	302	5/5	0.97	0.10	95,96,96,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.