



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 06:49 PM EDT

PDB ID : 4QB5
Title : Crystal structure of a glyoxalase/bleomycin resistance protein from *Albidiferax ferrireducens* T118
Authors : Kumaran, D.; Chamala, S.; Evans, B.; Foti, R.; Gizzi, A.; Hillerich, B.; Kar, A.; Lafleur, J.; Seidel, R.; Villigas, G.; Zencheck, W.; Al Obaidi, N.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2014-05-06
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	
1	B	148	
1	C	148	
1	D	148	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyoxalase/bleomycin resistance protein/dioxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	120	893	563	156	169	3	2	0	0	0
1	B	124	915	576	161	173	3	2	0	0	0
1	C	121	902	569	158	170	3	2	0	0	0
1	D	125	922	581	162	174	3	2	0	0	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MSE	-	expression tag	UNP Q21R90
A	-20	HIS	-	expression tag	UNP Q21R90
A	-19	HIS	-	expression tag	UNP Q21R90
A	-18	HIS	-	expression tag	UNP Q21R90
A	-17	HIS	-	expression tag	UNP Q21R90
A	-16	HIS	-	expression tag	UNP Q21R90
A	-15	HIS	-	expression tag	UNP Q21R90
A	-14	SER	-	expression tag	UNP Q21R90
A	-13	SER	-	expression tag	UNP Q21R90
A	-12	GLY	-	expression tag	UNP Q21R90
A	-11	VAL	-	expression tag	UNP Q21R90
A	-10	ASP	-	expression tag	UNP Q21R90
A	-9	LEU	-	expression tag	UNP Q21R90
A	-8	GLY	-	expression tag	UNP Q21R90
A	-7	THR	-	expression tag	UNP Q21R90
A	-6	GLU	-	expression tag	UNP Q21R90
A	-5	ASN	-	expression tag	UNP Q21R90
A	-4	LEU	-	expression tag	UNP Q21R90
A	-3	TYR	-	expression tag	UNP Q21R90
A	-2	PHE	-	expression tag	UNP Q21R90
A	-1	GLN	-	expression tag	UNP Q21R90

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q21R90
B	-21	MSE	-	expression tag	UNP Q21R90
B	-20	HIS	-	expression tag	UNP Q21R90
B	-19	HIS	-	expression tag	UNP Q21R90
B	-18	HIS	-	expression tag	UNP Q21R90
B	-17	HIS	-	expression tag	UNP Q21R90
B	-16	HIS	-	expression tag	UNP Q21R90
B	-15	HIS	-	expression tag	UNP Q21R90
B	-14	SER	-	expression tag	UNP Q21R90
B	-13	SER	-	expression tag	UNP Q21R90
B	-12	GLY	-	expression tag	UNP Q21R90
B	-11	VAL	-	expression tag	UNP Q21R90
B	-10	ASP	-	expression tag	UNP Q21R90
B	-9	LEU	-	expression tag	UNP Q21R90
B	-8	GLY	-	expression tag	UNP Q21R90
B	-7	THR	-	expression tag	UNP Q21R90
B	-6	GLU	-	expression tag	UNP Q21R90
B	-5	ASN	-	expression tag	UNP Q21R90
B	-4	LEU	-	expression tag	UNP Q21R90
B	-3	TYR	-	expression tag	UNP Q21R90
B	-2	PHE	-	expression tag	UNP Q21R90
B	-1	GLN	-	expression tag	UNP Q21R90
B	0	SER	-	expression tag	UNP Q21R90
C	-21	MSE	-	expression tag	UNP Q21R90
C	-20	HIS	-	expression tag	UNP Q21R90
C	-19	HIS	-	expression tag	UNP Q21R90
C	-18	HIS	-	expression tag	UNP Q21R90
C	-17	HIS	-	expression tag	UNP Q21R90
C	-16	HIS	-	expression tag	UNP Q21R90
C	-15	HIS	-	expression tag	UNP Q21R90
C	-14	SER	-	expression tag	UNP Q21R90
C	-13	SER	-	expression tag	UNP Q21R90
C	-12	GLY	-	expression tag	UNP Q21R90
C	-11	VAL	-	expression tag	UNP Q21R90
C	-10	ASP	-	expression tag	UNP Q21R90
C	-9	LEU	-	expression tag	UNP Q21R90
C	-8	GLY	-	expression tag	UNP Q21R90
C	-7	THR	-	expression tag	UNP Q21R90
C	-6	GLU	-	expression tag	UNP Q21R90
C	-5	ASN	-	expression tag	UNP Q21R90
C	-4	LEU	-	expression tag	UNP Q21R90
C	-3	TYR	-	expression tag	UNP Q21R90

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	PHE	-	expression tag	UNP Q21R90
C	-1	GLN	-	expression tag	UNP Q21R90
C	0	SER	-	expression tag	UNP Q21R90
D	-21	MSE	-	expression tag	UNP Q21R90
D	-20	HIS	-	expression tag	UNP Q21R90
D	-19	HIS	-	expression tag	UNP Q21R90
D	-18	HIS	-	expression tag	UNP Q21R90
D	-17	HIS	-	expression tag	UNP Q21R90
D	-16	HIS	-	expression tag	UNP Q21R90
D	-15	HIS	-	expression tag	UNP Q21R90
D	-14	SER	-	expression tag	UNP Q21R90
D	-13	SER	-	expression tag	UNP Q21R90
D	-12	GLY	-	expression tag	UNP Q21R90
D	-11	VAL	-	expression tag	UNP Q21R90
D	-10	ASP	-	expression tag	UNP Q21R90
D	-9	LEU	-	expression tag	UNP Q21R90
D	-8	GLY	-	expression tag	UNP Q21R90
D	-7	THR	-	expression tag	UNP Q21R90
D	-6	GLU	-	expression tag	UNP Q21R90
D	-5	ASN	-	expression tag	UNP Q21R90
D	-4	LEU	-	expression tag	UNP Q21R90
D	-3	TYR	-	expression tag	UNP Q21R90
D	-2	PHE	-	expression tag	UNP Q21R90
D	-1	GLN	-	expression tag	UNP Q21R90
D	0	SER	-	expression tag	UNP Q21R90

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	72	Total O 72 72	0	0
4	B	45	Total O 45 45	0	0
4	C	49	Total O 49 49	0	0
4	D	80	Total O 80 80	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glyoxalase/bleomycin resistance protein/dioxygenase




- Molecule 1: Glyoxalase/bleomycin resistance protein/dioxygenase



- Molecule 1: Glyoxalase/bleomycin resistance protein/dioxygenase



- Molecule 1: Glyoxalase/bleomycin resistance protein/dioxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.95Å 89.90Å 131.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.72 – 2.05 39.72 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.8 (39.72-2.05) 96.8 (39.72-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.57 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.172 , 0.211 0.187 , 0.224	Depositor DCC
R_{free} test set	1734 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtrriage
Anisotropy	0.538	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3897	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.96	1/914 (0.1%)	0.94	4/1236 (0.3%)
1	B	0.89	2/936 (0.2%)	0.96	4/1265 (0.3%)
1	C	0.93	1/923 (0.1%)	0.92	2/1247 (0.2%)
1	D	1.01	2/944 (0.2%)	1.00	5/1277 (0.4%)
All	All	0.95	6/3717 (0.2%)	0.96	15/5025 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	119	CYS	CB-SG	-11.50	1.62	1.82
1	D	29	GLY	N-CA	7.77	1.57	1.46
1	A	119	CYS	CB-SG	-7.75	1.69	1.82
1	C	29	GLY	N-CA	6.69	1.56	1.46
1	B	29	GLY	N-CA	6.60	1.55	1.46
1	B	119	CYS	CB-SG	-6.25	1.71	1.82

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	MSE	CG-SE-CE	6.79	113.84	98.90
1	D	26	LEU	CB-CG-CD1	6.60	122.22	111.00
1	B	110	LEU	CA-CB-CG	6.10	129.34	115.30
1	C	111	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	D	29	GLY	N-CA-C	-5.97	98.18	113.10
1	A	26	LEU	CB-CG-CD1	5.86	120.96	111.00
1	D	112	ASP	CB-CG-OD2	-5.80	113.08	118.30
1	B	111	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	C	37	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	29	GLY	N-CA-C	-5.53	99.28	113.10
1	B	119	CYS	N-CA-CB	-5.44	100.81	110.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	114	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	111	ARG	NE-CZ-NH1	5.19	122.90	120.30
1	D	20	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	29	GLY	N-CA-C	-5.17	100.17	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	893	0	856	7	0
1	B	915	0	883	18	0
1	C	902	0	869	9	0
1	D	922	0	890	14	0
2	B	5	0	0	1	0
2	D	10	0	0	0	0
3	D	4	0	6	0	0
4	A	72	0	0	0	0
4	B	45	0	0	1	0
4	C	49	0	0	0	0
4	D	80	0	0	3	0
All	All	3897	0	3504	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:93:GLU:HG3	1:C:94:PRO:HD2	1.58	0.85
1:B:10:ASN:HD21	1:B:58:ALA:H	1.35	0.73
1:C:10:ASN:HD21	1:C:58:ALA:H	1.35	0.73
1:A:5:VAL:O	1:A:6:MSE:HE2	1.93	0.68
1:D:10:ASN:HD21	1:D:58:ALA:H	1.43	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:ALA:O	1:B:102:VAL:HG23	1.95	0.66
1:B:123:ARG:C	4:B:304:HOH:O	2.35	0.65
1:C:10:ASN:ND2	1:C:58:ALA:H	2.01	0.57
1:A:1:MSE:HE3	1:C:42:SER:HB3	1.87	0.56
1:B:10:ASN:ND2	1:B:58:ALA:H	2.04	0.56
1:D:37:ARG:HD3	4:D:367:HOH:O	2.06	0.55
1:B:101:ALA:HB3	1:D:34:ASN:HD21	1.70	0.55
1:A:1:MSE:HE3	1:C:42:SER:CB	2.38	0.54
1:B:91:CYS:HA	2:B:201:SO4:O1	2.08	0.53
1:D:36:SER:C	1:D:37:ARG:HG3	2.31	0.51
1:B:1:MSE:HE3	1:D:42:SER:HB3	1.91	0.50
1:D:10:ASN:ND2	1:D:58:ALA:H	2.07	0.49
1:B:23:LEU:HB3	1:B:28:ILE:HB	1.94	0.49
1:B:1:MSE:HE3	1:D:42:SER:CB	2.43	0.49
1:B:101:ALA:O	1:B:102:VAL:CG2	2.60	0.48
1:B:100:GLY:O	1:B:102:VAL:N	2.46	0.48
1:C:80:HIS:HE1	1:C:90:THR:OG1	1.97	0.47
1:B:42:SER:OG	1:D:1:MSE:HE3	2.14	0.47
1:A:1:MSE:CE	1:C:42:SER:HB3	2.45	0.46
1:D:107:LEU:HD21	1:D:121:LEU:HD12	1.97	0.46
1:B:101:ALA:C	1:B:102:VAL:HG23	2.36	0.45
1:A:80:HIS:O	1:A:84:ILE:HG12	2.16	0.45
1:B:112:ASP:HB2	1:B:113:PRO:CD	2.47	0.44
1:D:0:SER:N	4:D:378:HOH:O	2.51	0.43
1:D:10:ASN:HD21	1:D:58:ALA:N	2.13	0.43
1:D:98:ARG:NH1	4:D:377:HOH:O	2.51	0.43
1:A:1:MSE:HE3	1:C:42:SER:OG	2.19	0.43
1:A:42:SER:HB2	1:A:43:PRO:HD2	2.01	0.43
1:B:73:PRO:HD3	1:B:122:HIS:CD2	2.54	0.42
1:B:1:MSE:CE	1:D:42:SER:HB3	2.47	0.42
1:B:1:MSE:HE2	1:D:28:ILE:HD12	2.01	0.42
1:B:10:ASN:HD21	1:B:58:ALA:N	2.11	0.40
1:C:16:LYS:HD2	1:C:38:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/148 (78%)	113 (97%)	3 (3%)	0	100	100
1	B	122/148 (82%)	115 (94%)	5 (4%)	2 (2%)	8	2
1	C	117/148 (79%)	112 (96%)	5 (4%)	0	100	100
1	D	123/148 (83%)	117 (95%)	6 (5%)	0	100	100
All	All	478/592 (81%)	457 (96%)	19 (4%)	2 (0%)	30	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	101	ALA
1	B	102	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/111 (83%)	89 (97%)	3 (3%)	33	27
1	B	93/111 (84%)	91 (98%)	2 (2%)	47	43
1	C	93/111 (84%)	90 (97%)	3 (3%)	34	28
1	D	94/111 (85%)	91 (97%)	3 (3%)	34	28
All	All	372/444 (84%)	361 (97%)	11 (3%)	36	31

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	37	ARG
1	A	118	ILE
1	B	37	ARG
1	B	110	LEU
1	C	37	ARG
1	C	92	GLU
1	C	93	GLU
1	D	26	LEU
1	D	104	LYS
1	D	112	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	86	ASN
1	A	116	ASN
1	A	122	HIS
1	B	10	ASN
1	B	80	HIS
1	B	86	ASN
1	B	116	ASN
1	B	122	HIS
1	C	10	ASN
1	C	80	HIS
1	C	86	ASN
1	C	116	ASN
1	D	10	ASN
1	D	60	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	D	203	-	3,3,3	0.64	0	2,2,2	0.27	0
2	SO4	B	201	-	4,4,4	0.43	0	6,6,6	0.82	0
2	SO4	D	201	-	4,4,4	0.57	0	6,6,6	0.49	0
2	SO4	D	202	-	4,4,4	0.50	0	6,6,6	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	203	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	201	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/148 (79%)	-0.32	2 (1%) 69 71	12, 19, 42, 64	0
1	B	122/148 (82%)	0.14	10 (8%) 19 20	16, 24, 56, 108	0
1	C	119/148 (80%)	-0.27	3 (2%) 58 60	13, 19, 42, 69	0
1	D	123/148 (83%)	-0.24	3 (2%) 59 61	11, 19, 50, 88	0
All	All	482/592 (81%)	-0.17	18 (3%) 45 47	11, 20, 49, 108	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	101	ALA	6.7
1	B	102	VAL	5.1
1	D	102	VAL	5.1
1	B	100	GLY	5.0
1	C	99	ASP	3.4
1	D	101	ALA	3.3
1	C	92	GLU	3.3
1	D	100	GLY	3.3
1	B	43	PRO	3.1
1	B	99	ASP	3.0
1	B	98	ARG	2.7
1	B	123	ARG	2.5
1	B	29	GLY	2.5
1	C	98	ARG	2.4
1	B	44	ALA	2.2
1	B	103	GLY	2.1
1	A	93	GLU	2.1
1	A	98	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	201	5/5	0.90	0.11	55,58,62,68	0
3	EDO	D	203	4/4	0.95	0.07	22,22,23,26	0
2	SO4	D	201	5/5	0.96	0.08	30,35,35,42	0
2	SO4	D	202	5/5	0.97	0.11	30,31,34,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.