



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 09:51 PM EDT

PDB ID : 4QUE
Title : Caspase-3 Y195FV266H
Authors : Cade, C.; Swartz, P.D.; MacKenzie, S.H.; Clark, A.C.
Deposited on : 2014-07-10
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1150 (1.84-1.84)
Clashscore	180529	1248 (1.84-1.84)
Ramachandran outliers	177936	1240 (1.84-1.84)
Sidechain outliers	177891	1240 (1.84-1.84)
RSRZ outliers	164620	1149 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	 71% 13% 16%
1	C	277	 68% 15% 16%
2	H	6	 83% 17%
2	J	6	 50% 50%
3	D	5	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	5	 20% 20% 40% 20%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4343 atoms, of which 1 is hydrogen and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	1948	1236	339	357	16	0	9	0
1	C	232	1928	1225	333	354	16	0	8	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	PHE	TYR	engineered mutation	UNP P42574
A	266	HIS	VAL	engineered mutation	UNP P42574
C	195	PHE	TYR	engineered mutation	UNP P42574
C	266	HIS	VAL	engineered mutation	UNP P42574


- Molecule 2 is a protein called ACE-ASP-GLU-VAL-ASP-CHLOROMETHYLKETONE INHIBITOR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	6	36	21	4	11	0	0	1
2	J	6	36	21	4	11	0	0	1

- Molecule 3 is a protein called SHORT PEPTIDE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	D	5	38	20	1	5	11	1	0	0	0
3	E	5	39	22	5	11	1	0	0	0	

- Molecule 4 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: C₄H₁₀O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 5 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	N	0	0
			3	3		
5	C	1	Total	N	0	0
			3	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	156	Total O 156 156	0	0
6	C	129	Total O 129 129	0	0
6	H	6	Total O 6 6	0	0
6	J	7	Total O 7 7	0	0
6	D	4	Total O 4 4	0	0
6	E	2	Total O 2 2	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Caspase-3

Chain A: 




- Molecule 1: Caspase-3

Chain C: 



- Molecule 2: ACE-ASP-GLU-VAL-ASP-CHLOROMETHYLKETONE INHIBITOR

Chain H: 



- Molecule 2: ACE-ASP-GLU-VAL-ASP-CHLOROMETHYLKETONE INHIBITOR

Chain J: 



- Molecule 3: SHORT PEPTIDE

Chain D:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: SHORT PEPTIDE

Chain E:  20% 20% 40% 20%

V178	D179	D180	D181	V182
------	------	------	------	------

4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.12Å 96.36Å 68.83Å 90.00° 127.10° 90.00°	Depositor
Resolution (Å)	34.84 – 1.84 34.84 – 1.84	Depositor EDS
% Data completeness (in resolution range)	97.6 (34.84-1.84) 92.5 (34.84-1.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.84Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.180 , 0.223 0.182 , 0.221	Depositor DCC
R_{free} test set	1890 reflections (4.17%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.187 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4343	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DTT, 0QE, ACE, AZI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2005	0.54	0/2691
1	C	0.39	0/1983	0.52	0/2660
2	H	1.84	0/32	1.59	0/43
2	J	1.85	1/32 (3.1%)	1.61	0/43
3	D	0.24	0/36	0.63	0/47
3	E	0.25	0/38	0.62	0/50
All	All	0.44	1/4126 (0.0%)	0.56	0/5534

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	2	ASP	C-N	5.18	1.46	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1948	0	1939	34	0
1	C	1928	0	1909	57	0
2	H	36	0	26	1	0
2	J	36	0	26	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	37	1	22	0	0
3	E	39	0	29	4	0
4	A	8	0	10	2	0
5	A	3	0	0	0	0
5	C	3	0	0	1	0
6	A	156	0	0	8	0
6	C	129	0	0	2	0
6	D	4	0	0	0	0
6	E	2	0	0	0	0
6	H	6	0	0	0	0
6	J	7	0	0	0	0
All	All	4342	1	3961	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:120:SER:HB3	1:C:127[A]:ILE:CD1	2.10	0.82
1:A:238[B]:ARG:NH1	6:A:470:HOH:O	2.14	0.80
1:A:118:LEU:HB3	1:A:127:ILE:CD1	2.12	0.80
1:A:187[A]:ILE:HG13	1:A:188:PRO:HD2	1.65	0.79
1:A:118:LEU:HB3	1:A:127:ILE:HD12	1.67	0.76
1:C:61[B]:MET:HE3	1:C:121:HIS:HB2	1.68	0.75
3:E:179:ASP:O	3:E:180:ASP:HB2	1.86	0.75
1:A:149[B]:ARG:NH1	6:A:444:HOH:O	2.17	0.70
1:A:241:ARG:NH1	1:C:271:LYS:HA	2.07	0.69
1:C:80:ASN:ND2	6:C:509:HOH:O	2.18	0.69
1:A:241:ARG:HD3	1:C:270:THR:O	1.91	0.69
1:C:127[B]:ILE:CD1	1:C:134:VAL:HG13	2.25	0.67
1:A:34:ASP:O	1:A:35:ASN:HB2	1.96	0.65
1:A:101[B]:ARG:NH2	1:A:102:ASP:OD1	2.31	0.63
1:A:72:ALA:HB2	3:E:178:VAL:HG11	1.81	0.63
1:C:127[B]:ILE:CG2	1:C:136:LEU:HD21	2.27	0.63
1:C:120:SER:HB3	1:C:127[A]:ILE:HD11	1.81	0.62
6:A:537:HOH:O	3:E:178:VAL:HG12	2.00	0.62
1:A:163:CYS:SG	2:J:5:ASP:C	2.78	0.62
1:C:207:ARG:HA	1:C:213:SER:HA	1.81	0.62
1:C:100:MET:HG3	1:C:139:ILE:HG23	1.81	0.61
1:A:93:ARG:HB2	1:A:134:VAL:HG12	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:88:LYS:HE2	1:C:88:LYS:HA	1.81	0.59
1:A:184:CYS:O	6:A:548:HOH:O	2.17	0.59
1:C:163:CYS:SG	2:H:5:ASP:C	2.80	0.59
1:C:127[B]:ILE:HD11	1:C:134:VAL:HG13	1.85	0.59
1:C:61[B]:MET:CE	1:C:121:HIS:HB2	2.33	0.59
1:C:127[B]:ILE:HG23	1:C:136:LEU:HD21	1.85	0.58
1:A:187[A]:ILE:HG13	1:A:188:PRO:CD	2.32	0.58
1:C:61[B]:MET:HE3	1:C:121:HIS:CB	2.34	0.57
1:A:101[A]:ARG:O	1:A:105:LYS:HG2	2.05	0.56
1:C:127[A]:ILE:HG22	1:C:134:VAL:HG13	1.88	0.54
1:C:53:LYS:HE3	5:C:301:AZI:N1	2.22	0.54
1:C:172:ILE:O	1:C:173[B]:GLU:HB3	2.08	0.54
1:C:61[B]:MET:HE1	1:C:128:PHE:CB	2.38	0.54
1:A:207:ARG:HA	1:A:213:SER:HA	1.90	0.53
1:C:228:ASP:OD1	1:C:229:LYS:HG3	2.08	0.53
1:A:98:GLU:HG3	1:A:101[B]:ARG:HH22	1.74	0.53
1:C:127[B]:ILE:HD11	1:C:139:ILE:CD1	2.41	0.51
1:A:224[A]:LYS:HG2	6:A:552:HOH:O	2.10	0.51
1:C:61[B]:MET:HE1	1:C:128:PHE:HB3	1.94	0.49
1:A:264:CYS:SG	1:C:266[B]:HIS:CE1	3.06	0.49
1:A:135[A]:ASP:OD1	6:A:517:HOH:O	2.20	0.49
1:A:101[B]:ARG:O	1:A:105:LYS:HG2	2.11	0.49
1:C:127[B]:ILE:HG21	1:C:136:LEU:HD21	1.95	0.49
1:C:61[B]:MET:HE2	1:C:121:HIS:HD2	1.78	0.48
1:C:231:GLU:HG3	1:C:272:GLU:HB3	1.95	0.48
1:C:127[B]:ILE:HD11	1:C:139:ILE:HD11	1.95	0.48
1:C:194:LEU:CD1	1:C:233:MET:CE	2.92	0.48
1:A:228:ASP:OD1	1:A:229:LYS:NZ	2.44	0.48
1:A:93:ARG:NH1	6:A:550:HOH:O	2.47	0.47
1:C:195:PHE:HB2	1:C:266[B]:HIS:HB3	1.97	0.47
1:C:157:LEU:HD11	1:C:273:LEU:HD21	1.97	0.47
1:A:231:GLU:HG3	1:A:272:GLU:HB3	1.97	0.47
1:C:264:CYS:HG	1:C:266[A]:HIS:CD2	2.34	0.46
1:C:74:LEU:HD13	1:C:117:VAL:HG11	1.97	0.45
4:A:301:DTT:S4	2:J:6:0QE:C1	3.05	0.45
4:A:301:DTT:H42	2:J:6:0QE:C1	2.47	0.44
1:C:194:LEU:HD13	1:C:233:MET:HE3	2.00	0.44
1:C:172:ILE:HG13	1:C:173[B]:GLU:N	2.33	0.44
1:C:193:PHE:HB2	1:C:268:MET:HE3	2.00	0.44
1:A:87:ASN:O	3:E:180:ASP:HA	2.17	0.44
1:A:241:ARG:NH1	1:C:270:THR:O	2.38	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:CYS:O	1:A:224[A]:LYS:HG3	2.17	0.43
1:C:61[B]:MET:HE2	1:C:121:HIS:CD2	2.53	0.43
1:C:127[B]:ILE:H	1:C:127[B]:ILE:HD13	1.83	0.43
1:C:159:ILE:HD13	1:C:219:LEU:HD21	2.01	0.43
1:C:172:ILE:HG13	1:C:173[A]:GLU:N	2.34	0.42
1:A:149[B]:ARG:HA	1:A:149[B]:ARG:HD2	1.80	0.42
1:C:172:ILE:O	1:C:173[A]:GLU:HB3	2.19	0.42
1:A:266:HIS:HD1	1:C:266[A]:HIS:CE1	2.36	0.42
1:C:210:LYS:HD2	1:C:211:ASP:OD1	2.20	0.42
1:C:59:THR:HG21	1:C:61[B]:MET:SD	2.60	0.42
1:C:127[B]:ILE:HG23	1:C:136:LEU:CD2	2.49	0.42
1:A:149[A]:ARG:HD3	6:A:520:HOH:O	2.19	0.42
1:C:127[B]:ILE:HD13	1:C:134:VAL:HG13	1.99	0.42
1:C:195:PHE:HE2	1:C:268:MET:HE1	1.83	0.42
1:C:194:LEU:HD13	1:C:233:MET:CE	2.50	0.42
1:C:124:GLU:OE2	6:C:481:HOH:O	2.21	0.41
1:C:264:CYS:SG	1:C:266[A]:HIS:CD2	3.14	0.41
1:C:140:THR:HG21	1:C:195:PHE:CE1	2.56	0.41
1:C:195:PHE:HB2	1:C:266[A]:HIS:HB2	2.02	0.41
1:C:194:LEU:CD1	1:C:233:MET:HE3	2.51	0.41
1:A:74:LEU:HD13	1:A:117:VAL:HG11	2.02	0.41
1:A:140:THR:HG21	1:A:195:PHE:CE1	2.56	0.40
1:A:218:SER:HB3	1:A:242:LYS:HD3	2.03	0.40
1:A:163:CYS:SG	2:J:5:ASP:N	2.94	0.40
1:C:55:PHE:CD2	1:C:130:THR:HA	2.56	0.40
1:C:140:THR:HG21	1:C:195:PHE:HE1	1.86	0.40
1:C:127[B]:ILE:HD12	1:C:136:LEU:HD23	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/277 (86%)	234 (98%)	4 (2%)	1 (0%)	30	18
1	C	235/277 (85%)	228 (97%)	7 (3%)	0	100	100
2	H	3/6 (50%)	3 (100%)	0	0	100	100
2	J	3/6 (50%)	3 (100%)	0	0	100	100
3	D	3/5 (60%)	3 (100%)	0	0	100	100
3	E	3/5 (60%)	1 (33%)	1 (33%)	1 (33%)	0	0
All	All	486/576 (84%)	472 (97%)	12 (2%)	2 (0%)	30	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
3	E	180	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/249 (87%)	215 (99%)	2 (1%)	75	67
1	C	215/249 (86%)	211 (98%)	4 (2%)	52	36
2	H	4/4 (100%)	4 (100%)	0	100	100
2	J	4/4 (100%)	4 (100%)	0	100	100
3	D	4/5 (80%)	4 (100%)	0	100	100
3	E	5/5 (100%)	1 (20%)	4 (80%)	0	0
All	All	449/516 (87%)	439 (98%)	10 (2%)	48	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	VAL
1	A	174	THR
1	C	58	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	59	THR
1	C	127[A]	ILE
1	C	127[B]	ILE
3	E	178	VAL
3	E	179	ASP
3	E	180	ASP
3	E	181	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DTT	A	301	-	7,7,7	0.63	0	4,8,8	1.14	0
5	AZI	A	302	-	2,2,2	4.31	1 (50%)	0,1,1	-	-
5	AZI	C	301	-	2,2,2	3.53	1 (50%)	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	A	301	-	-	0/8/8/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	302	AZI	N1-N2	-5.96	1.10	1.23
5	C	301	AZI	N1-N2	-4.59	1.13	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	301	DTT	2	0
5	C	301	AZI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/277 (84%)	-1.29	0 100 100	10, 20, 34, 53	9 (3%)
1	C	232/277 (83%)	-1.27	0 100 100	11, 21, 38, 65	8 (3%)
2	H	4/6 (66%)	-1.48	0 100 100	22, 23, 24, 24	0
2	J	4/6 (66%)	-1.48	0 100 100	16, 17, 19, 22	0
3	D	5/5 (100%)	-0.46	0 100 100	41, 42, 50, 50	0
3	E	5/5 (100%)	-0.39	0 100 100	42, 49, 54, 63	0
All	All	484/576 (84%)	-1.27	0 100 100	10, 21, 39, 65	17 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DTT	A	301	8/8	0.97	0.08	51,62,72,73	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	AZI	C	301	3/3	0.98	0.07	21,21,31,32	0
5	AZI	A	302	3/3	0.99	0.07	17,17,28,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.