



# Full wwPDB X-ray Structure Validation Report i

Sep 16, 2023 – 09:51 PM EDT

PDB ID : 4TXH  
Title : Crystal structure of uridine phosphorylase from Schistosoma mansoni in APO form  
Authors : Torini, J.; Romanello, L.; Marinho, A.; Brandao-Neto, J.; Cassago, A.; De-Marco, R.; Pereira, H.M.  
Deposited on : 2014-07-03  
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)


There are 3 unique types of molecules in this entry. The entry contains 9360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total 2237	C 1418	N 388	O 417	S 14	0	2	0
1	B	290	Total 2246	C 1427	N 389	O 416	S 14	0	1	0
1	C	286	Total 2087	C 1323	N 360	O 391	S 13	0	1	0
1	D	280	Total 2003	C 1256	N 354	O 381	S 12	0	2	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	298	Total O 298 298	0	0
3	B	258	Total O 258 258	0	0
3	C	102	Total O 102 102	0	0
3	D	119	Total O 119 119	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase



- Molecule 1: Uridine phosphorylase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.07 Å    109.03 Å    119.04 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	80.40 – 1.89 80.40 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.9 (80.40-1.89) 99.9 (80.40-1.89)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.38 (at 1.90 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
$R$ , $R_{free}$	0.215 , 0.239 0.215 , 0.237	Depositor DCC
$R_{free}$ test set	4998 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9360	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/2280	0.46	0/3082
1	B	0.26	0/2286	0.45	0/3087
1	C	0.25	0/2125	0.43	0/2883
1	D	0.26	0/2039	0.44	0/2761
All	All	0.26	0/8730	0.44	0/11813

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	0	2215	6	0
1	B	2246	0	2244	9	0
1	C	2087	0	1983	27	0
1	D	2003	0	1786	49	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	298	0	0	3	1
3	B	258	0	0	3	0
3	C	102	0	0	12	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	119	0	0	24	1
All	All	9360	0	8228	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:45:CYS:SG	3:D:335:HOH:O	2.26	0.94
1:D:42:VAL:HG12	1:D:87:LEU:HB2	1.48	0.91
1:A:162:GLN:OE1	3:A:401:HOH:O	1.90	0.90
1:D:252:VAL:O	3:D:301:HOH:O	1.95	0.83
1:D:253:CYS:HA	3:D:301:HOH:O	1.80	0.81
1:D:52:HIS:HB3	3:D:310:HOH:O	1.82	0.79
1:D:285:GLU:N	3:D:303:HOH:O	2.05	0.78
1:D:137:LEU:HB3	1:D:252:VAL:HG12	1.65	0.77
1:A:196:ASP:OD1	3:A:402:HOH:O	2.01	0.77
1:D:178:GLU:OE2	3:D:302:HOH:O	2.01	0.77
1:C:213:LYS:NZ	3:C:305:HOH:O	2.17	0.77
1:D:233:MET:SD	3:D:415:HOH:O	2.42	0.76
1:D:141:CYS:SG	1:D:239:THR:OG1	2.43	0.76
1:D:95:ILE:HG23	1:D:237:ALA:HB2	1.67	0.75
3:C:319:HOH:O	1:D:96:PRO:HG2	1.89	0.73
1:C:184:CYS:SG	3:C:312:HOH:O	2.46	0.73
1:B:216:LYS:NZ	3:B:402:HOH:O	2.20	0.73
1:D:201:GLN:HB3	1:D:259:ARG:HH12	1.53	0.72
1:D:149:TYR:O	3:D:304:HOH:O	2.08	0.71
1:C:199:GLU:OE2	3:C:302:HOH:O	2.10	0.69
1:C:194:ALA:O	3:C:303:HOH:O	2.11	0.69
1:D:247:HIS:O	3:D:305:HOH:O	2.12	0.68
1:C:215:GLU:O	3:C:304:HOH:O	2.12	0.67
1:D:133:GLY:O	1:D:278:ARG:NH1	2.28	0.66
1:D:143:ASN:HD21	1:D:147:GLU:HB2	1.62	0.65
1:D:153:CYS:SG	3:D:332:HOH:O	2.54	0.64
1:D:248:ARG:HA	3:D:305:HOH:O	2.00	0.61
1:D:286:TYR:O	1:D:290:ASN:ND2	2.30	0.60
1:D:241:HIS:ND1	3:D:320:HOH:O	2.30	0.60
1:C:86:ILE:HD13	1:C:287:LEU:HD13	1.85	0.59
1:C:147:GLU:OE2	1:C:147:GLU:N	2.30	0.59
1:D:196:ASP:HB3	1:D:200:GLU:HB2	1.86	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:50[A]:ARG:O	3:D:307:HOH:O	2.18	0.57
1:D:151:GLU:O	3:D:306:HOH:O	2.17	0.57
1:D:114:CYS:O	3:D:309:HOH:O	2.18	0.56
1:D:74:SER:O	3:D:308:HOH:O	2.18	0.56
1:C:107:LYS:NZ	3:C:313:HOH:O	2.31	0.55
1:D:50[B]:ARG:O	3:D:307:HOH:O	2.18	0.54
1:C:214:GLU:OE2	3:C:306:HOH:O	2.18	0.54
1:D:136:VAL:HG11	1:D:230:ASN:HB3	1.88	0.54
1:B:140:ARG:NH2	3:B:401:HOH:O	2.02	0.54
1:A:95:ILE:HG23	1:A:237:ALA:HB2	1.91	0.53
1:B:80:LEU:HD11	1:B:87:LEU:HG	1.91	0.52
1:C:95:ILE:HG23	1:C:237:ALA:HB2	1.90	0.52
1:D:174:LYS:O	1:D:177:SER:OG	2.25	0.52
1:C:144:THR:HG21	1:D:153:CYS:O	2.11	0.51
1:D:49:GLY:O	3:D:310:HOH:O	2.19	0.51
1:C:138:SER:O	1:C:165:VAL:HG11	2.11	0.50
1:D:256:ALA:N	3:D:315:HOH:O	2.44	0.50
1:C:224:TYR:N	3:C:301:HOH:O	2.45	0.49
1:D:223:ALA:HB1	1:D:228:ILE:HD12	1.94	0.49
1:D:249:ALA:HB3	3:D:343:HOH:O	2.12	0.49
1:C:215:GLU:C	3:C:304:HOH:O	2.49	0.48
1:D:91:HIS:HB2	1:D:98:ALA:HA	1.96	0.48
1:D:195:ASN:HA	3:D:314:HOH:O	2.14	0.47
1:C:196:ASP:OD2	1:C:199:GLU:N	2.47	0.47
1:C:18:ASP:OD2	1:C:111:TYR:OH	2.16	0.46
1:B:196:ASP:HB3	1:B:200:GLU:HB2	1.96	0.46
1:A:196:ASP:HB3	1:A:200:GLU:HB2	1.97	0.46
1:D:95:ILE:HD13	1:D:234:GLU:HA	1.97	0.46
1:B:47:SER:HB3	1:B:50[B]:ARG:HG3	1.98	0.46
1:C:229:ARG:HA	1:C:229:ARG:HD3	1.77	0.45
1:C:20:PHE:HD2	1:C:104:GLU:HG3	1.81	0.45
1:D:95:ILE:HB	1:D:96:PRO:HD3	1.99	0.45
1:A:183:GLU:HG2	3:A:550:HOH:O	2.17	0.44
1:B:7:ILE:HD12	1:B:8:VAL:HG23	1.99	0.44
1:D:195:ASN:N	1:D:200:GLU:OE1	2.42	0.44
1:C:196:ASP:HB3	1:C:200:GLU:HG3	2.01	0.43
1:C:202:GLY:O	1:C:216:LYS:HG3	2.18	0.43
1:B:196:ASP:OD1	3:B:403:HOH:O	2.21	0.43
1:C:221:GLN:C	3:C:301:HOH:O	2.57	0.43
1:D:184:CYS:SG	3:D:375:HOH:O	2.61	0.43
1:D:81:TYR:HB2	1:D:88:PHE:HB2	2.01	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:154:ILE:HD12	1:C:159:VAL:HG21	1.99	0.43
1:D:52:HIS:O	1:D:56:ILE:HG13	2.19	0.43
1:D:137:LEU:HA	1:D:251:LEU:O	2.19	0.42
1:D:221:GLN:O	1:D:224:TYR:N	2.51	0.42
1:D:27:THR:HA	1:D:32:PHE:CE1	2.55	0.42
1:D:119:PHE:O	1:D:249:ALA:HA	2.18	0.42
1:C:22:HIS:N	1:C:104:GLU:OE1	2.50	0.42
1:D:136:VAL:HG23	3:D:321:HOH:O	2.18	0.42
1:C:165:VAL:O	1:C:167:LEU:N	2.52	0.42
1:C:289:ARG:NH1	3:C:309:HOH:O	2.26	0.42
1:D:284:GLY:N	3:D:303:HOH:O	2.53	0.42
1:D:128:LEU:N	1:D:229:ARG:O	2.50	0.41
1:B:18:ASP:HB3	1:B:27:THR:HG23	2.03	0.41
1:A:128:LEU:HD23	1:A:259:ARG:HB2	2.04	0.40
1:B:95:ILE:HG23	1:B:237:ALA:HB2	2.03	0.40
1:C:134:THR:HG23	1:C:185:SER:O	2.21	0.40
1:C:27:THR:HA	1:C:32:PHE:CE1	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:614:HOH:O	3:D:414:HOH:O[4_445]	2.17	0.03

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/296 (98%)	279 (96%)	10 (4%)	0	100 100
1	B	289/296 (98%)	283 (98%)	6 (2%)	0	100 100
1	C	283/296 (96%)	270 (95%)	12 (4%)	1 (0%)	34 24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	271/296 (92%)	261 (96%)	10 (4%)	0	100	100
All	All	1132/1184 (96%)	1093 (97%)	38 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	166	ASP

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/252 (96%)	239 (99%)	3 (1%)	71	70
1	B	244/252 (97%)	242 (99%)	2 (1%)	81	82
1	C	211/252 (84%)	209 (99%)	2 (1%)	78	79
1	D	188/252 (75%)	179 (95%)	9 (5%)	25	16
All	All	885/1008 (88%)	869 (98%)	16 (2%)	59	55

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	PHE
1	A	66	LEU
1	A	232	GLU
1	B	30	PHE
1	B	232	GLU
1	C	104	GLU
1	C	211	PHE
1	D	30	PHE
1	D	136	VAL
1	D	137	LEU
1	D	146	LEU
1	D	231	MET
1	D	232	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	251	LEU
1	D	268	SER
1	D	273	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	301	-	4,4,4	0.10	0	6,6,6	0.13	0
2	SO4	A	301	-	4,4,4	0.15	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	289/296 (97%)	-0.24	0 [100] [100]	16, 25, 42, 55	0
1	B	290/296 (97%)	-0.20	0 [100] [100]	17, 29, 52, 70	0
1	C	286/296 (96%)	0.65	22 (7%) [13] [15]	27, 53, 78, 86	0
1	D	280/296 (94%)	1.53	90 (32%) [0] [0]	24, 70, 100, 119	0
All	All	1145/1184 (96%)	0.42	112 (9%) [7] [8]	16, 38, 86, 119	0

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	83	VAL	6.9
1	D	260	LEU	6.8
1	D	231	MET	6.5
1	D	276	ALA	6.3
1	D	225	GLU	5.3
1	D	123	GLY	4.9
1	D	44	VAL	4.9
1	D	250	ILE	4.9
1	D	249	ALA	4.8
1	D	169	THR	4.4
1	D	252	VAL	4.2
1	D	167	LEU	4.2
1	D	251	LEU	4.2
1	D	124	THR	4.1
1	D	129	GLY	4.1
1	D	136	VAL	4.0
1	D	148	PRO	4.0
1	D	283	VAL	4.0
1	D	67	PRO	3.9
1	C	75	HIS	3.9
1	D	56	ILE	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	135	ILE	3.9
1	D	228	ILE	3.9
1	D	294	ILE	3.9
1	D	134	THR	3.8
1	D	192	ILE	3.8
1	D	153	CYS	3.8
1	D	287	LEU	3.6
1	D	127	GLY	3.6
1	D	128	LEU	3.6
1	D	120	ILE	3.6
1	D	279	PRO	3.5
1	D	74	SER	3.4
1	D	59	ALA	3.4
1	C	182	LEU	3.4
1	C	159	VAL	3.3
1	D	149	TYR	3.3
1	C	20	PHE	3.3
1	D	187	VAL	3.3
1	D	253	CYS	3.3
1	D	257	VAL	3.2
1	D	86	ILE	3.2
1	D	126	GLY	3.2
1	D	43	CYS	3.1
1	D	88	PHE	3.1
1	D	185	SER	3.1
1	D	141	CYS	3.1
1	C	223	ALA	3.1
1	C	148	PRO	3.0
1	D	68	VAL	3.0
1	D	119	PHE	3.0
1	D	139	ASP	3.0
1	D	54	PHE	2.9
1	D	219	PHE	2.9
1	D	230	ASN	2.9
1	C	153	CYS	2.9
1	D	176	LEU	2.9
1	D	272	PHE	2.9
1	C	21	HIS	2.9
1	D	186	VAL	2.9
1	C	222	SER	2.8
1	D	256	ALA	2.8
1	D	178	GLU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	191	THR	2.7
1	C	249	ALA	2.7
1	C	154	ILE	2.7
1	D	243	TYR	2.7
1	D	170	VAL	2.7
1	C	98	ALA	2.6
1	D	222	SER	2.6
1	D	144	THR	2.5
1	C	144	THR	2.5
1	D	152	LEU	2.5
1	D	217	LEU	2.5
1	D	133	GLY	2.5
1	D	164	ILE	2.4
1	C	156	GLY	2.4
1	D	168[A]	ASN	2.4
1	D	64	LEU	2.4
1	C	244	LEU	2.3
1	D	262	GLY	2.3
1	D	233	MET	2.3
1	D	254	VAL	2.3
1	D	237	ALA	2.3
1	D	202	GLY	2.3
1	D	137	LEU	2.3
1	D	204	LEU	2.3
1	C	158	PRO	2.3
1	D	122	LEU	2.3
1	C	236	THR	2.2
1	D	130	VAL	2.2
1	D	278	ARG	2.2
1	D	132	PRO	2.2
1	D	177	SER	2.2
1	D	211	PHE	2.2
1	D	289	ARG	2.1
1	C	110	HIS	2.1
1	D	245	THR	2.1
1	D	284	GLY	2.1
1	D	173	LEU	2.1
1	D	57	SER	2.1
1	D	285	GLU	2.1
1	D	165	VAL	2.1
1	D	291	ASN	2.1
1	D	282	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	10	SER	2.0
1	D	142	VAL	2.0
1	D	238	ILE	2.0
1	C	73	GLY	2.0
1	D	227	GLY	2.0
1	C	180	LEU	2.0
1	C	220	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	301	5/5	0.99	0.11	18,22,23,25	0
2	SO4	B	301	5/5	0.99	0.10	21,22,24,24	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.