



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 07:22 pm GMT

PDB ID : 4UWY  
Title : FGFR1 Apo structure  
Authors : Thiyagarajan, N.; Bunney, T.; Katan, M.  
Deposited on : 2014-08-15  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	
1	B	310	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FIBROBLAST GROWTH FACTOR RECEPTOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	2396	1527	412	439	18	0	4	0
1	B	273	2171	1380	372	402	17	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	456	MET	-	expression tag	UNP P11362
A	457	VAL	-	expression tag	UNP P11362
A	488	ALA	CYS	conflict	UNP P11362
A	584	SER	CYS	conflict	UNP P11362
B	456	MET	-	expression tag	UNP P11362
B	457	VAL	-	expression tag	UNP P11362
B	488	ALA	CYS	conflict	UNP P11362
B	584	SER	CYS	conflict	UNP P11362

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 4 3	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Cl 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	87	Total O 87 87	0	0
4	B	58	Total O 58 58	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1



- Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	211.84Å 49.79Å 66.05Å 90.00° 107.40° 90.00°	Depositor
Resolution (Å)	27.82 – 2.31 27.82 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.7 (27.82-2.31) 90.5 (27.82-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 2.31Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.211 , 0.267 0.220 , 0.214	Depositor DCC
$R_{free}$ test set	1311 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.5	Xtrriage
Anisotropy	0.574	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 39.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4722	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/2463	0.44	0/3331
1	B	0.28	0/2212	0.50	0/2985
All	All	0.25	0/4675	0.47	0/6316

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2396	0	2420	32	2
1	B	2171	0	2194	36	0
2	A	7	0	10	0	0
3	A	3	0	0	0	0
4	A	87	0	0	2	0
4	B	58	0	0	2	0
All	All	4722	0	4624	68	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:576:ARG:HH11	1:B:576:ARG:HG2	1.26	0.97
1:A:656:LYS:HG2	1:A:662:LEU:CD2	1.99	0.92
1:A:470:ARG:NH2	4:A:2006:HOH:O	2.04	0.89
1:A:656:LYS:HG2	1:A:662:LEU:HD21	1.55	0.88
1:B:598:LYS:HE2	1:B:761:THR:O	1.77	0.83
1:A:656:LYS:CG	1:A:662:LEU:HD21	2.08	0.83
1:A:664:VAL:HG23	1:A:672:LEU:HD21	1.67	0.75
1:B:664:VAL:HG23	1:B:672:LEU:HD21	1.67	0.74
1:B:598:LYS:HE3	1:B:762:SER:CB	2.19	0.72
1:B:577:ARG:HB3	1:B:696:LEU:CD1	2.20	0.71
1:B:577:ARG:HG3	1:B:578:PRO:HD2	1.74	0.68
1:B:598:LYS:HE3	1:B:762:SER:CA	2.26	0.66
1:B:664:VAL:HG22	1:B:709:LEU:HD23	1.79	0.65
1:B:598:LYS:HE3	1:B:762:SER:HA	1.79	0.64
1:B:576:ARG:HG2	1:B:576:ARG:NH1	2.02	0.64
1:A:598:LYS:HD2	1:A:762:SER:HA	1.80	0.61
1:A:484[B]:LEU:HD12	1:A:492:VAL:HB	1.83	0.61
1:A:570:ARG:HD2	1:A:627:ARG:HD3	1.84	0.60
1:A:656:LYS:CG	1:A:662:LEU:CD2	2.72	0.59
1:B:554:ASP:OD2	4:B:2027:HOH:O	2.16	0.59
1:B:761:THR:O	1:B:762:SER:HB2	2.06	0.56
1:A:650:HIS:O	4:A:2057:HOH:O	2.18	0.55
1:B:501:ASP:HB3	1:B:503:ASP:OD1	2.07	0.54
1:B:598:LYS:CE	1:B:761:THR:O	2.53	0.54
1:B:641:ASP:HB3	1:B:644:LEU:HD22	1.91	0.53
1:B:576:ARG:HH11	1:B:576:ARG:CG	2.10	0.53
1:B:598:LYS:NZ	4:B:2033:HOH:O	1.68	0.52
1:B:498:ILE:HG12	1:B:508:VAL:HG12	1.91	0.51
1:A:656:LYS:HG3	1:A:662:LEU:HD21	1.90	0.51
1:A:475:ARG:NE	1:A:553:GLN:O	2.30	0.51
1:B:534:MET:HG2	1:B:643:GLY:HA2	1.94	0.49
1:A:475:ARG:NH2	1:A:556:PRO:O	2.31	0.49
1:A:607:VAL:HG13	1:A:639[B]:ILE:HD11	1.97	0.47
1:B:730:TYR:CZ	1:B:734:ARG:HD3	2.49	0.47
1:B:576:ARG:NH1	1:B:576:ARG:CG	2.72	0.47
1:A:517:LYS:HE3	1:A:519:ASP:OD1	2.15	0.46
1:A:694:PHE:CZ	1:A:729:LEU:HD13	2.50	0.46
1:B:577:ARG:HB3	1:B:696:LEU:HD13	1.97	0.46
1:B:503:ASP:OD1	1:B:504:LYS:N	2.49	0.46
1:A:500:LEU:HD12	1:A:509:THR:HB	1.99	0.45
1:B:577:ARG:HB3	1:B:696:LEU:HD11	1.97	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:699:SER:HA	1:A:700:PRO:HD3	1.76	0.45
1:A:576:ARG:HG2	1:A:593:GLU:O	2.16	0.45
1:B:461:SER:HA	1:B:464:GLU:O	2.16	0.45
1:B:694:PHE:CZ	1:B:729:LEU:HD13	2.52	0.44
1:A:656:LYS:HE2	1:A:662:LEU:HD11	1.98	0.44
1:A:627:ARG:HG3	1:A:661:ARG:HD3	2.00	0.44
1:B:597:SER:O	1:B:601:VAL:HG23	2.17	0.44
1:B:572:TYR:O	1:B:576:ARG:HG3	2.18	0.43
1:B:465:LEU:HA	1:B:466:PRO:HD3	1.91	0.43
1:A:578:PRO:HA	1:A:579:PRO:HD3	1.90	0.43
1:A:656:LYS:HG2	1:A:662:LEU:HD23	1.92	0.43
1:B:459:GLY:C	1:B:461:SER:H	2.21	0.42
1:A:471:TRP:CD1	1:A:536:LYS:HE2	2.53	0.42
1:A:707:GLU:H	1:A:707:GLU:HG3	1.54	0.42
1:A:551:CYS:HB2	1:A:558:TYR:HB2	2.02	0.42
1:A:653:TYR:HE2	1:A:675:ARG:HE	1.69	0.41
1:A:577:ARG:HA	1:A:578:PRO:HD2	1.67	0.41
1:B:740:VAL:O	1:B:743:GLN:HB2	2.21	0.41
1:A:671:ALA:O	1:A:675:ARG:HA	2.21	0.41
1:A:629:VAL:HG22	1:A:639[B]:ILE:HD13	2.03	0.40
1:B:573:LEU:HD13	1:B:693:ILE:HA	2.02	0.40
1:A:641:ASP:HB3	1:A:644:LEU:HD22	2.04	0.40
1:B:504:LYS:HD2	1:B:507:ARG:CZ	2.51	0.40
1:B:711:LYS:HE3	1:B:711:LYS:HB2	1.72	0.40
1:A:500:LEU:HD23	1:A:500:LEU:HA	1.94	0.40
1:B:676:ILE:O	1:B:676:ILE:HG13	2.22	0.40
1:B:712:LEU:O	1:B:715:GLU:HG2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:507:ARG:NH1	1:A:507:ARG:NH1[2_556]	0.59	1.61
1:A:507:ARG:CZ	1:A:507:ARG:NH1[2_556]	1.80	0.40

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/310 (96%)	285 (96%)	11 (4%)	0	100	100
1	B	265/310 (86%)	260 (98%)	5 (2%)	0	100	100
All	All	561/620 (90%)	545 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/271 (98%)	265 (100%)	1 (0%)	91	96
1	B	239/271 (88%)	237 (99%)	2 (1%)	81	91
All	All	505/542 (93%)	502 (99%)	3 (1%)	86	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	760	LEU
1	B	576	ARG
1	B	760	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PEG	A	1763	-	6,6,6	0.43	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	1763	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	296/310 (95%)	0.36	25 (8%) <b>11</b> <b>15</b>	28, 41, 75, 84	0
1	B	273/310 (88%)	0.56	26 (9%) <b>8</b> <b>11</b>	30, 47, 78, 111	0
All	All	569/620 (91%)	0.46	51 (8%) <b>9</b> <b>12</b>	28, 45, 76, 111	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	673	PHE	12.5
1	B	578	PRO	6.9
1	A	483	PRO	6.6
1	B	593	GLU	6.0
1	B	676	ILE	5.6
1	B	710	PHE	5.6
1	B	646	ARG	4.8
1	B	460	VAL	4.5
1	B	662	LEU	4.4
1	A	656	LYS	4.4
1	B	502	LYS	4.2
1	B	577	ARG	4.0
1	B	675	ARG	4.0
1	B	594	GLN	3.8
1	A	762	SER	3.7
1	A	581	LEU	3.7
1	B	674	ASP	3.6
1	B	598	LYS	3.5
1	B	595	LEU	3.5
1	A	648	ILE	3.5
1	A	463	TYR	3.5
1	A	659	ASN	3.3
1	B	672	LEU	3.3
1	A	491	GLN	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	658	THR	3.2
1	A	482	LYS	3.2
1	A	470	ARG	3.1
1	A	590	ASN	3.1
1	A	579	PRO	3.0
1	B	486	GLU	2.8
1	A	582	GLU	2.8
1	B	740	VAL	2.7
1	A	589	HIS	2.6
1	B	490	GLY	2.6
1	A	657	THR	2.5
1	A	493	VAL	2.5
1	A	503	ASP	2.5
1	B	485	GLY	2.4
1	A	518	SER	2.4
1	B	743	GLN	2.4
1	B	706	VAL	2.3
1	A	462	GLU	2.3
1	A	688	VAL	2.2
1	A	492	VAL	2.2
1	B	707	GLU	2.2
1	B	639	ILE	2.1
1	A	484[A]	LEU	2.1
1	B	459	GLY	2.1
1	A	519	ASP	2.1
1	B	762	SER	2.0
1	A	683	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PEG	A	1763	7/7	0.73	0.22	52,59,64,65	0
3	CL	A	1765	1/1	0.96	0.19	52,52,52,52	0
3	CL	A	1766	1/1	0.96	0.17	60,60,60,60	0
3	CL	A	1764	1/1	0.97	0.18	66,66,66,66	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.