



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2023 – 12:43 AM EDT

PDB ID : 4Y19
Title : immune complex
Authors : Beringer, D.X.; Petersen, J.; Reid, H.H.; Rossjohn, J.
Deposited on : 2015-02-07
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	 8% 91% 5%
2	B	200	 2% 84% 8% 8%
3	C	16	 69% 6% 6% 19%
4	D	210	 9% 91% 6%
5	E	243	 88% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	7	 100%
7	G	2	 50% 50%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1469	952	239	273	5	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	THR	-	expression tag	UNP P01903
A	183	SER	-	expression tag	UNP P01903
A	184	GLY	-	expression tag	UNP P01903
A	185	ASP	-	expression tag	UNP P01903
A	186	ASP	-	expression tag	UNP P01903
A	187	ASP	-	expression tag	UNP P01903
A	188	ASP	-	expression tag	UNP P01903
A	189	LYS	-	expression tag	UNP P01903

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DRB1-4 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	184	1515	957	266	287	5	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P13760
B	0	SER	-	expression tag	UNP P13760
B	191	THR	-	expression tag	UNP P13760
B	192	GLY	-	expression tag	UNP P13760
B	193	GLY	-	expression tag	UNP P13760
B	194	ASP	-	expression tag	UNP P13760
B	195	ASP	-	expression tag	UNP P13760
B	196	ASP	-	expression tag	UNP P13760

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	197	ASP	-	expression tag	UNP P13760
B	198	LYS	-	expression tag	UNP P13760

- Molecule 3 is a protein called Insulin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	13	94	58	18	18	0	0	0


- Molecule 4 is a protein called FS18_alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	205	1594	1009	261	315	9	0	1	0

- Molecule 5 is a protein called FS18_beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	241	1934	1218	341	364	11	0	2	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	F	7	83	46	2	35	0	0	0

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
7	G	2	28	16	2	10	0	0	0

- Molecule 8 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	B	1	7	3	4	0	0
8	B	1	7	3	4	0	0
8	D	1	7	3	4	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	72	72	72	0	0
9	B	67	67	67	0	0
9	C	7	7	7	0	0
9	D	57	57	57	0	0
9	E	103	103	103	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A: 




- Molecule 2: HLA class II histocompatibility antigen, DRB1-4 beta chain

Chain B: 





- Molecule 3: Insulin

Chain C: 



- Molecule 4: FS18_alpha

Chain D: 



- Molecule 5: FS18_beta

Chain E: 



- Molecule 6: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-

D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2
BKAS
MAMA
MANS
MANG
MANT

- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50%  50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	115.76Å 152.74Å 153.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.60 – 2.50 58.91 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.60-2.50) 99.9 (58.91-2.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.51Å)	Xtrriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.160 , 0.196 0.158 , 0.188	Depositor DCC
R_{free} test set	2351 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtrriage
Anisotropy	0.793	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 61.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.012 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7044	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, NAG, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1514	0.69	0/2065
2	B	0.50	0/1555	0.68	0/2112
3	C	0.54	0/94	0.76	0/123
4	D	0.53	0/1633	0.68	0/2219
5	E	0.53	0/1986	0.70	0/2698
All	All	0.52	0/6782	0.69	0/9217

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1469	0	1401	3	0
2	B	1515	0	1422	6	0
3	C	94	0	100	1	0
4	D	1594	0	1500	5	0
5	E	1934	0	1847	12	0
6	F	83	0	70	0	0
7	G	28	0	25	0	0
8	B	14	0	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	D	7	0	2	1	0
9	A	72	0	0	0	0
9	B	67	0	0	0	0
9	C	7	0	0	0	0
9	D	57	0	0	0	0
9	E	103	0	0	0	0
All	All	7044	0	6371	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:116:VAL:HG22	2:B:160:MET:HG2	1.76	0.68
1:A:58:GLY:HA3	3:C:3:LEU:HD21	1.81	0.63
4:D:37:ASP:HB2	4:D:109:TYR:H	1.65	0.61
1:A:118:ASN:HB2	1:A:166:GLU:HB2	1.83	0.60
5:E:101:VAL:HG13	5:E:121:ARG:HD3	1.83	0.59
4:D:38:TYR:CD1	4:D:109:TYR:HD2	2.26	0.54
2:B:103:PRO:HG3	2:B:188:TRP:CZ2	2.45	0.52
4:D:144:ASP:HB3	4:D:147:SER:O	2.09	0.52
5:E:186:GLN:HE21	5:E:187:PRO:HD2	1.75	0.51
8:D:301:MLI:O8	5:E:121:ARG:HD2	2.11	0.50
2:B:163:THR:HG21	2:B:171:TYR:HE2	1.76	0.50
5:E:228:LEU:HD22	5:E:241:PRO:HD2	1.93	0.49
5:E:214:ASN:OD1	5:E:216:ARG:HG2	2.15	0.46
2:B:10:GLN:HB2	2:B:31:PHE:HB2	1.96	0.46
2:B:96:TYR:HA	2:B:179:SER:OG	2.16	0.46
5:E:146:ILE:HG23	5:E:209:ALA:HB1	1.98	0.45
4:D:37:ASP:CB	4:D:109:TYR:H	2.29	0.45
1:A:57:GLN:HG2	5:E:37:GLU:HB3	2.00	0.43
5:E:191:GLN:HB2	5:E:194:LEU:HD22	2.00	0.43
5:E:133:PRO:HD3	5:E:241:PRO:HB3	2.02	0.42
2:B:163:THR:HG21	2:B:171:TYR:CE2	2.55	0.41
5:E:213:GLN:HA	5:E:253:ARG:O	2.21	0.41
5:E:43:ARG:HB3	5:E:53:ILE:HD11	2.02	0.41
4:D:173:TYR:O	4:D:194:ALA:HA	2.22	0.40
5:E:234:TRP:CZ2	5:E:241:PRO:HD3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/189 (94%)	174 (98%)	3 (2%)	0	100	100
2	B	180/200 (90%)	175 (97%)	4 (2%)	1 (1%)	25	43
3	C	11/16 (69%)	11 (100%)	0	0	100	100
4	D	204/210 (97%)	198 (97%)	6 (3%)	0	100	100
5	E	241/243 (99%)	235 (98%)	6 (2%)	0	100	100
All	All	813/858 (95%)	793 (98%)	19 (2%)	1 (0%)	51	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	33	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/173 (94%)	160 (98%)	3 (2%)	59	81
2	B	164/178 (92%)	160 (98%)	4 (2%)	49	74
3	C	9/12 (75%)	7 (78%)	2 (22%)	1	1
4	D	179/189 (95%)	172 (96%)	7 (4%)	32	57
5	E	210/210 (100%)	199 (95%)	11 (5%)	23	44
All	All	725/762 (95%)	698 (96%)	27 (4%)	34	60

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	71	GLU
1	A	95	SER
2	B	39	ARG
2	B	113	ASN
2	B	140	THR
2	B	182	SER
3	C	3	LEU
3	C	10	ARG
4	D	38	TYR
4	D	51	THR
4	D	83	SER
4	D	87	LEU
4	D	131	ASN
4	D	161	GLN
4	D	199	SER
5	E	11	ARG
5	E	19	MET
5	E	25	GLN
5	E	65	THR
5	E	91	LEU
5	E	121	ARG
5	E	186	GLN
5	E	189	LYS
5	E	191	GLN
5	E	204	ARG
5	E	255	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
5	E	186	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	F	1	1,6	14,14,15	0.30	0	17,19,21	0.99	2 (11%)
6	NAG	F	2	6	14,14,15	0.25	0	17,19,21	0.82	2 (11%)
6	BMA	F	3	6	11,11,12	0.30	0	15,15,17	0.76	1 (6%)
6	MAN	F	4	6	11,11,12	0.31	0	15,15,17	0.94	1 (6%)
6	MAN	F	5	6	11,11,12	0.31	0	15,15,17	0.88	1 (6%)
6	MAN	F	6	6	11,11,12	0.40	0	15,15,17	1.01	1 (6%)
6	MAN	F	7	6	11,11,12	0.42	0	15,15,17	0.80	1 (6%)
7	NAG	G	1	7,1	14,14,15	0.31	0	17,19,21	1.16	1 (5%)
7	NAG	G	2	7	14,14,15	0.26	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	0/6/23/26	0/1/1/1
6	BMA	F	3	6	-	0/2/19/22	0/1/1/1
6	MAN	F	4	6	-	0/2/19/22	0/1/1/1
6	MAN	F	5	6	-	2/2/19/22	0/1/1/1
6	MAN	F	6	6	-	0/2/19/22	0/1/1/1
6	MAN	F	7	6	-	1/2/19/22	0/1/1/1
7	NAG	G	1	7,1	-	2/6/23/26	0/1/1/1
7	NAG	G	2	7	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	1	NAG	C1-O5-C5	3.95	117.55	112.19
6	F	6	MAN	C1-O5-C5	3.77	117.30	112.19
6	F	5	MAN	C1-O5-C5	3.07	116.36	112.19
6	F	4	MAN	C1-O5-C5	3.04	116.31	112.19
6	F	1	NAG	O5-C1-C2	-2.56	107.24	111.29
6	F	7	MAN	C1-O5-C5	2.50	115.58	112.19
6	F	3	BMA	C1-O5-C5	2.28	115.28	112.19
6	F	2	NAG	C1-C2-N2	2.28	114.38	110.49
6	F	1	NAG	C1-O5-C5	2.23	115.21	112.19
6	F	2	NAG	O5-C1-C2	-2.01	108.12	111.29

There are no chirality outliers.


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	F	1	NAG	O5-C5-C6-O6
6	F	5	MAN	O5-C5-C6-O6
6	F	1	NAG	C4-C5-C6-O6
6	F	5	MAN	C4-C5-C6-O6
6	F	7	MAN	O5-C5-C6-O6
7	G	1	NAG	C4-C5-C6-O6
7	G	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	MLI	B	201	-	6,6,6	1.28	0	7,7,7	0.97	0
8	MLI	B	202	-	6,6,6	1.30	0	7,7,7	0.91	0
8	MLI	D	301	-	6,6,6	1.25	0	7,7,7	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	MLI	B	201	-	-	2/4/4/4	-
8	MLI	B	202	-	-	4/4/4/4	-
8	MLI	D	301	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	201	MLI	C2-C1-C3-O9
8	B	201	MLI	C2-C1-C3-O8
8	B	202	MLI	C3-C1-C2-O7
8	B	202	MLI	C2-C1-C3-O9
8	B	202	MLI	C3-C1-C2-O6
8	B	202	MLI	C2-C1-C3-O8
8	D	301	MLI	C2-C1-C3-O8
8	D	301	MLI	C2-C1-C3-O9
8	D	301	MLI	C3-C1-C2-O7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	D	301	MLI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/189 (94%)	0.36	16 (8%) 9 9	14, 29, 69, 116	0
2	B	184/200 (92%)	-0.12	4 (2%) 62 65	13, 27, 59, 89	0
3	C	13/16 (81%)	-0.20	0 100 100	19, 21, 47, 56	0
4	D	205/210 (97%)	0.40	18 (8%) 10 10	17, 35, 74, 87	0
5	E	241/243 (99%)	-0.03	1 (0%) 92 93	14, 27, 51, 100	0
All	All	822/858 (95%)	0.14	39 (4%) 31 33	13, 29, 67, 116	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	149	ASP	5.0
1	A	100	ARG	5.0
4	D	199	SER	4.7
5	E	255	ASP	4.3
1	A	129	THR	4.1
4	D	148	SER	3.9
1	A	128	VAL	3.6
1	A	158	GLU	3.5
4	D	183	ARG	3.5
1	A	181	ASP	3.5
1	A	130	THR	3.4
1	A	180	PHE	3.3
4	D	211	ILE	3.3
2	B	113	ASN	3.2
1	A	154	LEU	3.2
4	D	133	GLN	3.0
4	D	204	ALA	2.9
4	D	198	LYS	2.8
4	D	200	ASP	2.8
1	A	157	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	147	SER	2.7
4	D	1	MET	2.7
1	A	178	TRP	2.6
1	A	160	VAL	2.4
4	D	145	SER	2.3
4	D	146	LYS	2.3
4	D	209	ASN	2.3
2	B	164	VAL	2.3
1	A	126	LYS	2.2
4	D	207	PHE	2.2
1	A	99	LEU	2.2
4	D	112	GLY	2.2
2	B	140	THR	2.2
1	A	131	GLY	2.1
4	D	109	TYR	2.1
1	A	123	ARG	2.1
1	A	122	LEU	2.0
2	B	143	VAL	2.0
4	D	197	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MAN	F	5	11/12	0.88	0.20	73,74,76,77	0
6	MAN	F	7	11/12	0.89	0.24	57,63,75,79	0
6	MAN	F	4	11/12	0.93	0.13	48,53,57,66	0
6	MAN	F	6	11/12	0.94	0.17	54,59,64,64	0
7	NAG	G	2	14/15	0.94	0.15	42,50,57,58	0
7	NAG	G	1	14/15	0.96	0.18	32,41,47,52	0
6	NAG	F	1	14/15	0.97	0.14	27,32,38,45	0
6	NAG	F	2	14/15	0.97	0.13	33,36,40,40	0
6	BMA	F	3	11/12	0.98	0.11	40,44,47,52	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	MLI	B	201	7/7	0.83	0.35	48,59,70,71	0
8	MLI	D	301	7/7	0.90	0.23	60,64,71,74	0
8	MLI	B	202	7/7	0.96	0.21	43,46,57,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.