



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 03:48 PM EDT

PDB ID : 4YC3
Title : CDK1/CyclinB1/CKS2 Apo
Authors : Brown, N.R.; Korolchuk, S.; Martin, M.P.; Stanley, W.; Moukhametzianov, R.; Noble, M.E.M.; Endicott, J.A.
Deposited on : 2015-02-19
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
2	B	273	
3	C	84	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	2404	1548	408	439	9	0	2	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P06493
A	-3	PRO	-	expression tag	UNP P06493
A	-2	LEU	-	expression tag	UNP P06493
A	-1	GLY	-	expression tag	UNP P06493
A	0	SER	-	expression tag	UNP P06493

- Molecule 2 is a protein called G2/mitotic-specific cyclin-B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	265	2149	1385	361	387	16	0	1	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	160	GLY	-	expression tag	UNP P14635
B	161	SER	-	expression tag	UNP P14635
B	162	HIS	-	expression tag	UNP P14635
B	163	MET	-	expression tag	UNP P14635
B	166	SER	CYS	engineered mutation	UNP P14635
B	237	SER	CYS	engineered mutation	UNP P14635
B	349	SER	CYS	engineered mutation	UNP P14635


- Molecule 3 is a protein called Cyclin-dependent kinases regulatory subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	72	638	415	113	107	3	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	GLY	-	expression tag	UNP P33552
C	-3	PRO	-	expression tag	UNP P33552
C	-2	LEU	-	expression tag	UNP P33552
C	-1	GLY	-	expression tag	UNP P33552
C	0	SER	-	expression tag	UNP P33552

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	8	6	2	0	0
4	B	1	8	6	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	91	Total	O	0	0
			91	91		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	98	Total	O	0	0
			98	98		
5	C	23	Total	O	0	0
			23	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 1



- Molecule 2: G2/mitotic-specific cyclin-B1



- Molecule 3: Cyclin-dependent kinases regulatory subunit 2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.20Å 70.15Å 156.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.19 – 2.70 52.19 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (52.19-2.70) 98.2 (52.19-2.70)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.211 , 0.260 0.216 , 0.265	Depositor DCC
R_{free} test set	1100 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtrriage
Anisotropy	0.157	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 32.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.032 for k,h,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5419	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.24	13/2457 (0.5%)	1.22	19/3318 (0.6%)
2	B	1.11	8/2194 (0.4%)	1.14	15/2967 (0.5%)
3	C	1.19	5/661 (0.8%)	1.29	10/893 (1.1%)
All	All	1.18	26/5312 (0.5%)	1.20	44/7178 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	GLU	CG-CD	17.89	1.78	1.51
1	A	59[A]	ARG	C-O	15.02	1.51	1.23
1	A	59[B]	ARG	C-O	15.02	1.51	1.23
1	A	57	GLU	CD-OE2	12.95	1.39	1.25
1	A	57	GLU	CD-OE1	12.03	1.38	1.25
3	C	8	TYR	CE1-CZ	-10.56	1.24	1.38
2	B	327	GLU	CD-OE1	9.44	1.36	1.25
1	A	51	GLU	CD-OE2	7.65	1.34	1.25
1	A	209	GLU	CD-OE1	6.46	1.32	1.25
2	B	183	GLU	CD-OE1	6.37	1.32	1.25
2	B	166	SER	CA-CB	6.21	1.62	1.52
3	C	8	TYR	CG-CD2	-6.12	1.31	1.39
1	A	51	GLU	CD-OE1	5.86	1.32	1.25
1	A	207	ASP	CB-CG	5.70	1.63	1.51
1	A	160	TYR	CE1-CZ	5.69	1.46	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	394	THR	CB-CG2	-5.59	1.33	1.52
3	C	63	GLU	CD-OE2	5.38	1.31	1.25
3	C	7	TYR	CE1-CZ	5.33	1.45	1.38
1	A	41	GLU	CG-CD	5.28	1.59	1.51
2	B	360	TRP	CG-CD2	-5.23	1.34	1.43
2	B	298	ARG	CZ-NH1	5.22	1.39	1.33
2	B	430	ALA	N-CA	5.14	1.56	1.46
1	A	57	GLU	CB-CG	5.09	1.61	1.52
3	C	19	TYR	CG-CD1	-5.08	1.32	1.39
2	B	373	GLU	CD-OE2	5.04	1.31	1.25
1	A	38	GLU	CG-CD	5.04	1.59	1.51

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	LEU	CB-CG-CD1	13.94	134.69	111.00
1	A	57	GLU	OE1-CD-OE2	-13.20	107.46	123.30
2	B	180	LEU	CB-CG-CD2	11.87	131.17	111.00
2	B	300	LEU	CB-CG-CD1	11.34	130.28	111.00
2	B	181	GLU	OE1-CD-OE2	-10.68	110.48	123.30
1	A	22	ARG	NE-CZ-NH1	9.71	125.15	120.30
2	B	282	ARG	NE-CZ-NH2	-9.64	115.48	120.30
3	C	73	LEU	CB-CG-CD2	9.29	126.79	111.00
1	A	256	LEU	CB-CG-CD1	8.77	125.91	111.00
2	B	282	ARG	NE-CZ-NH1	8.76	124.68	120.30
3	C	70	ARG	NE-CZ-NH1	8.74	124.67	120.30
1	A	22	ARG	NE-CZ-NH2	-8.50	116.05	120.30
2	B	330	MET	CG-SD-CE	8.41	113.66	100.20
2	B	298	ARG	NE-CZ-NH1	8.40	124.50	120.30
2	B	298	ARG	CG-CD-NE	7.95	128.49	111.80
2	B	298	ARG	NE-CZ-NH2	-7.44	116.58	120.30
1	A	167	LEU	CB-CG-CD2	7.19	123.23	111.00
3	C	26	ARG	NE-CZ-NH1	7.14	123.87	120.30
2	B	192	LEU	CB-CG-CD2	6.91	122.75	111.00
1	A	167	LEU	CA-CB-CG	6.87	131.11	115.30
2	B	233	MET	CG-SD-CE	6.84	111.14	100.20
1	A	247	GLY	N-CA-C	-6.74	96.25	113.10
2	B	206	ASP	CB-CG-OD1	-6.56	112.39	118.30
2	B	290	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	A	151	ARG	NE-CZ-NH2	-6.41	117.09	120.30
1	A	248	SER	N-CA-C	-6.35	93.85	111.00
1	A	37	LEU	CB-CG-CD2	6.16	121.48	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	ASP	CB-CG-OD1	6.16	123.85	118.30
1	A	75	ARG	NE-CZ-NH1	6.11	123.35	120.30
3	C	20	ARG	NE-CZ-NH1	6.07	123.34	120.30
1	A	57	GLU	CB-CG-CD	6.00	130.40	114.20
3	C	71	ARG	NE-CZ-NH1	-5.94	117.33	120.30
3	C	73	LEU	CB-CG-CD1	-5.78	101.17	111.00
3	C	13	PHE	CB-CG-CD1	-5.72	116.80	120.80
3	C	26	ARG	N-CA-CB	5.71	120.87	110.60
2	B	360	TRP	CA-CB-CG	-5.43	103.38	113.70
3	C	26	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	A	236	ASP	CB-CG-OD1	5.37	123.13	118.30
1	A	128	ASP	CB-CG-OD2	-5.34	113.50	118.30
1	A	226	GLU	CA-CB-CG	5.17	124.78	113.40
3	C	70	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	247	GLY	O-C-N	5.15	130.94	122.70
1	A	123	ARG	NE-CZ-NH1	5.08	122.84	120.30
2	B	427	LYS	CB-CG-CD	5.00	124.60	111.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	246	PRO	Peptide
1	A	59[A]	ARG	Mainchain
1	A	59[B]	ARG	Mainchain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2404	0	2447	18	1
2	B	2149	0	2196	28	0
3	C	638	0	623	7	1
4	A	8	0	14	0	0
4	B	8	0	14	0	0
5	A	91	0	0	7	0
5	B	98	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	23	0	0	0	0
All	All	5419	0	5294	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:GLU:CD	1:A:57:GLU:CG	1.78	1.49
2:B:360:TRP:CZ3	2:B:370:TYR:O	2.23	0.91
1:A:35:ILE:O	5:A:401:HOH:O	1.95	0.84
2:B:174[A]:TYR:CE1	5:B:601:HOH:O	2.31	0.83
2:B:360:TRP:HZ3	2:B:370:TYR:O	1.60	0.81
1:A:56:LYS:HE3	2:B:293:ASN:O	1.84	0.76
2:B:174[B]:TYR:CZ	2:B:307:ARG:HD2	2.21	0.75
2:B:174[B]:TYR:CE2	2:B:307:ARG:HD2	2.29	0.68
2:B:297:GLY:O	2:B:298:ARG:HD2	1.97	0.65
2:B:298:ARG:O	2:B:300:LEU:HD13	1.99	0.62
1:A:125:LEU:HD21	1:A:183:THR:HA	1.82	0.60
2:B:416:GLN:HE21	2:B:416:GLN:H	1.48	0.60
1:A:75:ARG:HA	5:A:401:HOH:O	2.03	0.57
1:A:170:ARG:HG3	1:A:170:ARG:HH11	1.70	0.56
1:A:59[B]:ARG:NH1	1:A:66:LEU:O	2.39	0.55
2:B:174[A]:TYR:CZ	5:B:601:HOH:O	2.51	0.54
1:A:289:ASP:HB3	5:A:465:HOH:O	2.07	0.54
1:A:100:MET:HE3	1:A:104:LEU:HD13	1.90	0.53
3:C:27:GLU:N	3:C:27:GLU:OE2	2.42	0.53
1:A:219:ALA:O	1:A:248:SER:HB3	2.08	0.53
3:C:41:GLU:OE1	3:C:41:GLU:N	2.39	0.52
2:B:391:GLN:HE22	2:B:423:GLN:HE22	1.58	0.51
1:A:290:LEU:HD12	5:A:447:HOH:O	2.09	0.51
2:B:360:TRP:CH2	2:B:370:TYR:O	2.63	0.50
1:A:59[B]:ARG:HD3	5:A:450:HOH:O	2.11	0.50
3:C:38:MET:HE2	3:C:43:TRP:HA	1.94	0.49
3:C:73:LEU:HB3	3:C:74:PRO:HD2	1.94	0.49
2:B:274:ASN:HB3	5:B:654:HOH:O	2.13	0.48
2:B:391:GLN:NE2	2:B:423:GLN:HE22	2.11	0.48
3:C:51:SER:HG	3:C:54:TRP:HE1	1.62	0.48
3:C:38:MET:HE3	3:C:42:GLU:HB3	1.95	0.48
2:B:390:ASN:HD22	2:B:418:ASN:HD21	1.61	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ARG:HB2	5:A:412:HOH:O	2.15	0.46
2:B:429:VAL:HG12	2:B:430:ALA:N	2.31	0.46
2:B:345:ALA:HB2	5:B:619:HOH:O	2.15	0.46
1:A:205:HIS:O	1:A:205:HIS:CG	2.68	0.46
2:B:416:GLN:H	2:B:416:GLN:NE2	2.14	0.45
2:B:360:TRP:CZ3	2:B:365:GLN:HA	2.52	0.44
2:B:390:ASN:HD22	2:B:418:ASN:ND2	2.15	0.44
2:B:187:ARG:NH2	2:B:234:GLN:O	2.49	0.44
2:B:274:ASN:CB	5:B:654:HOH:O	2.66	0.44
1:A:100:MET:HE3	1:A:108:TYR:CD2	2.52	0.44
2:B:204:LEU:HD22	2:B:243:LEU:HG	2.01	0.43
3:C:74:PRO:O	3:C:75:LYS:HB2	2.18	0.43
1:A:296:LYS:HD2	1:A:296:LYS:C	2.40	0.42
2:B:300:LEU:N	2:B:301:PRO:HD2	2.34	0.42
2:B:429:VAL:CG1	2:B:430:ALA:N	2.82	0.42
2:B:300:LEU:H	2:B:301:PRO:HD2	1.85	0.42
1:A:75:ARG:NH1	1:A:75:ARG:HG2	2.35	0.41
2:B:350:LEU:O	2:B:354:ILE:HG12	2.21	0.41
2:B:180:LEU:HD13	2:B:300:LEU:HD11	2.03	0.41
1:A:82:PHE:N	5:A:405:HOH:O	2.44	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:THR:O	3:C:45:ARG:NH2[1_455]	1.76	0.44

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/302 (97%)	274 (94%)	19 (6%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	264/273 (97%)	261 (99%)	3 (1%)	0	100	100
3	C	70/84 (83%)	68 (97%)	2 (3%)	0	100	100
All	All	627/659 (95%)	603 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/270 (99%)	244 (91%)	23 (9%)	10	24
2	B	236/242 (98%)	225 (95%)	11 (5%)	26	54
3	C	70/79 (89%)	64 (91%)	6 (9%)	10	24
All	All	573/591 (97%)	533 (93%)	40 (7%)	16	35

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	28	GLN
1	A	37	LEU
1	A	38	GLU
1	A	39	SER
1	A	41	GLU
1	A	71	MET
1	A	75	ARG
1	A	92	ASP
1	A	104	LEU
1	A	123	ARG
1	A	132[A]	GLN
1	A	132[B]	GLN
1	A	160	TYR
1	A	167	LEU
1	A	170	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	205	HIS
1	A	233	SER
1	A	245	LYS
1	A	256	LEU
1	A	279	LYS
1	A	290	LEU
1	A	295	LYS
2	B	166	SER
2	B	241	LYS
2	B	242	MET
2	B	271	VAL
2	B	298	ARG
2	B	300	LEU
2	B	360	TRP
2	B	372	GLU
2	B	373	GLU
2	B	395	LYS
2	B	416	GLN
3	C	4	LYS
3	C	5	GLN
3	C	26	ARG
3	C	61	GLU
3	C	68	LEU
3	C	75	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	72	GLN
1	A	114	GLN
2	B	184	GLN
2	B	318	GLN
2	B	391	GLN
2	B	416	GLN
2	B	418	ASN
3	C	21	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	B	501	-	7,7,7	0.89	0	9,10,10	1.03	1 (11%)
4	MPD	A	301	-	7,7,7	0.64	0	9,10,10	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	B	501	-	-	0/5/5/5	-
4	MPD	A	301	-	-	1/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	MPD	O2-C2-C1	-2.00	101.66	108.08

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	MPD	C2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/302 (97%)	0.09	11 (3%) 41 41	20, 42, 85, 106	0
2	B	265/273 (97%)	-0.29	1 (0%) 92 93	15, 28, 59, 75	0
3	C	72/84 (85%)	0.02	0 100 100	31, 52, 73, 107	0
All	All	632/659 (95%)	-0.08	12 (1%) 66 69	15, 36, 79, 107	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	THR	4.1
1	A	37	LEU	4.0
1	A	38	GLU	4.0
1	A	25	THR	3.1
1	A	13	GLY	2.9
2	B	314	VAL	2.8
1	A	161	THR	2.7
1	A	139	LYS	2.5
1	A	36	ARG	2.4
1	A	1	MET	2.4
1	A	23	HIS	2.4
1	A	39	SER	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	B	501	8/8	0.79	0.25	48,52,54,54	0
4	MPD	A	301	8/8	0.81	0.25	33,49,62,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.