

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 19, 2024 – 11:10 AM EDT

PDB ID	:	4Z7O
Title	:	Integrin alphaIIbbeta3 in complex with AGDV peptide
Authors	:	Lin, F.Y.; Zhu, J.; Springer, T.A.
Deposited on	:	2015-04-07
Resolution	:	2.85  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
IVIETIC	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	164625	1268 (2.88-2.84)
Clashscore	180529	1351 (2.88-2.84)
Ramachandran outliers	177936	1318 (2.88-2.84)
Sidechain outliers	177891	1319 (2.88-2.84)
RSRZ outliers	164620	1269 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	455	% 96%	•
1	С	455	% 96%	•
2	В	469	% 94%	5% •
2	D	469	2% 97%	•
3	Е	221	95%	••



Mol	Chain	Length	Quality of chain	
3	Н	221	94%	•••
4	F	214	% 	·
4	L	214	99%	
5	G	4	25%	
5	Ι	4	25%	
6	J	5	40% 40%	20%
7	Κ	2	100%	
7	Ν	2	100%	
8	М	4	75%	25%



# 2 Entry composition (i)

There are 14 unique types of molecules in this entry. The entry contains 41839 atoms, of which 20230 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Integrin alpha-IIb.

Mol	Chain	Residues			Atom	S			ZeroOcc	AltConf	Trace
1	Δ	455	Total	С	Н	Ν	Ο	S	0	5	0
T	11	400	6886	2237	3369	607	665	8	0	5	U
1	C	453	Total	С	Η	Ν	Ο	$\mathbf{S}$	0	1	0
1	U	400	6796	2212	3315	600	661	8	0	1	0

• Molecule 2 is a protein called Integrin beta-3.

Mol	Chain	Residues			Atom	IS			ZeroOcc	AltConf	Trace
2	В	464	Total 7066	C 2230	Н 3484	N 612	O 707	S 33	0	1	0
2	D	469	Total 7140	C 2248	Н 3528	N 617	0 713	S 34	0	0	0

• Molecule 3 is a protein called Monoclonal antibody 10E5 Fab heavy chain.

Mol	Chain	Residues			Atom	S	ZeroOcc	AltConf	Trace		
3	Е	216	Total 3242	C 1041	Н 1600	N 266	O 329	S 6	0	0	0
3	Н	216	Total 3243	C 1041	Н 1601	N 266	O 329	S 6	0	0	0

• Molecule 4 is a protein called Monoclonal antibody 10E5 Fab light chain.

Mol	Chain	Residues			Atoms	5			ZeroOcc	AltConf	Trace
4	F	914	Total	С	Η	Ν	0	S	0	0	0
4	4 Γ	214	3190	1019	1553	268	341	9	0	0	0
4	т	914	Total	С	Η	Ν	0	S	0	0	0
4		214	3191	1019	1554	268	341	9	0	0	0

• Molecule 5 is a protein called Tetrapeptide ALA-GLY-ASP-VAL.



Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
F	С	4	Total	С	Η	Ν	0	0	0	0
0	5 G	4	45	14	20	4	$\overline{7}$	0	U	U
F	т	4	Total	С	Η	Ν	0	0	0	0
0		4	45	14	20	4	7	0	0	0

• Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues		At	$\mathbf{oms}$			ZeroOcc	AltConf	Trace
6	J	5	Total 118	C 34	Н 57	N 2	O 25	0	0	0

• Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues		At	$\mathbf{oms}$			ZeroOcc	AltConf	Trace
7	K	2	Total	С	Η	Ν	0	0	0	0
(	K 2	Δ	55	16	27	2	10	0	0	0
7	N	0	Total	С	Η	Ν	0	0	0	0
1	í N		55	16	27	2	10	0	0	0

• Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
8	М	4	Total 97	C 28	Н 47	N 2	O 20	0	0	0



• Molecule 9 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
9	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
9	L	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 10 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	4	Total Ca 4 4	0	0
10	С	4	Total Ca 4 4	0	0

• Molecule 11 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	В	3	Total Mn 3 3	0	0
11	D	3	Total Mn 3 3	0	0

• Molecule 12 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
19	В	1	Total	С	Η	Ν	0	0	0	
12	12 D	T	28	8	14	1	5	0		
19	Л	1	Total	С	Η	Ν	Ο	0	0	
12	D	L	28	8	14	1	5	0	0	

• Molecule 13 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	С	1	Total Cl 1 1	0	0

• Molecule 14 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	А	265	Total         O           265         265	0	0
14	В	117	Total         O           117         117	0	0
14	С	84	Total         O           84         84	0	0
14	D	64	$\begin{array}{cc} \text{Total} & \text{O} \\ 64 & 64 \end{array}$	0	0
14	Е	5	$\begin{array}{cc} \text{Total} & \text{O} \\ 5 & 5 \end{array}$	0	0
14	F	9	Total O 9 9	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	Н	22	$\begin{array}{cc} \text{Total} & \text{O} \\ 22 & 22 \end{array}$	0	0
14	L	16	Total O 16 16	0	0
14	G	2	Total O 2 2	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Integrin alpha-IIb



• Molecule 3: Monoclonal antibody 10E5 Fab heavy chain



Chain H:		94%		
E1 D102 D107 P125 V133 C134 C134 C134	T138 T138 1138 1149 149 149 149 149 1487 187 187 187 18219	PRO		
• Molecule 4: N	Ionoclonal antibod	y 10E5 Fab light chain		
Chain F:		99%		!
D1 180 1180 1183 1183 1183 1183 1213 1213 1213 1213				
• Molecule 4: N	Ionoclonal antibod	y 10E5 Fab light chain		
Chain L:		99%		
D1 N157 C214				
• Molecule 5: 7	Cetrapeptide ALA-O	GLY-ASP-VAL		
Chain G:	25%	100%		•
▲408 ● V411				
• Molecule 5: 7	Cetrapeptide ALA-O	GLY-ASP-VAL		
Chain I:	25%	100%		
• 4408 ● V411				
• Molecule 6: a se-(1-4)-2-aceta nose	lpha-D-mannopyra mido-2-deoxy-beta	nose-(1-3)-[alpha-D-ma -D-glucopyranose-(1-4)	unnopyranose-(1-6 )-2-acetamido-2-de	)]beta-D-mannopyrano eoxy-beta-D-glucopyra
Chain J:	40%	40%	20%	I.



• Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:

NAG:

100%



• Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:

100%

#### NAG1 NAG2

 $\bullet \ Molecule \ 8: \ alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose \\ eta-D-glucopyranose \ (1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose \ (1-4)-2-acetamido-2-deoxy-beta-D-glucopyra$ 

Chain M:

75%

25%





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	259.92Å 144.52Å 104.65Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{Posolution} \left( \overset{\circ}{\mathbf{A}} \right)$	49.03 - 2.85	Depositor
Resolution (A)	49.03 - 2.85	EDS
% Data completeness	97.8 (49.03-2.85)	Depositor
(in resolution range)	84.1 (49.03-2.85)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.28 (at 2.86 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.10_2142: ???)	Depositor
P. P.	0.223 , $0.257$	Depositor
$n, n_{free}$	0.236 , $0.261$	DCC
$R_{free}$ test set	2000 reflections $(2.18%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	59.4	Xtriage
Anisotropy	0.461	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, 74.1	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	41839	wwPDB-VP
Average B, all atoms $(Å^2)$	128.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAG, CA, MN, BMA, MAN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.29	0/3629	0.44	0/4944
1	С	0.26	0/3581	0.42	0/4880
2	В	0.26	0/3651	0.43	0/4950
2	D	0.26	0/3678	0.41	0/4986
3	Е	0.25	0/1684	0.42	0/2305
3	Н	0.24	0/1684	0.43	0/2305
4	F	0.25	0/1673	0.41	0/2269
4	L	0.25	0/1673	0.42	0/2269
5	G	0.19	0/24	0.45	0/30
5	Ι	0.25	0/24	0.48	0/30
All	All	0.26	0/21301	0.42	0/28968

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3517	3369	3370	10	0
1	С	3481	3315	3315	8	0
2	В	3582	3484	3495	10	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	3612	3528	3526	6	0
3	Е	1642	1600	1600	3	0
3	Н	1642	1601	1600	6	0
4	F	1637	1553	1553	1	0
4	L	1637	1554	1553	2	0
5	G	25	20	20	0	0
5	Ι	25	20	20	0	0
6	J	61	57	52	1	0
7	K	28	27	25	0	0
7	Ν	28	27	25	0	0
8	М	50	47	43	0	0
9	А	5	0	0	0	0
9	С	5	0	0	0	0
9	L	5	0	0	0	0
10	А	4	0	0	0	0
10	С	4	0	0	0	0
11	В	3	0	0	0	0
11	D	3	0	0	0	0
12	В	14	14	13	0	0
12	D	14	14	13	0	0
13	С	1	0	0	0	0
14	А	265	0	0	5	1
14	В	117	0	0	1	0
14	С	84	0	0	4	1
14	D	64	0	0	1	0
14	Е	5	0	0	0	0
14	F	9	0	0	0	0
14	G	2	0	0	0	0
14	Н	22	0	0	1	0
14	L	16	0	0	1	0
All	All	21609	20230	20223	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:GLN:OE1	14:A:601:HOH:O	2.13	0.67
1:A:15[B]:ASN:OD1	14:A:602:HOH:O	2.15	0.65
4:L:157:ASN:OD1	14:L:401:HOH:O	2.14	0.64



	A 4 arra 0	Interatomic	Clash		
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)		
2:B:320:ASN:ND2	6:J:1:NAG:O7	2.37	0.56		
1:A:142:GLU:OE1	14:A:603:HOH:O	2.18	0.56		
1:A:278[B]:HIS:NE2	14:A:614:HOH:O	2.34	0.53		
1:C:24:ASP:OD1	1:C:25:PHE:N	2.41	0.52		
1:C:428:ASP:OD2	14:C:601:HOH:O	2.19	0.50		
3:H:107:ASP:O	14:H:301:HOH:O	2.20	0.49		
1:A:224:ASP:OD1	1:A:225:SER:N	2.40	0.48		
3:H:149:LYS:NZ	3:H:177:GLN:OE1	2.38	0.48		
2:B:133:ASN:OD1	2:B:137:LYS:HG3	2.14	0.47		
1:C:373:ASP:O	14:C:602:HOH:O	2.20	0.47		
1:A:77:ARG:NH2	3:H:102:ASP:OD2	2.43	0.47		
2:B:159:LYS:O	2:B:161:VAL:N	2.47	0.47		
2:B:235:LYS:NZ	2:B:276:GLY:O	2.48	0.47		
2:B:138:LEU:O	2:B:141:GLN:N	2.49	0.46		
1:C:107:CYS:HA	1:C:130:CYS:HA	1.98	0.46		
1:C:364:GLY:N	14:C:602:HOH:O	2.43	0.46		
2:B:138:LEU:HA	2:B:341:LEU:CD1	2.46	0.45		
4:F:211:ARG:O	4:F:213:GLU:N	2.47	0.45		
2:D:127:ASP:N	2:D:127:ASP:OD1	2.50	0.45		
3:H:165:LEU:HD21	3:H:187:VAL:HG21	2.00	0.44		
1:C:227:ASN:ND2	14:C:613:HOH:O	2.49	0.44		
2:D:217:ASP:OD2	2:D:255:HIS:NE2	2.50	0.44		
1:A:53:VAL:HG21	1:A:106:ALA:CB	2.47	0.44		
1:A:160:PHE:O	14:A:605:HOH:O	2.21	0.44		
2:B:138:LEU:O	2:B:140:THR:N	2.51	0.43		
2:D:288:ASP:OD1	2:D:289:TYR:N	2.43	0.43		
1:C:35:ILE:O	1:C:56:CYS:N	2.46	0.43		
3:E:43:GLN:OE1	3:E:43:GLN:N	2.48	0.43		
3:E:125:PRO:HB2	3:E:148:VAL:HG13	2.01	0.43		
3:H:125:PRO:HB2	3:H:148:VAL:HG13	2.01	0.43		
1:C:2:ASN:OD1	1:C:2:ASN:N	2.51	0.42		
2:D:310:VAL:HB	2:D:314:VAL:HG23	2.02	0.42		
2:B:133:ASN:OD1	2:B:137:LYS:HE3	2.20	0.42		
2:D:150:ARG:NH1	14:D:2111:HOH:O	2.44	0.42		
2:B:334:SER:O	2:B:336:ASP:N	2.49	0.42		
2:B:191:LYS:NZ	14:B:2119:HOH:O	2.49	0.41		
1:A:107:CYS:HA	1:A:130:CYS:HA	2.02	0.41		
2:D:319:GLN:HA	2:D:330:VAL:HG21	2.03	0.40		
3:H:134:CYS:HB2	4:L:214:CYS:OXT	2.21	0.40		
3:E:133:VAL:O	3:E:134:CYS:SG	2.80	0.40		

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
14:A:833:HOH:O	14:C:650:HOH:O[1_554]	2.14	0.06	

metry operator and encoded unit-cell translations to be applied.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	458/455~(101%)	432 (94%)	26 (6%)	0	100	100
1	С	452/455~(99%)	429 (95%)	23~(5%)	0	100	100
2	В	463/469~(99%)	435 (94%)	26 (6%)	2~(0%)	30	49
2	D	467/469~(100%)	438 (94%)	28 (6%)	1 (0%)	44	63
3	Ε	212/221~(96%)	198 (93%)	13 (6%)	1 (0%)	25	43
3	Н	212/221~(96%)	202 (95%)	10 (5%)	0	100	100
4	F	212/214~(99%)	203~(96%)	9 (4%)	0	100	100
4	L	212/214~(99%)	204 (96%)	8 (4%)	0	100	100
5	G	2/4~(50%)	2 (100%)	0	0	100	100
5	Ι	2/4~(50%)	2 (100%)	0	0	100	100
All	All	2692/2726~(99%)	2545 (94%)	143 (5%)	4 (0%)	48	69

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	Ε	133	VAL
2	В	139	ALA
2	D	374	CYS
2	В	157	VAL



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	368/363~(101%)	362~(98%)	6(2%)	58	79
1	$\mathbf{C}$	362/363~(100%)	359~(99%)	3~(1%)	79	90
2	В	411/415~(99%)	405~(98%)	6 (2%)	60	81
2	D	415/415~(100%)	411 (99%)	4 (1%)	73	87
3	Ε	187/190~(98%)	187 (100%)	0	100	100
3	Η	187/190~(98%)	187 (100%)	0	100	100
4	F	188/188~(100%)	187~(100%)	1 (0%)	86	94
4	L	188/188 (100%)	187 (100%)	1 (0%)	86	94
5	G	2/2~(100%)	2~(100%)	0	100	100
5	Ι	2/2~(100%)	2 (100%)	0	100	100
All	All	2310/2316~(100%)	2289~(99%)	21 (1%)	75	88

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	23	LEU
1	А	166	TYR
1	А	190	TYR
1	А	252	GLU
1	А	267	VAL
1	А	288	TYR
2	В	132	GLN
2	В	136	THR
2	В	215	ASN
2	В	337	SER
2	В	343	LEU
2	В	447	ARG
1	С	23	LEU
1	С	166	TYR
1	С	288	TYR
2	D	127	ASP



Continued from previous page...

Mol	Chain	Res	Type
2	D	158	ASP
2	D	215	ASN
2	D	365	GLU
4	F	214	CYS
4	L	214	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	В	132	GLN
2	D	280	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Chain	Chain	Chain	Chain	Chain	Chain	Dec	Tink	Bo	Bond lengths			Bond angles		
	туре	Unam			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2							
6	NAG	J	1	2,6	14,14,15	0.40	0	17,19,21	0.67	1 (5%)							
6	NAG	J	2	6	14,14,15	0.23	0	17,19,21	0.49	0							
6	BMA	J	3	6	11,11,12	0.64	0	15,15,17	0.96	0							
6	MAN	J	4	6	11,11,12	0.48	0	$15,\!15,\!17$	0.89	1 (6%)							
6	MAN	J	5	6	11,11,12	0.64	0	15,15,17	1.26	2 (13%)							
7	NAG	K	1	7,2	14,14,15	0.48	0	17,19,21	0.47	0							



Mol Type		Chain	Chain	Chain	Chain	Chain Res	Ros Link	Bo	Bond lengths			Bond angles		
WIOI	Moi Type	n res		es Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2				
7	NAG	K	2	7	14,14,15	0.17	0	$17,\!19,\!21$	0.46	0				
8	NAG	М	1	2,8	14,14,15	0.55	0	$17,\!19,\!21$	0.51	0				
8	NAG	М	2	8	$14,\!14,\!15$	0.34	0	$17,\!19,\!21$	0.54	0				
8	BMA	М	3	8	$11,\!11,\!12$	0.71	0	$15,\!15,\!17$	0.78	0				
8	MAN	М	4	8	11,11,12	0.53	0	$15,\!15,\!17$	1.04	2 (13%)				
7	NAG	N	1	7,2	14,14,15	0.41	0	17,19,21	0.48	0				
7	NAG	N	2	7	14,14,15	0.28	0	17,19,21	0.44	0				

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	J	1	2,6	-	4/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
6	BMA	J	3	6	-	2/2/19/22	0/1/1/1
6	MAN	J	4	6	-	0/2/19/22	0/1/1/1
6	MAN	J	5	6	-	2/2/19/22	0/1/1/1
7	NAG	Κ	1	7,2	-	0/6/23/26	0/1/1/1
7	NAG	К	2	7	-	2/6/23/26	0/1/1/1
8	NAG	М	1	2,8	-	0/6/23/26	0/1/1/1
8	NAG	М	2	8	-	0/6/23/26	0/1/1/1
8	BMA	М	3	8	-	0/2/19/22	0/1/1/1
8	MAN	М	4	8	-	0/2/19/22	0/1/1/1
7	NAG	N	1	7,2	-	0/6/23/26	0/1/1/1
7	NAG	N	2	7	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
6	J	5	MAN	C1-O5-C5	3.85	117.35	112.19
8	М	4	MAN	C1-O5-C5	2.88	116.05	112.19
6	J	5	MAN	O2-C2-C3	-2.18	105.63	110.15
6	J	4	MAN	C1-O5-C5	2.13	115.04	112.19
8	М	4	MAN	O2-C2-C3	-2.10	105.79	110.15
6	J	1	NAG	C1-O5-C5	2.02	114.90	112.19

There are no chirality outliers.



Mol	Chain	Res	Type	Atoms
7	Κ	2	NAG	O5-C5-C6-O6
6	J	3	BMA	O5-C5-C6-O6
6	J	5	MAN	O5-C5-C6-O6
7	Κ	2	NAG	C4-C5-C6-O6
6	J	5	MAN	C4-C5-C6-O6
6	J	3	BMA	C4-C5-C6-O6
6	J	1	NAG	C3-C2-N2-C7
6	J	1	NAG	C1-C2-N2-C7
6	J	1	NAG	C4-C5-C6-O6
7	N	2	NAG	O5-C5-C6-O6
6	J	1	NAG	O5-C5-C6-O6

All (11) torsion outliers are listed below:

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	J	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















### 5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 15 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Ch		Chain	hain Res	Link	Bond lengths			Bond angles		
Moi Type	Unam	Counts			RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
12	NAG	В	2004	2	14,14,15	0.55	0	17,19,21	0.75	1 (5%)
12	NAG	D	2004	2	14,14,15	0.38	0	17,19,21	0.60	0
9	SO4	А	501	-	4,4,4	0.24	0	6,6,6	0.04	0
9	SO4	С	501	-	4,4,4	0.24	0	$6,\!6,\!6$	0.05	0
9	SO4	L	301	-	4,4,4	0.23	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	NAG	D	2004	2	-	2/6/23/26	0/1/1/1
12	NAG	В	2004	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
12	В	2004	NAG	C1-O5-C5	2.55	115.60	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	В	2004	NAG	O5-C5-C6-O6
12	D	2004	NAG	O5-C5-C6-O6
12	D	2004	NAG	C4-C5-C6-O6
12	В	2004	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	455/455~(100%)	-0.33	4 (0%) 81 79	34, 74, 108, 207	5(1%)
1	С	453/455~(99%)	0.00	3 (0%) 84 83	65, 105, 149, 194	1 (0%)
2	В	464/469~(98%)	-0.16	3 (0%) 85 84	52, 114, 200, 267	1 (0%)
2	D	469/469~(100%)	0.22	10 (2%) 63 60	80, 141, 201, 282	0
3	E	216/221~(97%)	0.31	0 100 100	121, 195, 279, 303	0
3	Н	216/221~(97%)	0.07	1 (0%) 87 86	80, 145, 210, 243	0
4	F	214/214~(100%)	0.28	3 (1%) 73 70	125, 187, 277, 329	0
4	L	214/214~(100%)	-0.11	0 100 100	85, 130, 165, 251	0
5	G	4/4~(100%)	0.78	1 (25%) 2 2	66, 89, 101, 102	0
5	Ι	4/4~(100%)	1.33	1 (25%) 2 2	101, 112, 116, 138	0
All	All	2709/2726 (99%)	0.00	26 (0%) 79 78	34, 123, 227, 329	7 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	454	VAL	5.4
2	D	151	ILE	3.8
1	А	455	LYS	3.6
2	D	375	LEU	3.3
1	С	1	LEU	3.3
4	F	180	THR	3.1
2	D	142	MET	3.1
1	С	148	GLY	3.0
5	Ι	408	ALA	3.0
1	С	453	VAL	2.9
4	F	181	LEU	2.7
2	В	128	LEU	2.6
2	D	118	MET	2.6



		1	1 0	
Mol	Chain	Res	Type	RSRZ
5	G	408	ALA	2.6
2	В	129	TRP	2.5
1	А	226	SER	2.5
1	А	90[A]	ARG	2.4
2	D	137	LYS	2.3
2	D	129	TRP	2.2
2	D	334	SER	2.2
4	F	193	THR	2.1
2	D	153	PHE	2.1
2	В	466	TRP	2.1
2	D	374	CYS	2.1
2	D	376	ASN	2.0
3	Н	133	VAL	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

SUGAR-RSR INFOmissingINFO

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
12	NAG	D	2004	14/15	0.64	0.11	114,142,170,175	0
11	MN	D	2002	1/1	0.65	0.23	431,431,431,431	0
12	NAG	В	2004	14/15	0.73	0.10	118,139,166,167	0
11	MN	В	2002	1/1	0.73	0.19	361,361,361,361	0
9	SO4	L	301	5/5	0.80	0.09	151,154,159,161	0
9	SO4	С	501	5/5	0.86	0.20	184,193,196,198	0
9	SO4	А	501	5/5	0.90	0.14	118,124,133,142	0
13	CL	С	502	1/1	0.91	0.11	85,85,85,85	0
11	MN	D	2003	1/1	0.95	0.10	216,216,216,216	0
10	CA	С	504	1/1	0.95	0.07	168,168,168,168	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
10	CA	С	506	1/1	0.98	0.06	108,108,108,108	0
10	CA	С	503	1/1	0.98	0.08	197,197,197,197	0
11	MN	D	2001	1/1	0.98	0.05	121,121,121,121	0
10	CA	А	504	1/1	0.98	0.04	57,57,57,57	0
10	CA	А	503	1/1	0.99	0.05	80,80,80,80	0
11	MN	В	2001	1/1	0.99	0.04	57,57,57,57	0
10	CA	А	502	1/1	0.99	0.03	77,77,77,77	0
11	MN	В	2003	1/1	0.99	0.03	90,90,90,90	0
10	CA	С	505	1/1	0.99	0.03	99,99,99,99	0
10	CA	А	505	1/1	1.00	0.02	66,66,66,66	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

