



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 11:53 AM EDT

PDB ID : 5F9O
Title : Crystal structure of broadly neutralizing VH1-46 germline-derived CD4-binding site-directed antibody CH235.09 in complex with HIV-1 clade A/E 93TH057 gp120
Authors : Joyce, M.G.; Mascola, J.R.; Kwong, P.D.
Deposited on : 2015-12-10
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	352	
2	H	234	
3	L	213	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EPE	G	513	-	-	X	-
5	EPE	G	514	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	G	341	2671	1678	464	506	23	0	0	0


- Molecule 2 is a protein called CH235.9 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	220	1661	1047	287	316	11	0	0	0

- Molecule 3 is a protein called CH235.09 Light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	210	1635	1022	285	323	5	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	G	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	G	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 6 is POLYETHYLENE GLYCOL (N=34) (three-letter code: 15P) (formula: $C_{69}H_{140}O_{35}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	G	1	Total	C	O	0	0
			23	15	8		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	G	191	Total	O	0	0
			191	191		
7	H	141	Total	O	0	0
			141	141		
7	L	128	Total	O	0	0
			128	128		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



- Molecule 2: CH235.9 Heavy chain



- Molecule 3: CH235.09 Light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.53Å 67.80Å 225.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.81 – 1.86 35.81 – 1.86	Depositor EDS
% Data completeness (in resolution range)	89.2 (35.81-1.86) 85.5 (35.81-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.85Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.204 , 0.221 0.213 , 0.229	Depositor DCC
R_{free} test set	3557 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtrriage
Anisotropy	0.192	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6648	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, NAG, 15P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.41	0/2727	0.48	0/3701
2	H	0.31	0/1702	0.54	0/2313
3	L	0.36	0/1670	0.51	0/2267
All	All	0.37	0/6099	0.51	0/8281

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	123	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2671	0	2610	26	1
2	H	1661	0	1624	11	0
3	L	1635	0	1596	9	0
4	G	168	0	156	9	1
5	G	30	0	35	0	18
6	G	23	0	29	0	0
7	G	191	0	0	8	0
7	H	141	0	0	1	0
7	L	128	0	0	4	0
All	All	6648	0	6050	47	19

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:325:ASP:OD1	7:G:601:HOH:O	1.88	0.91
3:L:141:ARG:NH1	7:L:301:HOH:O	2.10	0.84
1:G:106:GLU:OE2	7:G:602:HOH:O	1.98	0.81
1:G:442:LYS:NZ	7:G:603:HOH:O	2.18	0.77
2:H:69:SER:OG	2:H:84:ARG:NH1	2.18	0.75
4:G:507:NAG:H3	4:G:507:NAG:O7	1.92	0.69
3:L:5:THR:OG1	7:L:302:HOH:O	2.12	0.68
2:H:66:ASP:O	2:H:84:ARG:NH2	2.31	0.62
1:G:278:THR:HG22	1:G:280:ASN:H	1.65	0.61
4:G:505:NAG:H82	4:G:505:NAG:O3	2.01	0.60
1:G:409:GLY:N	7:G:607:HOH:O	2.36	0.57
4:G:502:NAG:H3	4:G:502:NAG:H82	1.87	0.56
1:G:278:THR:HG23	3:L:92:ASN:HA	1.87	0.55
4:G:508:NAG:H83	4:G:508:NAG:H3	1.88	0.54
1:G:465:ASN:OD1	1:G:465:ASN:N	2.41	0.54
3:L:141:ARG:NH2	7:L:308:HOH:O	2.41	0.53
1:G:88:ASN:HD22	4:G:501:NAG:H82	1.75	0.52
1:G:448:ASN:OD1	7:G:604:HOH:O	2.20	0.50
2:H:72:ARG:HD3	7:H:353:HOH:O	2.13	0.49
1:G:276:ASN:HB2	4:G:502:NAG:H83	1.95	0.48
1:G:448:ASN:OD1	4:G:512:NAG:H2	2.14	0.48
1:G:325:ASP:O	7:G:605:HOH:O	2.20	0.47
3:L:128:THR:OG1	7:L:303:HOH:O	2.13	0.47
1:G:112:TRP:CE3	1:G:116:LEU:HD11	2.50	0.46
1:G:205:CYS:HB2	7:G:761:HOH:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:126:LYS:HZ1	2:H:184:LEU:HD13	1.81	0.45
2:H:218:LYS:NZ	3:L:122:GLU:OE2	2.40	0.44
1:G:112:TRP:CE3	1:G:116:LEU:CD1	3.01	0.43
1:G:346:THR:HG23	1:G:359:ILE:HB	2.00	0.43
4:G:507:NAG:O7	4:G:507:NAG:C3	2.64	0.43
1:G:357:LYS:NZ	1:G:462:ASN:O	2.50	0.43
1:G:278:THR:HG22	1:G:279:ASN:N	2.34	0.42
2:H:34:ILE:HD13	2:H:79:LEU:CD2	2.49	0.42
2:H:3:ARG:HH11	2:H:4:LEU:H	1.67	0.42
3:L:13:VAL:HG21	3:L:19:VAL:HG23	2.00	0.42
2:H:64:PHE:HB3	2:H:68:VAL:HG23	2.01	0.42
1:G:448:ASN:OD1	4:G:512:NAG:C2	2.67	0.42
2:H:34:ILE:HD13	2:H:79:LEU:CD1	2.50	0.41
2:H:204:ILE:HG12	2:H:219:ARG:HB3	2.02	0.41
1:G:124:GLY:HA2	7:G:729:HOH:O	2.20	0.41
1:G:268:GLU:HB3	1:G:269:GLU:H	1.56	0.41
1:G:112:TRP:CZ3	1:G:116:LEU:HD11	2.55	0.41
1:G:278:THR:CG2	3:L:91:TYR:O	2.69	0.41
1:G:278:THR:HG21	3:L:91:TYR:O	2.21	0.41
2:H:34:ILE:HD13	2:H:79:LEU:HD21	2.01	0.41
1:G:123:THR:O	1:G:123:THR:CG2	2.65	0.40
1:G:123:THR:O	1:G:123:THR:HG22	2.21	0.40

All (19) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:G:513:EPE:O1S	5:G:514:EPE:O3S[3_554]	0.03	2.17
5:G:513:EPE:S	5:G:514:EPE:S[3_554]	0.07	2.13
5:G:513:EPE:C10	5:G:514:EPE:C10[3_554]	0.11	2.09
5:G:513:EPE:O8	5:G:514:EPE:O8[3_554]	0.15	2.05
5:G:513:EPE:O2S	5:G:514:EPE:O2S[3_554]	0.23	1.97
5:G:513:EPE:C3	5:G:514:EPE:C5[3_554]	0.24	1.96
5:G:513:EPE:O3S	5:G:514:EPE:O1S[3_554]	0.28	1.92
5:G:513:EPE:C6	5:G:514:EPE:C2[3_554]	0.34	1.86
5:G:513:EPE:C9	5:G:514:EPE:C9[3_554]	0.46	1.74
5:G:513:EPE:N1	5:G:514:EPE:N1[3_554]	0.50	1.70
5:G:513:EPE:N4	5:G:514:EPE:N4[3_554]	0.52	1.68
5:G:513:EPE:C5	5:G:514:EPE:C3[3_554]	0.61	1.59
5:G:513:EPE:C8	5:G:514:EPE:C8[3_554]	0.62	1.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:G:513:EPE:C2	5:G:514:EPE:C6[3_554]	0.69	1.51
5:G:513:EPE:C7	5:G:514:EPE:C7[3_554]	0.72	1.48
5:G:513:EPE:S	5:G:514:EPE:O1S[3_554]	1.29	0.91
5:G:513:EPE:O2S	5:G:514:EPE:S[3_554]	1.32	0.88
1:G:408:LYS:NZ	4:G:512:NAG:O4[3_644]	2.09	0.11
5:G:513:EPE:O2S	5:G:514:EPE:O1S[3_554]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	335/352 (95%)	327 (98%)	8 (2%)	0	100	100
2	H	216/234 (92%)	211 (98%)	5 (2%)	0	100	100
3	L	208/213 (98%)	203 (98%)	5 (2%)	0	100	100
All	All	759/799 (95%)	741 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	305/310 (98%)	291 (95%)	14 (5%)	23	9
2	H	183/196 (93%)	173 (94%)	10 (6%)	18	6
3	L	183/185 (99%)	171 (93%)	12 (7%)	14	3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	671/691 (97%)	635 (95%)	36 (5%)	18 6

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	44	VAL
1	G	46	LYS
1	G	87	GLU
1	G	117	GLN
1	G	123	THR
1	G	126	SER
1	G	226	LEU
1	G	231	LYS
1	G	244	SER
1	G	336	THR
1	G	407	MET
1	G	408	LYS
1	G	423	ILE
1	G	465	ASN
2	H	3	ARG
2	H	43	GLN
2	H	72	ARG
2	H	79	LEU
2	H	87	ARG
2	H	116	VAL
2	H	126	LYS
2	H	141	SER
2	H	159	VAL
2	H	208	ASN
3	L	11	LEU
3	L	14	SER
3	L	24	ARG
3	L	33	LEU
3	L	39	LYS
3	L	45	ARG
3	L	74	THR
3	L	80	SER
3	L	99	GLN
3	L	104	GLU
3	L	153	LEU
3	L	184	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	444	ASN
3	L	159	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	15P	G	515	-	22,22,103	0.85	0	21,21,102	0.27	0
4	NAG	G	505	1	14,14,15	1.02	1 (7%)	17,19,21	1.08	1 (5%)
4	NAG	G	509	1	14,14,15	0.45	0	17,19,21	0.56	0
4	NAG	G	510	1	14,14,15	0.29	0	17,19,21	0.56	0
4	NAG	G	503	1	14,14,15	0.28	0	17,19,21	0.55	0
4	NAG	G	506	1	14,14,15	0.23	0	17,19,21	0.56	0
4	NAG	G	507	1	14,14,15	0.29	0	17,19,21	0.56	0
4	NAG	G	501	1	14,14,15	0.29	0	17,19,21	0.56	0
4	NAG	G	508	1	14,14,15	0.17	0	17,19,21	1.28	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EPE	G	513	5	15,15,15	2.78	4 (26%)	19,20,20	1.20	2 (10%)
5	EPE	G	514	5	15,15,15	2.75	4 (26%)	19,20,20	1.66	4 (21%)
4	NAG	G	502	1	14,14,15	0.30	0	17,19,21	0.56	0
4	NAG	G	504	1	14,14,15	0.31	0	17,19,21	0.52	0
4	NAG	G	511	1	14,14,15	0.19	0	17,19,21	0.44	0
4	NAG	G	512	1	14,14,15	0.44	0	17,19,21	1.29	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	15P	G	515	-	-	0/20/20/101	-
4	NAG	G	505	1	-	4/6/23/26	0/1/1/1
4	NAG	G	509	1	-	2/6/23/26	0/1/1/1
4	NAG	G	510	1	-	0/6/23/26	0/1/1/1
4	NAG	G	503	1	-	5/6/23/26	0/1/1/1
4	NAG	G	506	1	-	0/6/23/26	0/1/1/1
4	NAG	G	507	1	-	3/6/23/26	0/1/1/1
4	NAG	G	501	1	-	4/6/23/26	0/1/1/1
4	NAG	G	508	1	-	6/6/23/26	0/1/1/1
5	EPE	G	513	5	-	1/9/19/19	0/1/1/1
5	EPE	G	514	5	-	4/9/19/19	0/1/1/1
4	NAG	G	502	1	-	5/6/23/26	0/1/1/1
4	NAG	G	504	1	-	2/6/23/26	0/1/1/1
4	NAG	G	511	1	-	2/6/23/26	0/1/1/1
4	NAG	G	512	1	-	2/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	513	EPE	C10-S	-9.38	1.64	1.77
5	G	514	EPE	C10-S	-8.85	1.65	1.77
5	G	514	EPE	O2S-S	3.60	1.55	1.45
5	G	514	EPE	O1S-S	-3.33	1.35	1.45
4	G	505	NAG	C1-C2	3.23	1.56	1.52
5	G	513	EPE	O2S-S	-3.19	1.36	1.45
5	G	513	EPE	O1S-S	-2.97	1.36	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	514	EPE	O3S-S	-2.74	1.36	1.47
5	G	513	EPE	O3S-S	2.60	1.57	1.47

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	512	NAG	C1-O5-C5	4.46	118.16	112.19
5	G	514	EPE	O2S-S-C10	4.32	113.25	106.73
4	G	508	NAG	C2-N2-C7	3.94	128.17	122.90
5	G	513	EPE	O2S-S-C10	3.68	112.29	106.73
4	G	505	NAG	C2-N2-C7	3.21	127.20	122.90
5	G	514	EPE	O1S-S-C10	2.76	110.89	106.73
4	G	508	NAG	C1-C2-N2	2.51	114.38	110.43
5	G	514	EPE	C9-N1-C2	-2.11	105.61	111.24
5	G	514	EPE	O2S-S-O1S	-2.01	107.28	113.82
5	G	513	EPE	O3S-S-C10	2.01	109.94	106.00

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	501	NAG	C8-C7-N2-C2
4	G	501	NAG	O7-C7-N2-C2
4	G	502	NAG	C1-C2-N2-C7
4	G	502	NAG	C8-C7-N2-C2
4	G	502	NAG	O7-C7-N2-C2
4	G	503	NAG	C1-C2-N2-C7
4	G	503	NAG	C8-C7-N2-C2
4	G	503	NAG	O7-C7-N2-C2
5	G	514	EPE	C10-C9-N1-C6
4	G	502	NAG	C4-C5-C6-O6
4	G	502	NAG	O5-C5-C6-O6
4	G	505	NAG	O5-C5-C6-O6
4	G	507	NAG	O5-C5-C6-O6
4	G	501	NAG	O5-C5-C6-O6
4	G	501	NAG	C4-C5-C6-O6
4	G	509	NAG	O5-C5-C6-O6
4	G	508	NAG	C4-C5-C6-O6
4	G	503	NAG	O5-C5-C6-O6
4	G	503	NAG	C4-C5-C6-O6
4	G	507	NAG	C4-C5-C6-O6
4	G	505	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	G	509	NAG	C4-C5-C6-O6
4	G	508	NAG	O5-C5-C6-O6
4	G	505	NAG	C8-C7-N2-C2
4	G	505	NAG	O7-C7-N2-C2
4	G	508	NAG	C8-C7-N2-C2
4	G	508	NAG	O7-C7-N2-C2
4	G	512	NAG	O5-C5-C6-O6
4	G	512	NAG	C4-C5-C6-O6
4	G	511	NAG	C4-C5-C6-O6
5	G	514	EPE	C9-C10-S-O3S
4	G	507	NAG	C3-C2-N2-C7
4	G	504	NAG	C4-C5-C6-O6
4	G	511	NAG	O5-C5-C6-O6
4	G	508	NAG	C1-C2-N2-C7
5	G	514	EPE	C9-C10-S-O1S
5	G	514	EPE	C9-C10-S-O2S
4	G	504	NAG	O5-C5-C6-O6
5	G	513	EPE	N4-C7-C8-O8
4	G	508	NAG	C3-C2-N2-C7

There are no ring outliers.

8 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	505	NAG	1	0
4	G	507	NAG	2	0
4	G	501	NAG	1	0
4	G	508	NAG	1	0
5	G	513	EPE	0	18
5	G	514	EPE	0	18
4	G	502	NAG	2	0
4	G	512	NAG	2	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	341/352 (96%)	0.57	28 (8%) 19 20	26, 40, 74, 146	0
2	H	220/234 (94%)	0.39	15 (6%) 25 25	25, 36, 60, 122	0
3	L	210/213 (98%)	0.43	8 (3%) 44 47	25, 40, 61, 85	0
All	All	771/799 (96%)	0.48	51 (6%) 26 27	25, 39, 65, 146	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	396	ILE	6.0
2	H	123	THR	4.6
1	G	410	CYS	3.9
2	H	142	GLY	3.9
2	H	141	SER	3.9
1	G	44	VAL	3.6
1	G	205	CYS	3.6
1	G	103	GLN	3.5
3	L	201	SER	3.4
1	G	406	THR	3.3
2	H	143	GLY	3.3
1	G	300	SER	3.3
2	H	223	LYS	3.3
1	G	119	CYS	3.0
3	L	2	ILE	3.0
3	L	153	LEU	3.0
1	G	460	ALA	2.9
1	G	228	CYS	2.9
1	G	409	GLY	2.8
1	G	125	GLY	2.6
2	H	64	PHE	2.6
2	H	219	ARG	2.6
3	L	1	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	40	ARG	2.5
1	G	462	ASN	2.5
2	H	212	SER	2.5
2	H	31	ASP	2.5
3	L	67	SER	2.4
1	G	412	GLY	2.4
1	G	463	THR	2.4
1	G	362	GLN	2.4
1	G	124	GLY	2.4
1	G	71	THR	2.3
1	G	239	CYS	2.3
3	L	68	GLY	2.3
1	G	445	CYS	2.2
3	L	122	GLU	2.2
2	H	85	SER	2.2
1	G	407	MET	2.2
2	H	213	ASN	2.1
1	G	236	THR	2.1
1	G	247	CYS	2.1
1	G	82	GLN	2.1
1	G	354	ASN	2.1
2	H	3	ARG	2.1
1	G	395	CYS	2.1
2	H	144	THR	2.1
2	H	111	HIS	2.0
1	G	45	TRP	2.0
1	G	325	ASP	2.0
2	H	195	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	G	509	14/15	0.18	0.18	111,119,121,122	0
4	NAG	G	501	14/15	0.41	0.16	115,116,117,118	0
4	NAG	G	507	14/15	0.58	0.19	61,67,68,70	0
4	NAG	G	508	14/15	0.62	0.21	85,94,99,99	0
4	NAG	G	505	14/15	0.63	0.20	113,127,130,131	0
4	NAG	G	512	14/15	0.68	0.17	95,101,105,107	0
4	NAG	G	503	14/15	0.70	0.16	78,81,95,96	0
4	NAG	G	502	14/15	0.70	0.26	97,102,105,105	0
4	NAG	G	511	14/15	0.81	0.11	86,90,96,98	0
4	NAG	G	510	14/15	0.82	0.17	50,54,59,59	0
4	NAG	G	506	14/15	0.83	0.12	46,51,62,62	0
5	EPE	G	514	15/15	0.87	0.22	100,114,150,151	15
5	EPE	G	513	15/15	0.88	0.21	109,114,124,135	0
4	NAG	G	504	14/15	0.88	0.11	39,46,51,53	0
6	15P	G	515	23/104	0.88	0.12	43,45,50,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.