



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 04:54 am GMT

PDB ID : 5G1M  
Title : Crystal structure of NagZ from Pseudomonas aeruginosa  
Authors : Acebron, I.; Artola-Recolons, C.; Mahasenan, K.; Mobashery, S.; Hermoso, J.A.  
Deposited on : 2016-03-28  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 9% 80% 14% • 5%
1	B	352	 8% 81% 13% • 5%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-HEXOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	333	2555	1599	479	463	14	0	2	0
1	B	333	2555	1599	479	463	14	0	2	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q9HZK0
A	-19	GLY	-	expression tag	UNP Q9HZK0
A	-18	SER	-	expression tag	UNP Q9HZK0
A	-17	SER	-	expression tag	UNP Q9HZK0
A	-16	HIS	-	expression tag	UNP Q9HZK0
A	-15	HIS	-	expression tag	UNP Q9HZK0
A	-14	HIS	-	expression tag	UNP Q9HZK0
A	-13	HIS	-	expression tag	UNP Q9HZK0
A	-12	HIS	-	expression tag	UNP Q9HZK0
A	-11	HIS	-	expression tag	UNP Q9HZK0
A	-10	SER	-	expression tag	UNP Q9HZK0
A	-9	SER	-	expression tag	UNP Q9HZK0
A	-8	GLY	-	expression tag	UNP Q9HZK0
A	-7	LEU	-	expression tag	UNP Q9HZK0
A	-6	VAL	-	expression tag	UNP Q9HZK0
A	-5	PRO	-	expression tag	UNP Q9HZK0
A	-4	ARG	-	expression tag	UNP Q9HZK0
A	-3	GLY	-	expression tag	UNP Q9HZK0
A	-2	SER	-	expression tag	UNP Q9HZK0
A	-1	HIS	-	expression tag	UNP Q9HZK0
B	-20	MET	-	expression tag	UNP Q9HZK0
B	-19	GLY	-	expression tag	UNP Q9HZK0
B	-18	SER	-	expression tag	UNP Q9HZK0
B	-17	SER	-	expression tag	UNP Q9HZK0
B	-16	HIS	-	expression tag	UNP Q9HZK0

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP Q9HZK0
B	-14	HIS	-	expression tag	UNP Q9HZK0
B	-13	HIS	-	expression tag	UNP Q9HZK0
B	-12	HIS	-	expression tag	UNP Q9HZK0
B	-11	HIS	-	expression tag	UNP Q9HZK0
B	-10	SER	-	expression tag	UNP Q9HZK0
B	-9	SER	-	expression tag	UNP Q9HZK0
B	-8	GLY	-	expression tag	UNP Q9HZK0
B	-7	LEU	-	expression tag	UNP Q9HZK0
B	-6	VAL	-	expression tag	UNP Q9HZK0
B	-5	PRO	-	expression tag	UNP Q9HZK0
B	-4	ARG	-	expression tag	UNP Q9HZK0
B	-3	GLY	-	expression tag	UNP Q9HZK0
B	-2	SER	-	expression tag	UNP Q9HZK0
B	-1	HIS	-	expression tag	UNP Q9HZK0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	306	Total O 306 306	0	0
5	B	264	Total O 264 264	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BETA-HEXOSAMINIDASE



- Molecule 1: BETA-HEXOSAMINIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.97Å 75.29Å 76.39Å 90.00° 110.10° 90.00°	Depositor
Resolution (Å)	46.96 – 1.80 46.96 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.96-1.80) 99.8 (46.96-1.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.79Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.196 , 0.241 0.199 , 0.243	Depositor DCC
$R_{free}$ test set	3181 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PEG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.49	4/2611 (0.2%)	0.65	2/3538 (0.1%)
1	B	0.42	0/2611	0.61	1/3538 (0.0%)
All	All	0.46	4/5222 (0.1%)	0.63	3/7076 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67	ARG	CD-NE	-7.37	1.33	1.46
1	A	67	ARG	NE-CZ	-7.11	1.23	1.33
1	A	67	ARG	CZ-NH2	-6.44	1.24	1.33
1	A	67	ARG	CZ-NH1	-6.15	1.25	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	LEU	CA-CB-CG	10.06	138.44	115.30
1	B	151	MET	CG-SD-CE	6.57	110.71	100.20
1	A	67	ARG	CA-CB-CG	6.06	126.72	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2555	0	2554	47	0
1	B	2555	0	2554	41	0
2	A	1	0	0	0	0
3	A	4	0	3	0	0
3	B	4	0	3	1	0
4	A	7	0	10	0	0
4	B	7	0	10	0	0
5	A	306	0	0	13	0
5	B	264	0	0	6	0
All	All	5703	0	5134	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:MET:SD	5:A:2243:HOH:O	2.29	0.90
1:B:114:PHE:HD1	1:B:159[A]:THR:HG21	1.42	0.83
1:B:177:ILE:HD11	1:B:211:ILE:HG13	1.60	0.83
1:B:309:ASN:HD21	1:B:312:TYR:HB3	1.45	0.81
1:A:159[B]:THR:HG22	1:A:204:ALA:HB3	1.63	0.79
1:A:70:ARG:HB2	1:B:70:ARG:HD2	1.66	0.78
1:A:133:PHE:HB3	5:A:2167:HOH:O	1.87	0.73
1:A:67:ARG:O	1:A:70:ARG:HG2	1.88	0.73
1:A:309:ASN:HD21	1:A:312:TYR:HB3	1.54	0.72
1:A:317:ARG:NH1	5:A:2305:HOH:O	2.26	0.68
1:B:249:ALA:HB1	1:B:260:ARG:HH22	1.59	0.67
1:A:209:HIS:HE1	1:A:251:ALA:HB3	1.61	0.64
1:A:63:GLN:CD	1:A:69:GLN:HE22	2.01	0.63
1:B:127:VAL:HG13	1:B:130:SER:HB3	1.78	0.63
1:B:258:ALA:O	1:B:262:GLU:HG3	1.97	0.63
1:A:31:ILE:HD11	1:A:62:ASP:HB2	1.81	0.63
1:A:128:VAL:HG23	5:A:2168:HOH:O	1.98	0.61
1:B:115:ALA:HB3	1:B:151:MET:HE3	1.81	0.61
1:A:63:GLN:OE1	1:A:69:GLN:NE2	2.34	0.61
1:B:131:ARG:N	5:B:2168:HOH:O	2.34	0.60
1:A:134:GLU:HB2	5:A:2170:HOH:O	2.01	0.60
1:A:106:GLN:HE22	1:A:306:GLY:H	1.50	0.59
1:A:64:GLU:HB2	1:A:67:ARG:HB3	1.84	0.58
1:A:122:HIS:O	1:A:124:ARG:NH1	2.37	0.58
1:B:1:MET:N	5:B:2001:HOH:O	2.37	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:ILE:HD11	1:B:62:ASP:HB2	1.85	0.57
1:B:304:GLY:HA3	3:B:1333:ACT:H2	1.87	0.56
1:A:71:LEU:HB3	1:B:70:ARG:NH1	2.20	0.56
1:B:127:VAL:CG1	1:B:130:SER:HB3	2.36	0.55
1:B:159[A]:THR:HG22	5:B:2154:HOH:O	2.07	0.54
1:A:71:LEU:HB3	1:B:70:ARG:CZ	2.37	0.54
1:A:106:GLN:NE2	1:A:306:GLY:H	2.06	0.53
1:B:207:PRO:HD3	1:B:242:PHE:O	2.08	0.53
1:B:114:PHE:CD1	1:B:159[A]:THR:HG21	2.33	0.53
1:A:278:ARG:O	1:A:282:GLU:HG3	2.09	0.52
1:B:114:PHE:HA	1:B:159[A]:THR:CG2	2.40	0.51
1:A:128:VAL:N	5:A:2168:HOH:O	2.40	0.51
1:B:249:ALA:HA	1:B:252:HIS:HB2	1.93	0.51
1:A:63:GLN:O	1:A:67:ARG:HB3	2.11	0.51
1:B:249:ALA:HB2	1:B:260:ARG:HH12	1.76	0.51
1:B:54:ARG:HB3	1:B:57[A]:LEU:HG	1.95	0.49
1:B:289:GLN:O	1:B:292:LYS:HE3	2.13	0.49
1:A:246:LEU:HB2	1:A:274:VAL:HG22	1.94	0.49
1:B:222:PHE:CE1	1:B:252:HIS:HE1	2.31	0.49
1:A:114:PHE:HA	1:A:159[B]:THR:OG1	2.12	0.48
1:A:64:GLU:HB2	1:A:67:ARG:CB	2.44	0.48
1:B:145:GLY:O	1:B:149:ARG:HG3	2.14	0.48
1:A:130:SER:N	5:A:2167:HOH:O	2.46	0.47
1:A:-1:HIS:HA	5:A:2002:HOH:O	2.13	0.47
1:A:71:LEU:HB3	1:B:70:ARG:NH2	2.29	0.47
1:A:207:PRO:HD3	1:A:242:PHE:O	2.15	0.47
1:A:316:PRO:O	1:A:320:GLU:HG3	2.15	0.46
1:B:22:LEU:HD21	1:B:30:LEU:HD12	1.96	0.46
1:A:286:ALA:O	1:A:290:ARG:HG3	2.16	0.46
1:A:67:ARG:NH1	1:A:67:ARG:HG3	2.30	0.46
1:B:53:ILE:HG22	1:B:54:ARG:HG2	1.98	0.45
1:A:64:GLU:O	1:A:67:ARG:HB2	2.16	0.45
1:B:31:ILE:HD12	1:B:114:PHE:HE2	1.81	0.45
1:A:92:GLU:HB2	5:A:2061:HOH:O	2.15	0.45
1:A:54:ARG:HB3	1:A:57[B]:LEU:HG	2.00	0.44
1:A:326:ARG:NE	1:A:332:ASP:OD2	2.47	0.44
1:B:313:ARG:HG3	5:B:2144:HOH:O	2.18	0.44
1:B:54:ARG:CB	1:B:57[A]:LEU:HG	2.48	0.44
1:B:130:SER:O	1:B:134:GLU:HG2	2.18	0.43
1:B:177:ILE:HG22	1:B:251:ALA:C	2.39	0.43
1:B:317:ARG:CZ	5:B:2270:HOH:O	2.66	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:HIS:ND1	1:A:174:HIS:O	2.52	0.43
1:A:185:GLU:H	1:A:185:GLU:CD	2.23	0.42
1:A:253:VAL:CG2	1:A:260:ARG:HD3	2.49	0.42
1:B:225:ARG:HD2	1:B:229:GLU:OE1	2.19	0.42
1:A:131:ARG:HA	5:A:2170:HOH:O	2.19	0.42
1:B:120:LEU:HD11	1:B:137:PRO:HB3	2.01	0.42
1:A:127:VAL:HA	5:A:2168:HOH:O	2.20	0.41
1:A:329:GLN:HA	5:A:2315:HOH:O	2.19	0.41
1:B:222:PHE:CZ	1:B:252:HIS:HE1	2.38	0.41
1:B:309:ASN:HD21	1:B:312:TYR:CB	2.25	0.41
1:B:177:ILE:HD13	1:B:219:PRO:HG3	2.02	0.41
1:A:209:HIS:CE1	1:A:251:ALA:HB3	2.48	0.41
1:B:139:ARG:NH2	5:B:2120:HOH:O	2.36	0.41
1:A:53:ILE:HG22	1:A:54:ARG:HG2	2.03	0.41
1:B:180:ASP:HB3	1:B:212:TYR:CD2	2.56	0.41
1:A:228:GLN:NE2	5:A:2232:HOH:O	2.41	0.40
1:A:63:GLN:O	1:A:67:ARG:N	2.54	0.40
1:A:331:ILE:O	1:A:332:ASP:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/352 (95%)	323 (97%)	10 (3%)	0	100	100
1	B	333/352 (95%)	324 (97%)	9 (3%)	0	100	100
All	All	666/704 (95%)	647 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/269 (95%)	252 (99%)	3 (1%)	71	65
1	B	255/269 (95%)	250 (98%)	5 (2%)	55	44
All	All	510/538 (95%)	502 (98%)	8 (2%)	62	54

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ARG
1	A	70	ARG
1	A	166	HIS
1	B	77	ARG
1	B	121	ASP
1	B	124	ARG
1	B	166	HIS
1	B	252	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	106	GLN
1	A	209	HIS
1	A	309	ASN
1	A	314	GLN
1	B	214	GLN
1	B	289	GLN
1	B	309	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	A	1334	-	3,3,3	0.70	0	3,3,3	1.48	0
4	PEG	A	1335	-	6,6,6	0.56	0	5,5,5	0.34	0
3	ACT	B	1333	-	3,3,3	0.86	0	3,3,3	1.38	0
4	PEG	B	1335	-	6,6,6	0.57	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	1335	-	-	1/4/4/4	-
4	PEG	A	1335	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1335	PEG	O2-C3-C4-O4
4	B	1335	PEG	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1333	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	333/352 (94%)	0.26	30 (9%) <b>9</b> <b>7</b>	11, 23, 78, 144	0
1	B	333/352 (94%)	0.32	27 (8%) <b>12</b> <b>9</b>	10, 25, 74, 159	0
All	All	666/704 (94%)	0.29	57 (8%) <b>10</b> <b>8</b>	10, 24, 76, 159	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	250	GLY	17.8
1	A	250	GLY	13.1
1	A	251	ALA	10.2
1	A	175	VAL	9.8
1	B	175	VAL	8.4
1	A	253	VAL	7.6
1	A	171	ALA	7.2
1	A	248	MET	7.1
1	B	70	ARG	7.1
1	A	125	SER	6.7
1	B	66	GLY	6.5
1	B	172	ASP	6.3
1	B	173	SER	6.2
1	A	174	HIS	6.0
1	A	249	ALA	6.0
1	B	251	ALA	5.6
1	B	125	SER	5.3
1	A	332	ASP	5.1
1	B	127	VAL	4.8
1	B	68	VAL	4.6
1	A	67	ARG	4.4
1	A	65	GLY	4.4
1	B	249	ALA	4.3
1	A	71	LEU	4.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	176	ALA	4.2
1	B	124	ARG	4.1
1	A	126	ALA	4.1
1	A	127	VAL	4.1
1	A	-1	HIS	4.1
1	A	252	HIS	4.1
1	A	128	VAL	4.0
1	B	174	HIS	4.0
1	B	67	ARG	3.9
1	A	64	GLU	3.7
1	B	176	ALA	3.7
1	A	70	ARG	3.7
1	B	69	GLN	3.6
1	B	169	ALA	3.4
1	A	131	ARG	3.4
1	A	172	ASP	3.3
1	A	173	SER	3.1
1	B	170	GLU	3.1
1	B	-1	HIS	3.0
1	B	177	ILE	3.0
1	A	254	VAL	2.8
1	A	129	GLY	2.8
1	B	128	VAL	2.8
1	B	126	ALA	2.7
1	A	123	GLN	2.6
1	A	130	SER	2.6
1	B	248	MET	2.5
1	B	252	HIS	2.5
1	B	168	TRP	2.5
1	B	171	ALA	2.4
1	A	247	SER	2.3
1	A	66	GLY	2.3
1	B	123	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PEG	B	1335	7/7	0.62	0.20	49,54,59,59	0
3	ACT	A	1334	4/4	0.83	0.14	32,44,45,52	0
4	PEG	A	1335	7/7	0.85	0.12	46,52,57,57	0
2	CL	A	1333	1/1	0.91	0.15	58,58,58,58	0
3	ACT	B	1333	4/4	0.92	0.10	18,24,30,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.