



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 05:27 AM EDT

PDB ID : 5J0L  
Title : De novo design of protein homo-oligomers with modular hydrogen bond network-mediated specificity  
Authors : Sankaran, B.; Zwart, P.H.; Pereira, J.H.; Baker, D.; Boyken, S.; Chen, Z.; Groves, B.; Langan, R.A.; Oberdorfer, G.; Ford, A.; Gilmore, J.; Xu, C.; DiMaio, F.; Seelig, G.  
Deposited on : 2016-03-28  
Resolution : 2.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 72% 21% 8%
1	B	141	 % 74% 10% 15%
1	C	141	 74% 9% 16%
1	D	141	 % 70% 13% 16%
1	E	141	 % 70% 8% 21%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	141	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '7%', followed by a large green segment labeled '59%', a yellow segment labeled '16%', and a grey segment at the end labeled '24%'. A small black dot is located on the boundary between the yellow and grey segments.</p>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called designed protein 3L6HC2\_2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	130	1020	617	188	211	4	0	0	0
1	B	120	946	575	175	193	3	0	0	0
1	C	118	937	569	171	194	3	0	0	0
1	D	119	936	569	174	190	3	0	0	0
1	E	111	871	529	163	176	3	0	0	0
1	F	107	856	519	161	173	3	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	21	Total	O	0	0
			21	21		
2	B	21	Total	O	0	0
			21	21		
2	C	33	Total	O	0	0
			33	33		
2	D	24	Total	O	0	0
			24	24		
2	E	22	Total	O	0	0
			22	22		
2	F	10	Total	O	0	0
			10	10		



- Molecule 1: designed protein 3L6HC2\_2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	152.65Å 152.65Å 70.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.10 – 2.63 66.10 – 2.63	Depositor EDS
% Data completeness (in resolution range)	100.0 (66.10-2.63) 100.0 (66.10-2.63)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.29 (at 2.62Å)	Xtrriage
Refinement program	PHENIX 1.7.2_869	Depositor
R, $R_{free}$	0.232 , 0.288 0.224 , 0.282	Depositor DCC
$R_{free}$ test set	2003 reflections (7.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.5	Xtrriage
Anisotropy	0.149	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5697	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/1020	0.43	0/1356
1	B	0.28	0/945	0.44	0/1255
1	C	0.27	0/935	0.40	0/1241
1	D	0.32	0/935	0.41	0/1241
1	E	0.29	0/870	0.41	0/1154
1	F	0.29	0/854	0.47	0/1132
All	All	0.29	0/5559	0.43	0/7379

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1020	0	1042	17	1
1	B	946	0	974	10	0
1	C	937	0	956	10	1
1	D	936	0	969	11	0
1	E	871	0	901	7	0
1	F	856	0	880	14	0
2	A	21	0	0	1	0
2	B	21	0	0	1	0
2	C	33	0	0	3	0
2	D	24	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	22	0	0	3	0
2	F	10	0	0	1	0
All	All	5697	0	5722	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:58:GLU:OE2	1:E:122:ARG:NH2	2.16	0.79
1:F:100:GLU:HG3	1:F:104:ARG:HH11	1.51	0.76
1:D:3:MET:HB2	1:D:85:VAL:HG11	1.73	0.69
1:A:2:LEU:HD12	1:A:89:ALA:HB1	1.74	0.68
1:F:48:LEU:HD13	1:F:130:GLU:HB2	1.77	0.66
1:E:35:LEU:HD23	1:E:52:MET:HG2	1.82	0.62
1:F:58:GLU:OE2	1:F:122:ARG:NH2	2.33	0.61
1:D:118:LYS:NZ	2:D:202:HOH:O	2.33	0.61
1:F:99:GLU:N	1:F:99:GLU:OE1	2.33	0.60
1:F:45:SER:N	2:F:202:HOH:O	2.34	0.59
1:C:91:GLU:N	2:C:204:HOH:O	2.34	0.59
1:F:125:TYR:HA	1:F:128:ASP:HB2	1.86	0.58
1:F:80:ARG:HH22	1:F:104:ARG:HH12	1.52	0.57
1:A:7:ALA:HB1	1:B:33:GLU:HG3	1.86	0.56
1:C:100:GLU:HG3	1:C:104:ARG:HE	1.72	0.54
1:A:110:GLU:OE1	1:A:113:ARG:NH2	2.33	0.54
1:A:5:LYS:NZ	1:A:95:ASP:OD1	2.27	0.54
1:F:21:ALA:HB2	1:F:67:ILE:HD13	1.91	0.52
1:E:51:LEU:N	2:E:201:HOH:O	2.42	0.52
1:E:117:LYS:NZ	2:E:202:HOH:O	2.42	0.52
1:D:23:GLU:OE2	1:E:15:ARG:NH2	2.33	0.52
1:B:89:ALA:HB1	1:B:93:GLU:HG2	1.91	0.52
1:C:124:LYS:NZ	2:C:207:HOH:O	2.43	0.51
1:F:38:GLU:OE1	1:F:41:ARG:NE	2.38	0.51
1:A:81:LEU:O	1:A:85:VAL:HG23	2.12	0.50
1:A:57:ARG:NH2	1:C:58:GLU:OE2	2.45	0.50
1:A:42:SER:O	1:B:86:LYS:NZ	2.41	0.48
1:A:42:SER:OG	1:A:43:SER:N	2.47	0.48
1:D:58:GLU:OE2	1:D:122:ARG:NH2	2.45	0.48
1:E:80:ARG:NH2	2:E:204:HOH:O	2.46	0.47
1:C:110:GLU:OE1	1:C:113:ARG:NH2	2.47	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:45:SER:OG	1:D:46:GLU:N	2.48	0.47
1:F:100:GLU:HG3	1:F:104:ARG:NH1	2.25	0.47
1:A:2:LEU:N	2:A:204:HOH:O	2.49	0.46
1:A:11:LYS:HG3	1:B:29:ALA:HB1	1.99	0.45
1:D:5:LYS:HE3	1:D:91:GLU:HG3	1.97	0.45
1:D:73:LEU:HD22	1:D:105:LEU:HG	1.99	0.45
1:D:35:LEU:HD12	1:D:124:LYS:HD3	2.00	0.44
1:B:115:GLU:O	1:B:119:ILE:HG13	2.18	0.44
1:D:30:LYS:NZ	1:D:34:ASP:OD1	2.50	0.44
1:C:100:GLU:HG3	1:C:104:ARG:NE	2.31	0.43
1:B:35:LEU:HB3	1:B:52:MET:HG3	2.00	0.43
1:A:30:LYS:NZ	1:A:34:ASP:OD1	2.52	0.43
1:A:92:LYS:O	1:A:96:GLU:HB2	2.19	0.42
1:A:53:ALA:HB1	1:A:57:ARG:NH1	2.35	0.42
1:D:47:ARG:HA	1:D:47:ARG:HD2	1.43	0.42
1:F:39:ILE:HD13	1:F:48:LEU:HG	2.02	0.42
1:F:9:ALA:O	1:F:13:LEU:HG	2.21	0.41
1:A:33:GLU:HG3	1:B:7:ALA:HB1	2.01	0.41
1:A:80:ARG:HA	1:A:83:GLU:OE1	2.20	0.41
1:B:73:LEU:HD12	1:B:105:LEU:HD23	2.03	0.41
1:C:105:LEU:HD23	1:C:105:LEU:HA	1.92	0.41
1:D:108:LEU:HD23	1:D:108:LEU:HA	1.91	0.41
1:F:73:LEU:HD21	1:F:104:ARG:HE	1.86	0.41
1:B:80:ARG:NH1	2:B:207:HOH:O	2.53	0.41
1:C:100:GLU:O	1:C:104:ARG:HG2	2.20	0.41
1:E:79:ILE:HG22	1:E:83:GLU:OE1	2.20	0.40
1:A:39:ILE:HD12	1:A:49:LEU:HD12	2.04	0.40
1:A:105:LEU:HD23	1:A:105:LEU:HA	1.92	0.40
1:B:47:ARG:HB2	1:B:48:LEU:H	1.56	0.40
1:C:24:LEU:HD11	1:C:112:SER:OG	2.22	0.40
1:F:46:GLU:O	1:F:50:GLU:HG2	2.21	0.40
1:C:46:GLU:N	2:C:210:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:TYR:OH	1:C:83:GLU:OE2[5_555]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/141 (91%)	122 (95%)	4 (3%)	2 (2%)	9	13
1	B	116/141 (82%)	113 (97%)	3 (3%)	0	100	100
1	C	112/141 (79%)	112 (100%)	0	0	100	100
1	D	115/141 (82%)	114 (99%)	1 (1%)	0	100	100
1	E	107/141 (76%)	105 (98%)	1 (1%)	1 (1%)	17	26
1	F	101/141 (72%)	97 (96%)	2 (2%)	2 (2%)	7	10
All	All	679/846 (80%)	663 (98%)	11 (2%)	5 (1%)	22	32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	GLU
1	F	126	GLU
1	F	125	TYR
1	A	88	GLY
1	E	52	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/115 (90%)	102 (98%)	2 (2%)	57	74
1	B	95/115 (83%)	92 (97%)	3 (3%)	39	56
1	C	95/115 (83%)	93 (98%)	2 (2%)	53	71

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	95/115 (83%)	91 (96%)	4 (4%)	30	45
1	E	87/115 (76%)	85 (98%)	2 (2%)	50	68
1	F	87/115 (76%)	84 (97%)	3 (3%)	37	53
All	All	563/690 (82%)	547 (97%)	16 (3%)	43	61

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	SER
1	A	99	GLU
1	B	47	ARG
1	B	49	LEU
1	B	86	LYS
1	C	60	GLN
1	C	91	GLU
1	D	47	ARG
1	D	117	LYS
1	D	124	LYS
1	D	125	TYR
1	E	91	GLU
1	E	105	LEU
1	F	8	ARG
1	F	73	LEU
1	F	111	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	130/141 (92%)	0.03	0 <b>100</b> <b>100</b>	24, 49, 83, 102	0
1	B	120/141 (85%)	-0.07	1 (0%) <b>86</b> <b>85</b>	25, 43, 82, 110	0
1	C	118/141 (83%)	-0.09	0 <b>100</b> <b>100</b>	21, 42, 75, 80	0
1	D	119/141 (84%)	-0.01	1 (0%) <b>86</b> <b>85</b>	22, 50, 83, 92	0
1	E	111/141 (78%)	-0.02	1 (0%) <b>84</b> <b>83</b>	28, 53, 81, 96	0
1	F	107/141 (75%)	0.47	10 (9%) <b>8</b> <b>6</b>	32, 61, 100, 120	0
All	All	705/846 (83%)	0.05	13 (1%) <b>68</b> <b>66</b>	21, 49, 83, 120	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	97	ILE	5.5
1	F	10	ALA	4.1
1	F	13	LEU	3.2
1	F	99	GLU	3.0
1	F	80	ARG	2.8
1	F	79	ILE	2.8
1	F	101	ALA	2.6
1	F	98	VAL	2.6
1	F	103	LYS	2.5
1	F	100	GLU	2.4
1	B	85	VAL	2.3
1	E	36	MET	2.2
1	D	48	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.