



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 07:53 PM EDT

PDB ID : 5KYM
Title : Crystal Structure of the 1-acyl-sn-glycerophosphate (LPA) acyltransferase, PlsC, from *Thermotoga maritima*
Authors : Robertson, R.M.; Yao, J.; Gajewski, S.; Kumar, G.; Martin, E.W.; Rock, C.O.; White, S.W.
Deposited on : 2016-07-21
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	 84% 11% ...
1	B	247	 12% 78% 15% • 6%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-acyl-sn-glycerol-3-phosphate acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	Total	C	N	O	S	0	0	0
			1973	1306	336	327	4			
1	B	232	Total	C	N	O	S	0	0	0
			1886	1246	319	317	4			

- Molecule 2 is 1-HEPTADECANOYL-2-TRIDECANOYL-3-GLYCEROL-PHOSPHONYL CHOLINE (three-letter code: HGX) (formula: $C_{38}H_{77}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	O	P	N		
2	A	1	Total	C	O	P	0	0	
			32	23	8	1			
2	A	1	Total	C	O	P	0	0	
			32	23	8	1			
2	A	1	Total	C	O	P	0	0	
			32	23	8	1			
2	B	1	Total	C	N	O	P	0	0
			44	34	1	8	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			43	33	1	8	1		
2	B	1	Total	C	O	P		0	0
			42	33	8	1			

- Molecule 3 is nonane (three-letter code: DD9) (formula: C₉H₂₀).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			9	9		
3	B	1	Total	C	0	0
			9	9		

- Molecule 4 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: C₂₄H₄₆O₁₁).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	C O	0	0
			35	24 11		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 1-acyl-sn-glycerol-3-phosphate acyltransferase

Chain A:  84% 11% ...



- Molecule 1: 1-acyl-sn-glycerol-3-phosphate acyltransferase

Chain B:  12% 78% 15% • 6%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	173.84Å 173.84Å 173.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.46 – 2.80 46.46 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.46-2.80) 100.0 (46.46-2.80)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.219 , 0.277 0.223 , 0.275	Depositor DCC
R_{free} test set	1678 reflections (7.76%)	wwPDB-VP
Wilson B-factor (Å ²)	65.4	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.028 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4138	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LMT, HGX, DD9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.94	1/2023 (0.0%)	1.04	6/2728 (0.2%)
1	B	0.75	0/1932	0.94	0/2608
All	All	0.85	1/3955 (0.0%)	0.99	6/5336 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	143	GLU	CD-OE2	5.44	1.31	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	A	37	ASP	CB-CG-OD2	-6.09	112.82	118.30
1	A	242	LYS	CD-CE-NZ	5.86	125.17	111.70
1	A	42	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	38	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	173	LEU	CA-CB-CG	-5.39	102.90	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	4	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	2053	17	0
1	B	1886	0	1947	28	0
2	A	96	0	105	0	0
2	B	129	0	183	1	0
3	A	9	0	20	0	0
3	B	9	0	20	1	0
4	B	35	0	46	0	0
5	A	1	0	0	0	0
All	All	4138	0	4374	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ILE:HB	1:B:91:PRO:CD	2.26	0.66
1:B:241:LEU:O	1:B:243:ALA:N	2.32	0.62
1:A:189:TRP:CH2	1:A:242:LYS:HE2	2.34	0.62
1:B:79:ILE:HD12	1:B:181:VAL:HG22	1.82	0.62
1:B:90:ILE:HG21	3:B:305:DD9:H9B	1.83	0.60
1:A:42:ARG:HH11	1:A:42:ARG:HG3	1.67	0.59
1:A:173:LEU:HD13	1:A:227:ARG:HA	1.86	0.58
1:B:90:ILE:HB	1:B:91:PRO:HD3	1.89	0.55
1:B:93:ILE:HD11	1:B:188:ILE:HD11	1.89	0.55
1:B:108:LEU:HD23	1:B:111:ILE:HD13	1.88	0.55
1:A:93:ILE:HD11	1:A:188:ILE:HD11	1.89	0.55
1:A:179:THR:OG1	1:A:181:VAL:HG13	2.07	0.54
1:B:217:PRO:HB3	1:B:226:LEU:HD22	1.90	0.54
1:A:135:VAL:HG12	1:A:135:VAL:O	2.09	0.52
1:A:104:ALA:HB1	1:A:108:LEU:HD12	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:TYR:CE1	1:B:16:ILE:HD11	2.45	0.52
1:A:139:ARG:O	1:A:143:GLU:HG2	2.10	0.51
1:A:76:ARG:HH11	1:A:76:ARG:HG3	1.77	0.50
1:B:79:ILE:CD1	1:B:181:VAL:HG22	2.42	0.50
1:A:76:ARG:HH11	1:A:76:ARG:CG	2.25	0.49
1:A:12:TYR:CE1	1:A:16:ILE:HD11	2.48	0.49
1:A:103:ILE:HA	1:A:124:VAL:O	2.13	0.48
1:B:158:THR:OG1	1:B:197:LYS:NZ	2.46	0.48
1:B:176:ALA:HB1	1:B:217:PRO:HG2	1.96	0.47
1:B:189:TRP:CE3	1:B:238:VAL:HG13	2.49	0.47
2:B:301:HGX:H233	2:B:301:HGX:C44	2.45	0.47
1:A:42:ARG:HG3	1:A:42:ARG:NH1	2.31	0.46
1:A:76:ARG:HG3	1:A:76:ARG:NH1	2.31	0.46
1:B:208:PHE:HB3	1:B:241:LEU:HD22	1.98	0.46
1:B:90:ILE:HD12	1:B:102:PHE:CE1	2.51	0.44
1:B:35:MET:CE	1:B:40:LYS:HG3	2.47	0.44
1:B:65:VAL:CG1	1:B:208:PHE:CD1	3.00	0.44
1:B:102:PHE:CE2	1:B:152:ILE:HD12	2.53	0.44
1:B:145:LEU:CD1	1:B:175:ILE:CG2	2.95	0.43
1:A:138:LEU:O	1:A:142:ILE:HG13	2.19	0.43
1:B:108:LEU:CD2	1:B:111:ILE:HD13	2.47	0.43
1:A:199:ARG:NH2	1:A:203:THR:HG23	2.34	0.43
1:B:176:ALA:HB1	1:B:217:PRO:CG	2.49	0.43
1:B:4:ILE:HD12	1:B:4:ILE:C	2.40	0.42
1:B:111:ILE:HD12	1:B:111:ILE:H	1.84	0.42
1:B:179:THR:OG1	1:B:181:VAL:HG13	2.20	0.42
1:A:81:VAL:HG13	1:A:155:PRO:CG	2.50	0.41
1:B:4:ILE:HD12	1:B:4:ILE:O	2.20	0.41
1:B:164:LYS:HE3	1:B:166:LEU:HD21	2.04	0.40
1:B:228:LYS:O	1:B:232:GLU:HG2	2.21	0.40
1:B:155:PRO:HG2	1:B:168:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/247 (96%)	227 (96%)	10 (4%)	0	100	100
1	B	226/247 (92%)	213 (94%)	12 (5%)	1 (0%)	34	66
All	All	463/494 (94%)	440 (95%)	22 (5%)	1 (0%)	47	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	242	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/218 (96%)	201 (96%)	9 (4%)	29	62
1	B	201/218 (92%)	196 (98%)	5 (2%)	47	80
All	All	411/436 (94%)	397 (97%)	14 (3%)	37	71

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	3	LYS
1	A	76	ARG
1	A	138	LEU
1	A	156	GLU
1	A	173	LEU
1	A	221	SER
1	A	224	GLU
1	A	232	GLU
1	B	107	GLU
1	B	135	VAL
1	B	181	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	184	LEU
1	B	244	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HGX	A	302	-	31,31,47	1.38	2 (6%)	34,36,55	1.42	3 (8%)
2	HGX	B	302	-	42,42,47	1.14	2 (4%)	48,50,55	1.03	3 (6%)
3	DD9	A	304	-	8,8,8	0.32	0	7,7,7	0.74	0
2	HGX	B	301	-	43,43,47	1.10	2 (4%)	49,51,55	0.99	3 (6%)
4	LMT	B	304	-	36,36,36	0.66	1 (2%)	47,47,47	1.16	4 (8%)
2	HGX	B	303	-	41,41,47	1.18	2 (4%)	44,46,55	1.12	4 (9%)
2	HGX	A	303	-	31,31,47	1.48	2 (6%)	34,36,55	1.42	4 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DD9	B	305	-	8,8,8	0.42	0	7,7,7	0.58	0
2	HGX	A	301	-	31,31,47	1.30	2 (6%)	34,36,55	1.26	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HGX	A	302	-	-	17/35/35/51	-
2	HGX	B	302	-	-	14/46/46/51	-
3	DD9	A	304	-	-	0/6/6/6	-
2	HGX	B	301	-	-	16/47/47/51	-
4	LMT	B	304	-	-	12/21/61/61	0/2/2/2
2	HGX	B	303	-	-	11/45/45/51	-
2	HGX	A	303	-	-	9/35/35/51	-
3	DD9	B	305	-	-	1/6/6/6	-
2	HGX	A	301	-	-	12/35/35/51	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	303	HGX	O2-C31	5.45	1.49	1.34
2	A	303	HGX	O3-C11	5.24	1.48	1.33
2	B	303	HGX	O3-C11	4.97	1.47	1.33
2	A	302	HGX	O2-C31	4.81	1.47	1.34
2	B	302	HGX	O3-C11	4.62	1.46	1.33
2	B	302	HGX	O2-C31	4.61	1.47	1.34
2	B	301	HGX	O3-C11	4.60	1.46	1.33
2	A	301	HGX	O3-C11	4.57	1.46	1.33
2	A	301	HGX	O2-C31	4.46	1.46	1.34
2	A	302	HGX	O3-C11	4.42	1.46	1.33
2	B	303	HGX	O2-C31	4.42	1.46	1.34
2	B	301	HGX	O2-C31	4.14	1.46	1.34
4	B	304	LMT	O1'-C1'	2.03	1.43	1.40

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	HGX	O2-C31-C32	6.16	124.77	111.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	303	HGX	O2-C31-C32	5.10	122.50	111.50
2	B	303	HGX	O2-C31-C32	4.39	120.97	111.50
2	A	301	HGX	O2-C31-C32	4.01	120.15	111.50
2	B	301	HGX	O3-C11-C12	3.74	123.66	111.91
2	B	302	HGX	O2-C31-C32	3.73	119.55	111.50
2	A	301	HGX	O3-C11-C12	3.73	123.62	111.91
2	B	302	HGX	O3-C11-C12	3.71	123.56	111.91
2	A	302	HGX	O2-C31-O31	-3.42	115.42	123.70
4	B	304	LMT	O1'-C1'-C2'	3.40	113.61	108.30
2	B	302	HGX	O3-C11-O11	-3.02	115.96	123.59
2	B	301	HGX	O3-C11-O11	-2.98	116.07	123.59
2	B	301	HGX	O2-C31-C32	2.98	117.92	111.50
4	B	304	LMT	O1B-C1B-C2B	2.91	115.64	108.10
2	B	303	HGX	O3-C11-C12	2.87	120.90	111.91
4	B	304	LMT	O5'-C1'-C2'	-2.77	104.49	110.35
2	B	303	HGX	C3-O3-C11	2.58	126.67	117.12
2	A	303	HGX	O3-C11-C12	2.52	119.81	111.91
2	A	303	HGX	C1-C2-C3	-2.40	106.12	111.79
2	B	303	HGX	O3-C11-O11	-2.32	117.75	123.59
2	A	303	HGX	C3-O3-C11	2.28	125.55	117.12
2	A	301	HGX	O3-C11-O11	-2.26	117.90	123.59
4	B	304	LMT	O5'-C5'-C4'	2.23	114.46	109.75
2	A	302	HGX	C3-O3-C11	2.01	124.56	117.12

There are no chirality outliers.

All (92) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	HGX	C4-O5-P1-O4
2	A	302	HGX	O2-C2-C3-O3
2	A	302	HGX	C1-O3P-P1-O4
2	A	302	HGX	C1-O3P-P1-O5
2	A	302	HGX	C32-C31-O2-C2
2	A	302	HGX	C4-O5-P1-O3P
2	A	302	HGX	C4-O5-P1-O1
2	A	302	HGX	C4-O5-P1-O4
2	B	301	HGX	C2-C1-O3P-P1
2	B	301	HGX	C1-O3P-P1-O1
2	B	302	HGX	C1-O3P-P1-O5
2	A	302	HGX	O31-C31-O2-C2
2	B	303	HGX	C32-C31-O2-C2
4	B	304	LMT	O5'-C5'-C6'-O6'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	303	HGX	O31-C31-O2-C2
2	B	301	HGX	C12-C11-O3-C3
2	B	301	HGX	O11-C11-O3-C3
2	B	301	HGX	C1-O3P-P1-O5
2	B	303	HGX	C4-O5-P1-O3P
4	B	304	LMT	O5B-C1B-O1B-C4'
2	B	301	HGX	C15-C16-C17-C18
2	B	301	HGX	C13-C14-C15-C16
4	B	304	LMT	C2B-C1B-O1B-C4'
2	B	302	HGX	O31-C31-O2-C2
4	B	304	LMT	C4-C5-C6-C7
2	B	302	HGX	C32-C31-O2-C2
2	B	301	HGX	C39-C40-C42-C43
2	B	302	HGX	C12-C13-C14-C15
2	A	302	HGX	C12-C11-O3-C3
2	A	302	HGX	O11-C11-O3-C3
2	B	301	HGX	C37-C38-C39-C40
2	A	303	HGX	C32-C31-O2-C2
2	A	303	HGX	O31-C31-O2-C2
2	A	301	HGX	C15-C16-C17-C18
2	A	303	HGX	C32-C33-C34-C35
4	B	304	LMT	C2-C3-C4-C5
4	B	304	LMT	C4'-C5'-C6'-O6'
4	B	304	LMT	C1-C2-C3-C4
2	B	302	HGX	C36-C37-C38-C39
2	B	301	HGX	C11-C12-C13-C14
2	A	301	HGX	C12-C13-C14-C15
2	B	303	HGX	C14-C15-C16-C17
2	B	301	HGX	O2-C31-C32-C33
2	A	302	HGX	C33-C34-C35-C36
2	A	301	HGX	C32-C33-C34-C35
2	B	302	HGX	C14-C15-C16-C17
2	B	302	HGX	C35-C36-C37-C38
2	B	303	HGX	O3P-C1-C2-O2
4	B	304	LMT	C9-C10-C11-C12
2	B	301	HGX	O2-C2-C3-O3
2	A	301	HGX	O31-C31-O2-C2
4	B	304	LMT	O5B-C5B-C6B-O6B
2	B	302	HGX	C17-C18-C19-C20
2	A	302	HGX	C1-C2-C3-O3
2	A	301	HGX	O3-C11-C12-C13
2	B	303	HGX	C33-C34-C35-C36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	302	HGX	C12-C13-C14-C15
2	A	301	HGX	C4-O5-P1-O3P
2	B	301	HGX	C1-O3P-P1-O4
2	B	302	HGX	C1-O3P-P1-O1
2	B	303	HGX	C4-O5-P1-O4
2	A	302	HGX	O3P-C1-C2-C3
2	A	301	HGX	C32-C31-O2-C2
4	B	304	LMT	C4B-C5B-C6B-O6B
2	B	302	HGX	C40-C42-C43-C44
2	A	302	HGX	O3P-C1-C2-O2
2	B	302	HGX	C16-C17-C18-C19
2	B	303	HGX	C40-C42-C43-C44
2	B	303	HGX	O3P-C1-C2-C3
2	B	302	HGX	C12-C11-O3-C3
2	A	303	HGX	C1-O3P-P1-O5
2	A	303	HGX	C4-O5-P1-O3P
2	B	302	HGX	O11-C11-O3-C3
2	B	303	HGX	C5-C4-O5-P1
2	B	303	HGX	C43-C44-C45-C46
2	B	301	HGX	O31-C31-C32-C33
2	A	301	HGX	C35-C36-C37-C38
2	B	301	HGX	C40-C42-C43-C44
2	A	302	HGX	O2-C31-C32-C33
2	B	301	HGX	C1-C2-C3-O3
2	B	302	HGX	C32-C33-C34-C35
2	A	303	HGX	O2-C31-C32-C33
2	A	302	HGX	O31-C31-C32-C33
2	A	301	HGX	C36-C37-C38-C39
4	B	304	LMT	C6-C7-C8-C9
2	A	301	HGX	C1-O3P-P1-O1
2	A	301	HGX	C1-O3P-P1-O4
2	A	303	HGX	C1-O3P-P1-O4
2	A	303	HGX	C4-O5-P1-O4
2	A	303	HGX	O31-C31-C32-C33
3	B	305	DD9	C5-C6-C7-C8
4	B	304	LMT	C2-C1-O1'-C1'

There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	HGX	1	0




Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	305	DD9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/247 (97%)	-0.32	0 100 100	35, 53, 92, 131	0
1	B	232/247 (93%)	0.47	29 (12%) 3 2	59, 97, 145, 157	0
All	All	473/494 (95%)	0.07	29 (6%) 21 13	35, 71, 131, 157	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	155	PRO	3.9
1	B	173	LEU	3.6
1	B	176	ALA	3.6
1	B	217	PRO	3.6
1	B	226	LEU	3.4
1	B	168	PHE	3.4
1	B	236	ARG	3.3
1	B	189	TRP	3.0
1	B	208	PHE	2.9
1	B	220	PHE	2.8
1	B	180	GLY	2.7
1	B	179	THR	2.7
1	B	241	LEU	2.6
1	B	215	VAL	2.6
1	B	185	PRO	2.6
1	B	221	SER	2.5
1	B	187	SER	2.4
1	B	210	LYS	2.4
1	B	181	VAL	2.4
1	B	223	GLU	2.3
1	B	39	GLU	2.3
1	B	206	LYS	2.2
1	B	184	LEU	2.2
1	B	183	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	209	LEU	2.1
1	B	205	GLY	2.1
1	B	224	GLU	2.1
1	B	177	VAL	2.1
1	B	227	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	LMT	B	304	35/35	0.66	0.28	87,108,141,146	0
2	HGX	A	303	32/48	0.76	0.33	60,98,208,217	0
2	HGX	B	302	43/48	0.78	0.30	89,117,204,231	0
2	HGX	B	301	44/48	0.78	0.29	93,115,170,188	0
2	HGX	B	303	42/48	0.82	0.24	81,115,204,244	0
2	HGX	A	302	32/48	0.82	0.26	69,87,141,146	0
2	HGX	A	301	32/48	0.86	0.23	66,83,147,149	0
3	DD9	B	305	9/9	0.88	0.33	64,68,93,95	0
3	DD9	A	304	9/9	0.96	0.26	50,54,56,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HGX B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HGX B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HGX B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around HGX A 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.