

Full wwPDB NMR Structure Validation Report (i)

Dec 25, 2024 – 04:10 AM EST

PDB ID : 5LG7 BMRB ID : 25248

Title : Solution NMR structure of Tryptophan to Arginine mutant of Arkadia RING

domain

Authors: Birkou, M.; Chasapis, C.T.; Loutsidou, A.K.; Bentrop, D.; Lelli, M.; Her-

rmann, T.; Episkopou, V.; Spyroulias, G.A.

Deposited on : 2016-07-06

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &: & v1.2 \\ BMRB \ Restraints \ Analysis &: & v1.2 \\ \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 49%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$ m NMR~archive \ (\#Entries)$		
Clashscore	210492	14027		
Ramachandran outliers	207382	12486		
Sidechain outliers	206894	12463		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	69	67%	7%	26%



2 Ensemble composition and analysis (i)

This entry contains 31 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid					
1	A:12-A:62 (51)	0.86	1		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 8, 11, 16, 18, 21, 29, 30
2	4, 5, 9, 15, 22, 25, 26, 27
3	6, 7, 17, 19, 24, 31
4	13, 14, 23, 28
5	3, 10
Single-model clusters	12; 20



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1079 atoms, of which 530 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called E3 ubiquitin-protein ligase Arkadia.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	60	Total	С	Н	N	О	S	0
1	1 A	A 69	1077	331	530	94	114	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference	
A	1	MET	-	initiating methionine	UNP Q6ZNA4	
A	47	ARG	TRP	engineered mutation	UNP Q6ZNA4	

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms
2	A	2	Total Zn 2 2




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

• Molecule 1: E3 ubiquitin-protein ligase Arkadia



4.2.2 Score per residue for model 2






4.2.3 Score per residue for model 3

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.4 Score per residue for model 4

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.5 Score per residue for model 5

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.6 Score per residue for model 6

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.7 Score per residue for model 7






4.2.8 Score per residue for model 8

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.9 Score per residue for model 9

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.10 Score per residue for model 10

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.11 Score per residue for model 11

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.12 Score per residue for model 12






4.2.13 Score per residue for model 13

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.14 Score per residue for model 14

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.15 Score per residue for model 15

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.16 Score per residue for model 16

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.17 Score per residue for model 17






4.2.18 Score per residue for model 18

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.19 Score per residue for model 19

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.20 Score per residue for model 20

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.21 Score per residue for model 21

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.22 Score per residue for model 22






4.2.23 Score per residue for model 23

• Molecule 1: E3 ubiquitin-protein ligase Arkadia



4.2.24 Score per residue for model 24

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.25 Score per residue for model 25

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.26 Score per residue for model 26

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.27 Score per residue for model 27





4.2.28 Score per residue for model 28

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.29 Score per residue for model 29

• Molecule 1: E3 ubiquitin-protein ligase Arkadia




4.2.30 Score per residue for model 30

• Molecule 1: E3 ubiquitin-protein ligase Arkadia



4.2.31 Score per residue for model 31





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: molecular mechanics.

Of the 31 calculated structures, 31 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
UNIO	structure calculation	10
Amber	refinement	5.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	427
Number of shifts mapped to atoms	427
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	49%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
MIOI		RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.47 ± 0.01	$0\pm0/416~(~0.0\pm~0.0\%)$	0.93 ± 0.04	$0\pm0/561~(~0.0\pm~0.1\%)$	
All	All	0.47	0/12896 (0.0%)	0.93	3/17391 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	0.2 ± 0.4
All	All	0	5

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain Res	Chain Res Type Atoms		Atoms	7	$Observed(^o)$	$Ideal(^{o})$	Models	
IVIOI	Chain	nes	Type	Atoms Z Observed(*) Ideal(ideai()	Worst	Total	
1	A	56	ILE	CB-CG1-CD1	8.48	137.64	113.90	1	1
1	A	40	HIS	CA-CB-CG	6.74	125.06	113.60	14	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	31	ARG	Sidechain	2
1	A	35	CYS	Peptide	1
1	A	54	CYS	Peptide	1
1	A	40	HIS	Sidechain	1



6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	411	410	410	1±1
All	All	12803	12710	12709	41

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom 2	Clash (Å)	Distance (Å)	Models	
		Clash(A)	$\operatorname{Distance}(\mathrm{\AA})$	Worst	Total
1:A:19:ILE:HG23	1:A:47:ARG:HB2	0.69	1.65	14	6
1:A:38:LEU:HD23	1:A:38:LEU:O	0.59	1.98	12	1
1:A:19:ILE:HG21	1:A:43:CYS:O	0.59	1.98	14	3
1:A:54:CYS:CB	1:A:57:CYS:HB2	0.55	2.32	3	1
1:A:38:LEU:HD12	1:A:38:LEU:C	0.53	2.24	29	3
1:A:54:CYS:CB	1:A:57:CYS:SG	0.52	2.95	22	2
1:A:54:CYS:HB3	1:A:57:CYS:CB	0.51	2.35	9	2
1:A:33:LEU:HD22	1:A:61:ILE:CG2	0.49	2.37	13	4
1:A:33:LEU:HD11	1:A:39:PHE:CE1	0.46	2.45	4	1
1:A:33:LEU:HB3	1:A:61:ILE:HG23	0.46	1.88	25	1
1:A:54:CYS:HB2	1:A:57:CYS:HB2	0.44	1.89	3	1
1:A:54:CYS:HB2	1:A:61:ILE:HD11	0.44	1.89	29	1
1:A:38:LEU:HD12	1:A:39:PHE:N	0.44	2.27	24	1
1:A:53:LYS:HE2	1:A:60:ASP:CG	0.43	2.32	3	1
1:A:33:LEU:HD11	1:A:39:PHE:CD1	0.43	2.48	4	1
1:A:54:CYS:SG	1:A:55:PRO:HD2	0.43	2.53	12	1
1:A:33:LEU:HD12	1:A:61:ILE:CG2	0.43	2.44	3	1
1:A:54:CYS:HB3	1:A:57:CYS:HB3	0.42	1.90	9	1
1:A:14:GLU:O	1:A:38:LEU:HD21	0.42	2.14	12	1
1:A:24:LEU:HD22	1:A:24:LEU:N	0.42	2.30	5	1
1:A:23:ILE:N	1:A:23:ILE:CD1	0.41	2.83	1	1
1:A:12:ASP:O	1:A:16:LYS:HE2	0.41	2.15	22	2
1:A:38:LEU:HD23	1:A:38:LEU:C	0.41	2.35	6	2
1:A:54:CYS:HB3	1:A:57:CYS:HB2	0.40	1.92	16	1
1:A:23:ILE:HD12	1:A:23:ILE:N	0.40	2.32	22	1



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	51/69 (74%)	34±3 (67±6%)	14±3 (28±5%)	3±1 (5±2%)	3 24
All	All	1581/2139 (74%)	1062 (67%)	440 (28%)	79 (5%)	3 24

All 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	MET	10
1	A	56	ILE	10
1	A	13	THR	10
1	A	55	PRO	9
1	A	26	GLU	8
1	A	25	GLU	6
1	A	52	LYS	5
1	A	57	CYS	4
1	A	12	ASP	4
1	A	62	GLU	4
1	A	49	ILE	4
1	A	21	LEU	2
1	A	38	LEU	2
1	A	41	GLN	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	A	50/65~(77%)	43±1 (85±3%)	7±1 (15±3%)	5	43
All	All	1550/2015 (77%)	1323 (85%)	227 (15%)	5	43



All 35 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	38	LEU	18
1	A	54	CYS	16
1	A	32	ARG	15
1	A	52	LYS	14
1	A	57	CYS	13
1	A	40	HIS	12
1	A	41	GLN	12
1	A	22	SER	9
1	A	15	GLU	9
1	A	31	ARG	9
1	A	53	LYS	8
1	A	33	LEU	8
1	A	37	HIS	8
1	A	62	GLU	6
1	A	56	ILE	5
1	A	45	ASP	5
1	A	29	ASP	5
1	A	35	CYS	5
1	A	46	GLN	5
1	A	59	VAL	5
1	A	20	CYS	5
1	A	14	GLU	4
1	A	36	MET	4
1	A	26	GLU	4
1	A	58	ARG	3
1	A	12	ASP	3
1	A	28	GLU	3
1	A	47	ARG	3
1	A	21	LEU	3
1	A	51	ASN	2
1	A	60	ASP	2
1	A	23	ILE	1
1	A	25	GLU	1
1	A	24	LEU	1
1	A	18	THR	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.



6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 49% for the well-defined parts and 46% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	427
Number of shifts mapped to atoms	427
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	68	-0.02 ± 0.18	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	59	0.09 ± 0.13	None needed (< 0.5 ppm)
¹³ C'	0		None (insufficient data)
^{15}N	64	0.14 ± 0.39	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 49%, i.e. 349 atoms were assigned a chemical shift out of a possible 718. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	149/252 (59%)	50/101 (50%)	51/102 (50%)	48/49 (98%)
Sidechain	200/442 (45%)	116/285 (41%)	84/139 (60%)	0/18 (0%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}\mathbf{N}$
Aromatic	0/24 (0%)	0/13 (0%)	0/9 (0%)	0/2 (0%)
Overall	349/718 (49%)	166/399 (42%)	135/250 (54%)	48/69 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 46%, i.e. 427 atoms were assigned a chemical shift out of a possible 928. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	198/342 (58%)	$66/138 \ (48\%)$	68/138 (49%)	64/66 (97%)
Sidechain	229/562 (41%)	132/360 (37%)	97/181 (54%)	0/21 (0%)
Aromatic	0/24 (0%)	0/13 (0%)	0/9 (0%)	0/2 (0%)
Overall	427/928 (46%)	198/511 (39%)	$165/328 \ (50\%)$	64/89 (72%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	575
Intra-residue ($ i-j =0$)	97
Sequential ($ i-j =1$)	245
Medium range ($ i-j >1$ and $ i-j <5$)	85
Long range (i-j ≥5)	148
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	8.1
Number of long range restraints per residue ¹	2.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.3	0.2
0.2-0.5 (Medium)	0.0	0.22
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1~Å are not included in the statistics.

Dordensinda dom o	Count	% ¹	${f Violated^3}$			Consis	tentl	$\overline{ m Violated}^4$
Restraints type	Count	%0°	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	97	16.9	1	1.0	0.2	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	89	15.5	1	1.1	0.2	0	0.0	0.0
Sidechain-Sidechain	8	1.4	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	245	42.6	9	3.7	1.6	0	0.0	0.0
Backbone-Backbone	101	17.6	5	5.0	0.9	0	0.0	0.0
Backbone-Sidechain	130	22.6	4	3.1	0.7	0	0.0	0.0
Sidechain-Sidechain	14	2.4	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	85	14.8	8	9.4	1.4	0	0.0	0.0
Backbone-Backbone	32	5.6	7	21.9	1.2	0	0.0	0.0
Backbone-Sidechain	43	7.5	1	2.3	0.2	0	0.0	0.0
Sidechain-Sidechain	10	1.7	0	0.0	0.0	0	0.0	0.0
Long range ($ i-j \ge 5$)	148	25.7	2	1.4	0.3	0	0.0	0.0
Backbone-Backbone	23	4.0	1	4.3	0.2	0	0.0	0.0
Backbone-Sidechain	77	13.4	1	1.3	0.2	0	0.0	0.0
Sidechain-Sidechain	48	8.3	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	575	100.0	20	3.5	3.5	0	0.0	0.0
Backbone-Backbone	156	27.1	13	8.3	2.3	0	0.0	0.0
Backbone-Sidechain	339	59.0	7	2.1	1.2	0	0.0	0.0
Sidechain-Sidechain	80	13.9	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1~Å are not included in the statistics.

Model ID	Model ID Nu			f viola	ations	5	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Wiodei 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
1	0	1	1	0	0	2	0.16	0.17	0.02	0.16
2	0	0	1	0	0	1	0.15	0.15	0.0	0.15
3	0	0	1	0	0	1	0.11	0.11	0.0	0.11
4	0	0	1	0	0	1	0.14	0.14	0.0	0.14
5	1	1	1	1	0	4	0.15	0.18	0.03	0.16
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	1	0	0	1	0.17	0.17	0.0	0.17
8	0	1	1	0	0	2	0.12	0.13	0.02	0.12
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	0	1	1	0	0	2	0.14	0.16	0.02	0.14

Continued on next page...




Continued from previous page...

Madal ID		Nun	nber o	f viola	ations	;	Mean (Å)	Mar (Å)	${ m SD}^6 \ (m \AA)$	Modian (Å)
Model ID	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (A)	Max (Å)	$SD^*(A)$	Median (Å)
11	0	1	0	0	0	1	0.18	0.18	0.0	0.18
12	0	1	1	0	0	2	0.12	0.13	0.02	0.12
13	0	0	3	0	0	3	0.14	0.17	0.02	0.12
14	0	1	0	0	0	1	0.18	0.18	0.0	0.18
15	0	1	0	0	0	1	0.15	0.15	0.0	0.15
16	0	0	0	0	0	0	0.0	0.0	0.0	0.0
17	0	1	0	0	0	1	0.13	0.13	0.0	0.13
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	1	0	0	1	0.12	0.12	0.0	0.12
20	0	1	0	0	0	1	0.15	0.15	0.0	0.15
21	0	0	1	0	0	1	0.11	0.11	0.0	0.11
22	0	0	1	0	0	1	0.13	0.13	0.0	0.13
23	0	2	2	1	0	5	0.14	0.18	0.03	0.13
24	0	2	0	0	0	2	0.12	0.14	0.02	0.12
25	0	0	0	0	0	0	0.0	0.0	0.0	0.0
26	0	1	0	0	0	1	0.2	0.2	0.0	0.2
27	0	0	0	0	0	0	0.0	0.0	0.0	0.0
28	0	1	1	0	0	2	0.16	0.17	0.01	0.16
29	0	0	1	0	0	1	0.11	0.11	0.0	0.11
30	0	1	1	0	0	2	0.13	0.16	0.03	0.13
31	0	2	1	0	0	3	0.15	0.22	0.05	0.12

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation



9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 555(IR:96, SQ:236, MR:77, LR:146, IC:0) restraints are not violated in the ensemble.

Nu	ımber	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	$ LR^4 $	$ IC^5 $	Total	Count ⁶	%
1	6	5	2	0	14	1	3.2
0	0	1	0	0	1	2	6.5
0	1	0	0	0	1	3	9.7
0	1	0	0	0	1	4	12.9
0	0	1	0	0	1	5	16.1
0	1	0	0	0	1	6	19.4

Continued on next page...




Continued from previous page...

		$\frac{\mathbf{of vio}}{\mathbf{of vio}}$				Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	0	0	0	0	0	7	22.6
0	0	0	0	0	0	8	25.8
0	0	1	0	0	1	9	29.0
0	0	0	0	0	0	10	32.3
0	0	0	0	0	0	11	35.5
0	0	0	0	0	0	12	38.7
0	0	0	0	0	0	13	41.9
0	0	0	0	0	0	14	45.2
0	0	0	0	0	0	15	48.4
0	0	0	0	0	0	16	51.6
0	0	0	0	0	0	17	54.8
0	0	0	0	0	0	18	58.1
0	0	0	0	0	0	19	61.3
0	0	0	0	0	0	20	64.5
0	0	0	0	0	0	21	67.7
0	0	0	0	0	0	22	71.0
0	0	0	0	0	0	23	74.2
0	0	0	0	0	0	24	77.4
0	0	0	0	0	0	25	80.6
0	0	0	0	0	0	26	83.9
0	0	0	0	0	0	27	87.1
0	0	0	0	0	0	28	90.3
0	0	0	0	0	0	29	93.5
0	0	0	0	0	0	30	96.8
0	0	0	0	0	0	31	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	9	0.13	0.02	0.13
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	6	0.16	0.04	0.14
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	5	0.14	0.02	0.15
(1,289)	1:35:A:CYS:H	1:36:A:MET:HA	4	0.12	0.02	0.11
(1,215)	1:28:A:GLU:HB2	1:29:A:ASP:H	3	0.17	0.01	0.18
(1,405)	1:54:A:CYS:HA	1:57:A:CYS:H	2	0.14	0.02	0.14


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	31	0.22
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	26	0.2
(1,215)	1:28:A:GLU:HB2	1:29:A:ASP:H	11	0.18
(1,215)	1:28:A:GLU:HB2	1:29:A:ASP:H	14	0.18
(1,185)	1:24:A:LEU:H	1:24:A:LEU:HG	5	0.18
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	23	0.18
(1,473)	1:67:A:SER:H	1:69:A:SER:H	1	0.17
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	13	0.17
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	28	0.17
(1,72)	1:16:A:LYS:HA	1:24:A:LEU:HG	5	0.17
(1,37)	1:12:A:ASP:H	1:15:A:GLU:H	7	0.17
(1,405)	1:54:A:CYS:HA	1:57:A:CYS:H	10	0.16
(1,238)	1:30:A:VAL:HB	1:31:A:ARG:H	30	0.16
(1,375)	1:52:A:LYS:H	1:53:A:LYS:HA	15	0.15
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	23	0.15
(1,289)	1:35:A:CYS:H	1:36:A:MET:HA	28	0.15

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,215)	1:28:A:GLU:HB2	1:29:A:ASP:H	20	0.15
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	2	0.15
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	5	0.15
(1,472)	1:66:A:PRO:HA	1:67:A:SER:H	1	0.14
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	24	0.14
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	4	0.14
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	12	0.13
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	17	0.13
(1,416)	1:57:A:CYS:H	1:58:A:ARG:H	23	0.13
(1,206)	1:26:A:GLU:HA	1:28:A:GLU:H	8	0.13
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	22	0.13
(1,368)	1:49:A:ILE:HA	1:51:A:ASN:H	13	0.12
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	31	0.12
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	13	0.12
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	19	0.12
(1,405)	1:54:A:CYS:HA	1:57:A:CYS:H	3	0.11
(1,396)	1:54:A:CYS:H	1:57:A:CYS:H	29	0.11
(1,312)	1:40:A:HIS:HB2	1:43:A:CYS:H	21	0.11
(1,289)	1:35:A:CYS:H	1:36:A:MET:HA	10	0.11
(1,289)	1:35:A:CYS:H	1:36:A:MET:HA	24	0.11
(1,289)	1:35:A:CYS:H	1:36:A:MET:HA	31	0.11
(1,179)	1:23:A:ILE:HB	1:24:A:LEU:H	5	0.11
(1,68)	1:16:A:LYS:HA	1:23:A:ILE:H	23	0.11
(1,21)	1:9:A:THR:H	1:10:A:GLU:H	23	0.11
(1,216)	1:28:A:GLU:HB2	1:29:A:ASP:HA	8	0.1
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	12	0.1
(1,5)	1:3:A:GLN:HA	1:5:A:GLY:H	30	0.1



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

